

By raising the subject of the role of primary care in managing chronic kidney disease, Brady and O'Donoghue may have sparked a timely debate.¹ It would also be good to hear from primary care physicians on this issue.

While they call for primary care to take ownership of the problem, it seems clear that colleagues I have spoken to are mystified and confused. Here is a new disease that apparently one in 10 people have, no longer the rare condition we were brought up with as students, and that clever kidney doctors managed with strange diets, pills, potions, and transfusions (not quite leeches). Little wonder we, as GPs, have been frightened off from believing we had a role, especially as the patients don't complain of anything. So now we have a role, what should it be? I believe it turns out to be rather easier than we imagine.

First, find the patients. We're doing the bloods anyway. They nearly all fall into just four categories.

These are people with diabetes, hypertension, older people, and then people with intrinsic kidney disease. The latter is for the clever kidney doctors and can be found because they have rapidly deteriorating kidney function and/or heavy proteinuria or blood, or a family history of polycystic kidneys.

The rest are for us to manage. We find them by looking for them in our populations with diabetes and hypertension. The older population with ageing kidneys will declare themselves along the way with high blood pressure. If these older people don't have high blood pressure there's nothing for us to do apart from give usual healthy living advice.

Once identified we need to treat their vascular risk factors, especially lifestyle and hypertension. Get their blood pressure to target levels of <140/90 mmHg, and if they have significant proteinuria, use ACE inhibitors or ARBs and aim for <130/80 mmHg. I have a low threshold for adding statins although the jury is still out about how effective they are in more severe renal impairment.² One meta-analysis showed benefit in all cause mortality, CVD mortality, non-fatal CVD events, and a reduction in 24-hour urinary protein

excretion.³ We don't need to worry about bicarbonates and bone disease at the stage we are dealing with, unless something crops up in our routine blood tests.

To keep the critics of primary care at bay, make sure patients records are Read Coded correctly, and the QOF will do the rest in terms of auditing the population.

Primary care is getting there but the mystique of CKD has to be removed. This will come with time and education.

Ivan Benett,

*GPwSI Cardiovascular Disease,
365 Wilbraham Road, Whalley Range,
Manchester, M16 8NG.*

E-mail: ivan.benett@nhs.net

REFERENCES

1. Brady M, O'Donoghue D. The role of primary care in managing chronic kidney disease. *Br J Gen Pract* 2010; **60**(575): 396–397.
2. Benett I. The cardiovascular implications of chronic kidney disease. *Prim Care Cardiovasc J* 2009; special issue: chronic kidney disease: 24–27.
3. Strippoli GF, Navaneethan SD, Johnson DE, *et al.* Effects of statins in patients with chronic kidney disease: meta-analysis and meta-regression of randomised controlled trials. *BMJ* 2008; **336**(7645): 645–651.

DOI: 10.3399/bjgp10X514846

MRSA screening: is it really necessary in primary care?

In a period of economic restrictions within the NHS, the cost of Meticillin Resistant *Staphylococcus aureus* (MRSA) screening for healthy individuals receiving minor surgical interventions is becoming increasingly difficult to justify; with lack of evidence to support the effectiveness of this intervention.

Reduction in healthcare associated infections, especially MRSA, is a government target and hence a priority for primary care trusts (PCTs). All patients undergoing elective surgery should now be screened and de-colonised of MRSA prior to surgery.⁵

MRSA colonisation increases the risk of developing infections (ranging from

superficial to invasive) following surgery,^{2,3} that can be difficult and costly to treat, resulting in prolonged hospital stays for affected in-patients.⁴

In NHS Walsall approximately 3000 patients are currently referred annually for all minor surgery in primary care. We audited the results of MRSA screens from 25 June 2009 to 7 December 2009 for patients undergoing vasectomy/carpel tunnel decompression to assess whether the policy was being adhered to and to make recommendations based on the findings.

The audit found that of the 230 patients screened for MRSA (72 carpel tunnel and 158 vasectomy) only one positive case (nasal swab) was identified from a vasectomy patient.

We calculated the cost of these tests to be approximately £3 each, totalling £690, not taking into account administration, transport, and other related costs.

In an attempt to add to the body of knowledge around screening in primary care we recognise that the financial implications to NHS Walsall are minimal, however, for larger organisations there may be savings if screening activities are reviewed.

The Department of Health reports that 30% and 3% of the general population carry *Staph. aureus* and MRSA, respectively.¹ Although our sample size was small, we found that one patient (0.4%) tested positive for MRSA; this is lower than the predicted value of 3% (approximately seven patients from our population sample).

However, our study population is not representative of the general population as they were offered surgery in primary care, indicating that they are healthier than patients undergoing this intervention in hospital.

Based on our findings we recommend a 'risk-based approach' to MRSA screening in primary care as it does not seem effective to screen relatively well patients with no risk factors or evidence of benefit.

In the current economic climate with financial constraints, it will become increasingly difficult to support