- risks of medications. N Engl J Med 2000; 342(22): 1645-1650
- Glenton C, Paulsen EJ, Oxman AD. Portals to Wonderland: health portals lead to confusing information about the effects of health care. BMC Med Inform Decis Mak 2005; 5(1): 7.
- 9. Lewis M, Orrock P, Myers S. Uncritical reverence in CM reporting: assessing the scientific quality of Australian news media reports. Health Sociol Rev 2010; 19(1): 57-72.
- 10. Scotto di Santolo A. Coronavirus treatment shows first 'encouraging results' since start of deadly pandemic. Express.co.uk 2020; 30 Apr: https://www.express.co.uk/news/uk/1275773/ coronavirus-treatment-ebola-antiviral-drug-Remdesivir-covid-19-latest-news (accessed 29 Mar 2022).
- 11. Cao B, Wang Y, Wen D, et al. A trial of Lopinavir-Ritonavir in adults hospitalized with severe Covid-19. N Engl J Med 2020; 382(19): 1787-1799.
- 12. Australian Associated Press. Melbourne scientists find head lice treatment can kill COVID-19 cells. Mail Online 2020; 4 Apr: https://www. dailymail.co.uk/news/article-8186287/Monash-University-scientists-anti-parasite-Ivermectinkill-COVID-19-cells-two-days.html (accessed 29 Mar 2022).
- 13. Adams RC, Challenger A, Bratton L, et al. Claims of causality in health news: a randomised trial. BMC Med 2019; 17(1): 91.
- 14. Sumner P, Vivian-Griffiths S, Boivin J, et al. Exaggerations and caveats in press releases and health-related science news. PLoS One 2016; 11(12): e0168217.
- 15. Bott L, Bratton L, Diaconu B, et al. Caveats in science-based news stories communicate caution without lowering interest. J Exp Psychol Appl 2019; 25(4): 517-542.

DOI: https://doi.org/10.3399/bjgp22X719261

GPs' understanding of the wider workforce in primary care

The workforce crisis within general practice has been well documented. One response to this by NHS England has been national funding for a wider multidisciplinary team with a forecast of 26 000 additional staff by 2024. However, the impact of a wider workforce on patient care and satisfaction is not clear. Gibson et al (2022) showed that having more GPs in the team was associated with an increase in patient satisfaction, but this was not true for increasing staff in other

A wider workforce has so much to offer but GPs cannot promote the roles of other colleagues if we do not understand their skill sets and the success of diversifying the

primary care workforce depends on good leadership.

We explored GPs understanding of the wider workforce. 21 newly qualified GPs on a fellowship programme were asked to rate their understanding of 16 roles and summarise their views on the key responsibilities for GPs. All had been working in primary care for at least 18 months. GPs' understanding of other roles was highly variable with reasonable understanding of some (for example, advanced nurse practitioners, first contact physiotherapists) but poor for others (for example, physician associates, pharmacy technicians, health and wellbeing coaches, care coordinators).

GPs perceived their key responsibilities to be leadership, giving advice, coordinating patient care, holistic management, and managing complexity. The GPs valued listening to short talks from colleagues with other roles and, following an afternoon workshop, felt more equipped to work in wide multidisciplinary teams.

GPs were enthusiastic diversification of the workforce but they cannot successfully lead teams they do not understand. Patient care and satisfaction will remain overly dependent on GPs until leaders are appropriately utilising the skill mix within their teams.² Further education is urgently needed to upskill the workforce so that they understand each others roles, and patients are directed to the right practitioner, at the right time.

Emer Forde.

GP Programme Director, HEE Thames Valley and Wessex Primary Care School. Email: Emer.Forde@hee.nhs.uk

Katie Collins,

GP Programme Director, HEE Thames Valley and Wessex Primary Care School; Clinical Lead, Hampshire and Isle of Wight New to Practice GP Fellowship Programme.

REFERENCES

- 1. Gibson J, Francetic I, Spooner S, et al. Associations of primary care workforce composition with population, professional and system outcomes Br J Gen Pract 2022; DOI: https://doi.org/10.3399/ BJGP.2021.0593.
- 2. Baird B, Lamming L, Beech J, et al. Integrating additional roles into primary care networks. 2022. https://www.kingsfund.org.uk/publications/ integrating-additional-roles-into-primary-carenetworks (accessed 6 Apr 2022).

DOI: https://doi.org/10.3399/bjgp22X719273

Verschlimmbesserung

I agree with David Zigmond that government has been 'trying to fix things but making them worse'. From the 1990s government promoted a consumer model. However, when patients took on the role of consumers, their expectations and demands could be impossible to fulfil, because of GPs' lack of time and funds. I found that the consequent anger of patients could be demoralising. Also, in those years, continuity of care became fragmented so that many patients did not know the GP and so had not formed a relationship of mutual understanding and trust. I felt that relationships with patients had changed from a solid growth in trust over time to something more fragile, ephemeral, and transactional, which was worse for both patients and GPs. If the government could stop making promises about what can never be, and educate and encourage patients to be realistic in their expectations, maybe GPs could lead happier lives and want to stay being GPs.

Hilary A Lavender, Retired GP. Email: hilary.lavender@gmail.com

REFERENCE

1. Zigmond D. More funding, training, and recruitment? Or just more Verschlimmbesserung? Br J Gen Pract 2022; DOI: https://doi.org/10.3399/ bjgp22X718697.

DOI: https://doi.org/10.3399/bjgp22X719465

Correction

In the Clinical Practice article: Nicholas R Jones and Thomas Round, Venous thromboembolism management and the new NICE guidance: what the busy GP needs to know. Br J Gen Pract 2021; DOI: https://doi.org/10.3399/bjgp21X716765. The duration of treatment had been incorrectly cited and should read: 'NICE recommend a minimum of 3 months of anticoagulation for a confirmed VTE, whether provoked or not. An unprovoked VTE may require 6 months of anticoagulation or longer, depending on the risks of recurrence.' The online version has been corrected

DOI: https://doi.org/10.3399/bjgp22X719285

Editorial note

In the April 2022 print Issue the DOIs were incorrect in the Life & Times articles on pages 169–179; the correct DOIs appear in the online version and in the electronic flipbook.

DOI: https://doi.org/10.3399/bjgp22X719297