

perusal of medicine bottles and repeat prescription cards. In only 2% of cases was there any difficulty and even in these cases it was possible to provide adequate emergency care based on clinical findings at the time. From a practical point of view therefore, general practitioners and deputies alike can manage perfectly well without seeing the records in approx 98% of emergency calls.

On the subject of continuity of care I think it is important to point out that the provision of continuity in out of hours care is not the prerogative of the group practice. Some of us who work as full-time deputies can claim continuity ratings roughly comparable. My own figures might be of interest. A study of 300 consecutive calls showed that I had recollection of previously visiting the homes of no less than 36% patients seen. Were I to repeat this study now that I have been nearly nine years in my present post, then I would anticipate a figure in excess of 40%.

Continuity is simply a reflection of time in post and the number of doctors providing the service. Obviously lower ratings would be recorded by those services employing large numbers of casual deputies, just as would be the case where calls were seen by trainees, new partners, locums and those from other practices working a large rota. There really is no difference.

P M JOHNSON

Tregonce Cottage
St Issey
Wadebridge
Cornwall

Surveillance of the over 75s

Sir,
Paul Wallace's editorial on surveillance of over 75 year olds (*July Journal*, p.267) was most discouraging as it was based entirely on theory and selected references. Doctors are not well served by the current negative thinking about reorganization, as this leads only to further discouragement.

I was glad he considers the eight parameters for surveillance, which I put forward to the RCGP in January 1988, to be appropriate. However, he then erects a number of barriers based upon theory, and not experience. Let us consider first the problem of intrusion into a patient's privacy. Such a phrase implies a barren doctor-patient relationship. My own experience of assessing over 70 year olds in my practice indicated that the older the patient the more likely it was that my visit would be welcomed. With the 30% of patients who lived alone, the difficulty was in ensuring that the visit had a medical as well as a social content. Wilkin and

Williams¹ excellent study of 17 771 consultations with patients aged over 65 years undertaken by 201 general practitioners showed that the mean time spent with each patient per year was 25 minutes divided between about five consultations — possibly not long enough to deal with complex medicosocial problems. My own figures showed that before my screening programme there were 6.6 consultations per patient per year in the 70 years and over age group. This figure fell to 3.8 in subsequent years so not only was much time saved, but the frustration of random symptom-oriented visits was avoided.

Dr Wallace quotes data from specialists which suggest that the full assessment would take 60–90 minutes. I found that, with practice, my examination took 20 minutes per patient on average, and was fully structured to geriatric anatomy and pathology. A standardized recording form is helpful both as an *aide-memoire* and for research — my record had space for identification data by the receptionist, the doctor's examination, the social worker's or rehabilitation officer's home assessment, and the nurse's record of urine results, blood tests and blood pressure.

Dr Wallace considers it unlikely that any other member of the existing primary health care team will have time to take on the additional workload. In my practice we were perhaps fortunate in that senior nursing officers knew of our programme, and appointed appropriate personnel. In addition, time devoted to home confinements in the past became available as this work became hospital oriented. The district nurse and I often visited together to plan future management with patients, having found case conferences to be a time-wasting autocracy without the patient's presence. Such joint visiting still seems to be a rare practice.

So far as materials are concerned, I have made my record system available for many years. However, the RCGP has now produced a screening record card for the elderly.

When Dr Wallace suggests the need for training in the appropriate use of instruments and record forms I assume that he does not mean for general practitioners but for 'link workers'. However, the use of link workers would once more push the elderly to the back of the queue. The differences in doctors' perception of their role, result in a wide variation in the pattern of care provided for older patients. Many doctors still recoil from the social dimension needed in geriatric practice. Without this, however, practice would be very dull.

M K THOMPSON

28 Steep Hill
Stanhope Road
Croydon CR0 5QS

Reference

1. Wilkin D, William EI. Patterns of care for the elderly in general practice. *J R Coll Gen Pract* 1986; 36: 567-570.

Involvement of clergy in patient care

Sir,

It was refreshing to read the report by Ward Jones (*July Journal*, p.280) concerning the attitudes of general practitioners to the involvement of the clergy in patient care. Although the paper is a welcome contribution to this neglected area of study, the method used is somewhat unsatisfactory.

The principal difficulty is the use of the word religious without any definition being offered. The use of a letter from the Bishop of Bristol's advisory group in the study implies that the study was cast within a Christian context. However, given that the survey was located mainly in the city of Bristol, one can safely assume that many of the general practitioners and the patients they treat would belong to other religious groups. This factor, we believe, would have affected both the response rate and the answers received. Although the author may have wished to set the study in a Christian context, lack of data on the beliefs of the doctors themselves adds to the confusion.

Again, little distinction was drawn between religious practice and spiritual belief. This undervalues a spiritual belief whereby a person may not adhere to any particular religious practice but, nevertheless, may search for existential meaning within the experience of illness. This is not a minor quibble; some form of spiritual belief is much more common in the population than religious practice. It is a pity that no attempt was made to determine whether the general practitioners understood this distinction and how it might have related to their willingness to refer patients to the clergy.

MICHAEL KING

PETER SPECK

Academic Department of Psychiatry
Royal Free Hospital
Pond Street
London NW3 2QG

Budget holding

Sir,

Not being a member of the RCGP I obviously have little to lose from Dr Sykes' recommendations (*April Journal* p.170). However Dr Sykes has obviously not considered the future of general practice in