explanation was available in at least two languages), and the mobile ones were able to correctly interpret untoward signs and help in the care of the less mobile.

I understand that neither staff nor patients were expected to speak outside of the hospital regarding the medical conditions of others, and perhaps therein lies the difference. We might not need confidentiality if we could trust everyone to mind their own business.

David Church

GP, Machynlleth.

E-mail: David.Church@gp-96014.wales.nhs.uk

REFERENCES

 Scott K, Dyas JV, Middlemass JB, Siriwardena AN. Confidentiality in the waiting room: an observational study in general practice. Br J Gen Pract 2007; 57: 490–493

Depression as chronic disease

Whether they intend it or not, Walters and Tylee's¹ argument that depression needs a chronic disease model (CDM), conjures up a vision of another contractual recall of patients in order to go through an exercise in superficial questioning and box ticking.

What Chew Graham et al,² and others cited by the authors, studied is better thought of in the UK as a collaborative care model, in which intensive (5–10 sessions) personal care is offered over months (but not the years that the CDM implies). Such successful models typically offer choices to patients of drug and non-drug treatments, such as problem solving which are at odds with the simple biomedical conceptualisation of depression as a brain disorder, but harder to deliver with limited resources.

Walters and Tylee point out the failure of current methods of treatment for depression, yet imply that more or greater intensity of the same is required. Given the relatively poor response to antidepressants over placebo, it is difficult to see how more (medical management) could be better in the context of primary care defined depressive disorders.

Depression is not the same as diabetes or asthma, in terms of its daily impact and the personal and social implications of the diagnosis. One of us has demonstrated the moral dilemma facing women in accepting help for depression, and in particular shown that in order to be acceptable, such interventions needed to be seen as short term and temporary.³

Patients with difficult lives meeting current conceptualisations of depression may well benefit from longitudinal care, but as Heath points out, human continuity easily becomes lost when medicine adopts disease based management.⁴ Such a dehumanising approach is in direct opposition to the approach expressed in Chew Graham's study: to 're-humanise' people with depression.

Chris Burton

Senior Research Fellow, University of Edinburgh, Community Health Sciences: General Practice Section. E-mail: chris.burton@ed.ac.uk

Margaret Maxwell

Senior Research Fellow

REFERENCES

- Tylee A, Walters P. We need a chronic disease management model for depression in primary care. Br J Gen Pract 2007; 57: 348–350.
- Chew-Graham CA, Lovell K, Roberts C, et al. A randomised controlled trial to test the feasibility of a collaborative care model for the management of depression in older people. Br J Gen Pract 2007; 57: 364–370.
- Maxwell M. Women's and doctors' accounts of their experiences of depression in primary care: the influence of social and moral reasoning on patients' and doctors' decisions. Chronic Illn 2005; 1(1): 61–71.
- Heath I, Nessa J. Objectification of physicians and loss of therapeutic power. Lancet 2007; 369(9565): 886–888.

Handshakes and dubious editing

You don't seem to have a 'quick response' section on your website, which is a bit of a shame as, although a rare contributor, I did feel the need of such a facility on reading the letter from Gary Parkes² and your subsequent Editorial comment. Don't you think there is a bit of a danger of taking yourselves FAR too seriously? In more blunt Yorkshire terms, you all seem to be in danger of disappearing up your own backsides.

I tolerate the *BJGP*, despite its overwhelming greyness, although I often wonder why. That letter from Dr Jenkins¹

was an unusual shaft of light illuminating the gloom, and making more sense than the most of the rest of the Journal put together. The perception seemed valid to me, representing one of those rewarding aspects of general practice that still happen occasionally even after almost 30 years, and worthy of comment. Whimsical perhaps, but nevertheless appropriate for some light-hearted (but never-the-less valuable) research.

I think that both Dr Parkes' letter and your rather lily-livered response could be actually quite hurtful to Dr Jenkins, if he makes the mistake of taking either seriously. A bit more real general practice such as humanity and humour, and less of this 'informed consent' and 'ethics committee' nonsense would not go amiss. To use words like 'fraud', 'deceive', 'insulting' and 'arrogant' is way over the top. It does make me wonder how many handshakes Dr Parkes gets, or whether he is just content to go home each night with a general feeling of self-satisfied smugness for putting another colleague (or even patient) well and truly in their place ...

Jim Newmark

GP for Refugees, Asylum Seekers, and Homeless, Bevan House, 152 Sunbridge Road, Bradford, BD1 2LT. E-mail: Newmark@jabt.demon.co.uk

REFERENCES

- Jenkins M. Authors response. Br J Gen Pract 2007; 57: 500.
- Parkes G. Handshakes and spoof publications. Br J Gen Pract 2007: 57: 411.

The *BJGP*'s discussion forum is available for immediate comment on the *BJGP* or other relevant topics at www.rcgp.org.uk/bjgp

Correction

In the letter by Campbell M and Freeman JV. Survival statistics. *Br J Gen Pract* 2007; **57:** 410. The author JV Freeman was incorrectly named as John. The author's name is Jenny V Freeman. In the article Bartholomeeusen S, Vandenbrouke J, Truyers C, Buntinx F. Time trends in the incidence of peptic ulcers and oesophagitis between 1994 and 2003. *Br J Gen Pract* 2007; **57:** 497–499. The author C Truyers was incorrectly named as Carl. The author's name is Carla Truyers.

The corrected versions are available online at www. rcqp.org.uk/bjqp/