pharmacists that will be beneficial for all. We are quite sure that if the new system is implemented and the pharmacist is ready to be moulded into a new shape, practical application of these ideas can be achieved even in the least developed areas of Pakistan. It's better to try than to sit and wait for a miracle to happen.

Fahad Saleem,

B-Pharm, MPhil, MBA(HRM), PhD candidate, Discipline of Social and Administrative Pharmacy, School of Pharmaceutical Sciences, Universiti Sains Malaysia. E-mail: fahaduob@gmail.com

Mohamed Azmi Ahmad Hassali,

Senior Lecturer/HOD, Discipline of Social and Administrative Pharmacy, School of Pharmaceutical Sciences, Universiti Sains Malaysia.

Asrul Akmal Shafie,

Senior Lecturer, Discipline of Social and Administrative Pharmacy, School of Pharmaceutical Sciences, Universiti Sains Malaysia.

REFERENCE

1. Saleem F, Hassali AA, Shafie AA. Hypertension in Pakistan: time to take some serious action. *Br J Gen Pract* 2010; **60**(575): 449–550.

DOI: 10.3399/bjgp10X515151

Access to computed tomography

Thomas *et al*'s study into primary care access to computed tomography (CT) scanning¹ (that found that the provision of open access CT for chronic headache reduced referrals to neurology by 88% in the scanned group) is a helpful addition to a sparsely researched area. Its premise and conclusion, however, ought not to be accepted without further consideration. The premise, that it is desirable to avoid referral to neurology, seems questionable as the aim must surely be to reach a diagnosis leading to treatment and, therefore, symptomatic improvement. A scan rarely did that. Of

the 30 patients that had scans and were also referred to neurologists, 13 (43%) had a consultation that lead to the diagnosis of a treatable condition, mainly migraine.

A scan may have stopped 88% of referrals to neurology but did the patients feel better? CT in patients without clinical indicators may have provided short-term relief for doctor and patient, but in the absence of definite neurological indications the study demonstrates the risk: 10.2% had incidental abnormalities and in 5% the scan triggered the referral. Greater CT availability would surely lead to greater usage, more incidental abnormalities, and more referrals; while the average GP requested fewer than two CT scans, one GP requested seven.

GPs may refer fewer chronic headaches if open access CT is available. However, our patients would benefit more from an informed and skilled consultation (whether in primary or secondary care) leading to diagnosis and treatment, than from a scan leading to incidental findings or, more commonly, nothing.

Nick Hall,

Department of Neurology, Gloucester Royal Hospital, Gloucester, GL1 3NN. E-mail: longhallage@googlemail.com

REFERENCE

1. Thomas R, Cook A, Main G, *et al*. Primary care access to computed tomography for chronic headache. *Br J Gen Pract* 2010; **60**(575): 426–429.

DOI: 10.3399/bjgp10X515160

Detection of congenital heart disease in mid-Essex

Congenital heart defects account for 3% of all infant deaths, with 18–25% of affected infants dying in the first year of life. Early (pre-symptomatic) identification helps improve outcome.¹ Routine prenatal anomaly scans detect less than half of

cardiac defects,^{2,3} hence detection after birth remains important. Only families with a high risk of congenital heart defects are offered foetal echocardiography.

The UK National Screening Committee's newborn screening policy comprises clinical examination at birth and at 6–8 weeks, with specific cardiac investigations for high-risk children.⁴

We conducted a 3-year retrospective study in mid-Essex evaluating the detection of congenital heart defects, during postnatal and 6–8 week examinations from September 2004 to August 2007 on 55 infants born in mid-Essex with a confirmed diagnosis of congenital heart defects. Twenty infants identified either antenatally or diagnosed while admitted in the neonatal unit for other indications or detected by active case finding because comorbid conditions were excluded.

Results showed an incidence of 5.5/1000 live-births/year. Postnatal examination detected sixteen (45%) and GPs detected 13 infants (37%) during the 6–8 week check. Six infants with significant congenital heart defects were missed on screening and presented later with symptoms. Cyanotic lesions accounted for just under a quarter of all cases (23.6%). The most common acyanotic lesion was ventricular septal defect (29%). Six infants with significant cardiac lesions were missed on screening and presented at varying ages from

9 days to 7 months of whom four presented with cyanosis. Although all these infants survived, the increased morbidity, use of emergency care resources, and parental stress could be avoided by early detection.

Newborn physical examination is increasingly done in the first 24–48 hours of life by doctors in training and midwives. A congenital heart defect is likely to be missed because signs and symptoms are uncommon in the first few days, infants may look well initially, cyanosis is difficult to detect, and femoral pulses may be felt even in left ventricular obstructive lesions as a result of shunting.