

All letters are subject to editing and may be shortened. Letters should be sent to the BJGP office by e-mail in the first instance, addressed to journal@rcgp.org.uk (please include your postal address). We regret that we cannot notify authors regarding publication. Letters not published in the Journal will be posted online on our Discussion Forum. For instructions please visit: <http://www.rcgp.org.uk/bjgp-discuss>

Editor's choice

The problem with the Liverpool Care Pathway is that someone felt the need to give it a name. Once it had a name it also developed boxes that needed ticking. Everything really went downhill from there. In nearly 30 years of general practice I've looked after many dying patients, but each individual's needs are different. Perhaps it's because I haven't given what I do a name, that I haven't stopped doing it and am going to continue until I retire, working in the same way. Of course, because there are no boxes, there are no QOF points to it; but, I'm rather pleased about that.

JA Glasspool,

*Victor Street Surgery, Victor Street,
Shirley, Southampton, SO15 5SY.*

DOI: 10.3399/bjgp13X675269

Proceed with caution

Beales and Tulloch's arguments about anticipatory care of older patients¹ represent the triumph of hope over experience. Anticipatory care for older people in the community has not yet been shown to be clinically or cost-effective in a thorough and less selective overview of the literature.² Trials of anticipatory care for older people in US, UK, and Denmark up to 1990 showed a rise in patients' morale, increased referrals to all agencies, reduced duration of in-patient stay (sometimes), increased in-patient rates (mostly respite care), reduction in mortality in some trials, but no improvement in functional ability and an increase in GP workload unless alternative services were provided.³

Evidence for the benefits of anticipatory care remains scarce. The UK MRC trial showed little or no benefits for quality of life or health outcomes for older people receiving comprehensive assessment.⁴ A systematic review of 15 trials of preventive home visits for older people showed no clear evidence of benefit⁵ while the ProAge trial

yielded no change in health-risk behaviours in older people.⁶ Case management has not reduced hospital admission rates for frail older people and may even cause disruption of established nursing teams and services.

There are signs that effective interventions are being developed but effect sizes in positive trials are often small and may not remain when interventions are transferred to routine practice. GPs should be cautious about committing time and resources to forms of anticipatory care for older patients that are plausible but untested.

Steve Iliffe,

*Professor of Primary Care for Older
People, University College London.*

E-mail: s.iliffe@ucl.ac.uk

REFERENCES

1. Beales D, Tulloch A. Community care of older people: a cause for concern. *Br J Gen Pract* 2013; **63**: 549–550.
2. Scottish Collaboration for Public Health Research and Policy. Promoting health & wellbeing in later life: interventions in primary care and community settings. <https://www.scphrp.ac.uk/node/198> (accessed 5 Nov 2013).
3. Stuck AE, Siu AL, Wieland GD, *et al*. Effects of a comprehensive geriatric assessment on survival, residence and function; a meta-analysis of controlled trials. *Lancet* 1993; **342**: 1032–1036.
4. Fletcher AE, Price GM, Ng ES, *et al*. Population-based multidimensional assessment of older people in UK general practice: a cluster-randomised factorial trial. *Lancet* 2004; **364**: 1667–1677.
5. van Haastregt JC, Diederiks JP, van Rossum E, *et al*. Effects of preventive home visits to elderly people living in the community: systematic review. *BMJ* 2000; **320**: 754–758.
6. Harari D, Iliffe S, Kharicha K, *et al*. Promotion of health in older people: a randomised controlled trial of health risk appraisal in British general practice. *Age Ageing* 2008; **37**: 565–571.

DOI: 10.3399/bjgp13X675278

The right to die peacefully

The editorial¹ and accompanying article² in the October edition of the *BJGP* highlighted the problems of advance care planning in older people. We detail below the tragic consequences of failure to have these conversations.

A review of case notes of patients registered with a local CCG, who were over 75 years of age and died after spending no more than 1 night in hospital between 1 January 2013 and 31 March 2013, showed that there were 31 such deaths. Of these, eight came from nursing or residential homes and five of these patients were recorded as being unresponsive or had a GCS of 3 when first seen by the ambulance crew. At least six of the patients would have met the Gold Standards Framework prognostic indicators criteria for being on the palliative care register, and in two the family requested admission or resuscitation in case of collapse.

At least four of these cases were pre-alerted to hospital and taken directly into the resuscitation area for multiple investigations and treatments: frail older patients, clearly nearing the end of life, precipitated into hospital where staff feel an obligation to try to preserve life. Most of these patients do not have the mental capacity to understand what is going on around them, and probably find the interventions extremely distressing. The whole process serves only to cause unnecessary suffering.

In some cases the family were not prepared for the patient's demise, and in most the care institutions were not confident in the management of patients nearing the end of life. The ambulance services are put under considerable pressure and without clarity from the carers will understandably default to an active resuscitation mode.

It is a challenge to primary care to champion the rights of older people to die peacefully.

Chris Gunstone,

E-mail: chris.gunstone@nhs.net

Liz Waddy,

*GP and Clinical Lead for End of Life Care
East Staffordshire CCG.*

REFERENCES

1. Eynon T, Lakhani MK, Baker R. Never mind the right time: advance care planning with frail older people. *Br J Gen Pract* 2013; **63**: 511–512.
2. Sharp T, Moran E, Kuhn I, Barclay S. Do the elderly have a voice? Advance care planning discussions with frail and older individuals: a systematic literature review and narrative synthesis. *Br J Gen Pract* 2013; DOI: 10.3399/bjgp13X673667.