



Relieving the bronchial tree in asthma

'ASMAC' TABLETS are formularized to provide symptomatic relief of the bronchial tree both during actual dyspnoeic attacks of bronchial asthma, and during remissions.

'Asmac' Tablets combine in a single prescription 'official' drugs recognized for their reliability to effect mental sedation, decongestion, expectoration and bronchodilatation. *

Packs and Cost to Pharmacists:

Standard Tube of 20: 3/-

Dispensing Bottles:

100, 12/-; 500, 52/6; 1000, 102/6

A. WANDER LIMITED

42 Upper Grosvenor Street,
Grosvenor Square, London W.1.

M. 372

Asmac

WANDER

Formula (each Tablet)

Allobarbitone B.P.C.	0.03 g. (0.46 grain)
Liquid extract of Ipecacuanha B.P.	0.02 ml. (0.34 minim)
Ephedrine Hydrochloride B.P.	0.015 g. (0.23 grain)
Caffeine B.P.	0.10 g. (1.54 grains)
Theophylline with Ethylenediamine B.P.	0.15 g. (2.31 grains)

P1, S1, S4. Permissible on N.H.S. scripts.

The Climacteric

"... a mixture of the male and female hormones was more efficacious than oestrogen alone."

Brit. Med. Journ. 1953 (July 25),
2.214

The association of methyltestosterone and ethinyl- α -estradiol in Mepilin produces a more complete response in the treatment of menopausal disorders than can be obtained by the use of α -oestrogens alone. An increased feeling of confidence and well-being is produced which is both mental and physical.

The presence of methyltestosterone enables a reduction in α -oestrogen dosage to be made; thus undesirable side effects such as breast turgidity and pelvic congestion are avoided and the risk of withdrawal bleeding is reduced.

'MEPILIN'

TRADE MARK

TABLETS & ELIXIR

Each teaspoonful (4ml.) of Mepilin Elixir and each Mepilin Tablet contains ethinyl- α -estradiol 0.01 mg. and methyltestosterone 3 mg.

'MEPILIN' TABLETS. Bottle of 25 at 4/-, 100 at 12/-
'MEPILIN' ELIXIR. Bottle of 4 fl. oz. at 6/- and
500 ml. at 20/6 *Basic N.H.S. prices.*

DOSAGE: Menopause and geriatric conditions: average cases—
3 tablets or 3 teaspoonfuls daily. Premenstrual tension and
dysmenorrhoea—2 tablets or 2 teaspoonfuls daily from 10th to
22nd day of the menstrual cycle.

Literature and specimen packings are available on request

THE BRITISH DRUG HOUSES LTD.
(Medical Department) LONDON N. 1

MEP/1/541