



Specially designed Insurance Policies from the Medical Professions' own Insurance Society

MEDICAL SICKNESS ANNUITY & LIFE ASSURANCE SOCIETY LIMITED 7-10 CHANDOS STREET, CAVENDISH SQUARE, LONDON, W.1. Tel: MUSeum 1686 '.... the use of Pentovis orally was effective in relieving the symptoms of senile vaginitis. In the series (30 patients) there was no evidence that the drug had any unpleasant side effects or produced withdrawal bleeding. Pentovis therefore merits further use, as it seems to be a valuable preparation in the treatment of a distressing and difficult condition.'

Ulster med. J. (1964), 33, 62,

- 'Before the present series, all cases of senile cervicitis or vaginitis or both (in my wards) were treated by vaginal pessaries containing 0.1 mg. oestrone (B. P. C.) daily. No oral treatment was given. There is no doubt that the results with Pentovis are better, in that more effective and quicker improvement is obtained.'
- J. Endocr. (1964) 29, 209.



PENTOVIS

the selective oral oestrogen for

Biopsies before and after treatment repeatedly showed that, while Pentovis has a beneficial effect on vaginal epithelium, endometrial changes do not take place and therefore withdrawal bleeding does not occur.


Pentovis provides oral treatment for

Atrophic vaginitis
Kraurosis vulvae
and related conditions

The side effects common to other oestrogens with multiple sites of action have not been seen with Pentovis. Nausea and gastric disturbances are very

other oestrogens with multiple sites of action have not been seen with Pentovis. Nausea and gastric disturbances are very uncommon. Subjectively, patients usually feel better when taking Pentovis. There are no objections to raising the dose, but this is ordinarily unnecessary.

Dosage: One 0.25 mg. capsule twice daily for fourteen days. This course may be repeated, if necessary. Packages: Pentovis (Quinestradol) is available in bottles of 30 and 250 capsules. The Basic NHS cost of 25 capsules is 8/5d. Pull clinical information available on request. William R Warner & Co Ltd Eastleigh Hampshire England Telephone Eastleigh 3131



In cases of dyspepsia with or without a radiologically proven ulcer Roter will normally provide prompt symptomatic relief and lead to rapid healing of the lesion.

The reputation of Roter is world wide. It is a complete treatment in itself and requires no additional antispasmodics or anticholinergics.



FORMULA: Each Roter tablet contains: Bismuth Subnitrate
Roter (350 mg.), Magnesium Carbonate (400 mg.), Sodium
Bicarbonate (200 mg.) and Cortex Rhamni Frangulae (25 mg.).

PACKINGS OF ROTER TABLETS: Tins of 40 and 120, also dispensing sizes, 360 and 720 (P.T. exempt).

ROTER IN PEPTIC ULCER

A spiral or concentric circles?


clinical studies[†] show pain relief is no illusion with

PONSTAN*

(mefenamic acid, Parke-Davis)

a new effective oral analgesic

†J. Pharmacol. exp. Ther., 139: 172, 1963; Arch. phys. Med., 8: 26, 1965; Practitioner, 194: 817, 1965 Ponstan Kapseals* 250 mg. are available in containers of 12, 100 and 500



Parke, Davis & Co. Hounslow, Middlesex. Tel: Hounslow 2361*Trade Mark

PARKE-DAVIS

TO-DAY'S DRUGS

The most up-to-date book of its kind

"... short, authoritative, and of a high standard."

Lancet (1965), 1, 893.

"... of the greatest value to doctors in general and hospital practice."

Brit. med. J. (1965), 1, 304.

"... well-planned and moderately priced book ..."

Medical News (1965), No. 119, 22.

"... a volume which every physician who considers himself up-to-date will want to have at hand."

Practitioner (1965), 194, 439.

Demy Octavo

Price 30s. net. (postage 2s. 6d. extra)

400 pages



The Publishing Manager, BRITISH MEDICAL JOURNAL, BMA House, Tavistock Square, London, W.C.1 or through any leading bookseller throughout the world

BRANDED CARPETS

WILTONS · AXMINSTERS · TUFTED · ORIENTALS

UP 30% DISCOUNT

- All makes available with full Manufacturers Guarantees.
- No imperfect goods sold ■ Free delivery in the U.K.
- Expert fitting service available most areas.

£100,000 carpets on display at our London Showrooms open 9.30 to 5.30 Mon. to Fri. 9.00 to 12 noon Sats.

DODSON-BULL CARPET CO

Showrooms: 2-4 Little Britain, London E.C.1 (Nr. St. Pauls.)
Offices: 37a Aldersgate St., London, E.C.1 Tel: MON 7171 (10 lines)

LEDERNYCIN 300 Demethylchlortetracycline Lederle fregd. trademark MG TABLETS

and 'Eaton Agent' pneumonia

Enter the Mycoplasma pneumoniae

There was a story about a scientist who discovered a cure for which there was no known disease. Though it was only meant for a laugh, this is not wholly incongruous. Something quite similar has actually happened.

When LEDERMYCIN Demethylchlortetracycline was introduced in 1959, we had full confidence it would prove extremely effective in a host of acute and chronic infections due to a wide range of Gram-positive and Gram-negative organisms. And it did. What we did not anticipate was that it would prove effective in pneumonia due to Mycoplasma pneumoniae.

The reason is simple: the disease itself was not invented – rather, it was not clearly defined – for another four years. Not until 1963 was enough known about it that the causative agent could be assigned a taxonomic designation.

Here are the principal links in the chain of events

1942 'Primary atypical pneumonia' officially established by the Surgeon General of the U.S. Army to distinguish nonbacterial pneumonias of unknown aetiology.

1944 'Eaton agent' isolated from pleural secretions of patients with nonbacterial pneumonia.

1957 'Eaton agent' definitely implicated as the cause of much primary atypical pneumonia.

1961LEDERMYCIN Demethylchlortetracycline shown highly effective in 'Eaton agent' pneumonia.

1962 'Eaton agent' identified as one of the so-called pleuropneumonia-like organisms (PPLO), already assigned the generic name Mycoplasma. 1963 Mycoplasma pneumoniae proposed as species name for 'Eaton agent'.

So, enter Mycoplasma pneumoniae.

And, as a corollary, exit 'Eaton agent'?

Not necessarily. Familiar names
often die hard.

Mycoplasma pneumoniae has ceased to be merely a clinical conversation piece. It is estimated that the newly-named organism may be responsible for as many as half of all hospitalized adult cases of pneumonia. It easily qualifies as a public health problem of some importance.

The recommended dosage for LEDERMYCIN in 'Eaton agent' pneumonia is 900 mg daily given as one 300 mg tablet T.I.D.

Lederle Laboratories



Peritrate SA aids survival

encourages the development of collateral circulation

increases myocardial blood flow

Peritrate SA long-acting vasodilator with twelve-hourly dosage schedule.

Each tablet of Peritrate SA (sustained action) contains 80 mg. pentaerythritol tetranitrate. William R. Warner & Co. Ltd., Eastleigh, Hampshire. Telephone Eastleigh 3131



In convalescence after serious illness **ANABOLEX**

ANABOLEX promotes return of appetite, increased weight and a sense of well-being

Convalescence can be shortened by the administration of ANABOLEX, which stimulates protein metabolism. ANABOLEX is indicated for all patients—adults and children—in negative nitrogen balance. Presentation: Tablets each containing 25 mg. androstanolone in vials of 25, 100 and 250. Dose: The maintenance dose of Anabolex is 2-3 tablets daily and, based on the new reduced NHS price, costs only 1/11d. a day.

LLOYD-HAMOL LIMITED, 36-38 BERKELEY SQ., W.1



SEIGY

Detailed literature describing any Geigy product will be supplied on request.

Geigy Pharmaceutical Company Ltd Manchester 23



in oral steroid therapy

8Mg LEDERCOR'

facilitates the use of new and convenient dosage schedules

once-a-day full daily dose in the morning


once-every-other-day

Full effectiveness maintained Undesirable effects minimised Especially useful in long term therapy Greater convenience for your patients Less chance of missed doses

Lederle Laboratories



a division of Cyanamid of Great Britain Ltd Bush House Aldwych London WC2




Because *Norgesic* goes to work in two ways it can reduce the period of incapacity resulting from acute or chronic painful muscular conditions. *Norgesic* provides the complementary actions of the highly effective skeletal muscle relaxant, orphenadrine citrate, and the anal-

gesic paracetamol, to relieve both local pain and the pain and stiffness associated with muscle spasm. *Norgesic* quickly restores mobility and speeds recovery.

Each tablet contains: orphenadrine citrate 35 mg. and paracetamol 450 mg. Dosage is two tablets three times daily.

RIKER LABORATORIES · LOUGHBOROUGH · ENGLAND

For greater simplicity in oral contraception... Ovulen—the low dosage oral contraceptive—now comes in this 3week wallet



Ovulen (brand of ethynodiol diacetate with mestranol) is the oral contraceptive with a progestin content of only 1 mg. per tablet. The new 3-week wallet means that Ovulen now combines the benefits of low dosage with a simple 3-week administration schedule. This routine can be easily explained to the patient in a few minutes.

SIMPLICITY The patient takes one tablet a day for 3 weeks, waits one week, then starts her next course of tablets. The complete 4-week cycle makes it easier for her to remember her 'pill routine', because her starting day will always be the same day of the week. Ovulen's low progestin content is consistently associated with a very low incidence of side effects.



Patients may not know that Duvadilan is a vasodilator which selectively stimulates the β-receptors which predominate in the smooth muscle cells of the deep blood vessels supplying the arms and legs and in the cerebral arteries

But they do know that Duvadilan
—without causing intense
flushing—makes cold, clammy
extremities warm and supple
again, relieves aching and
restless legs and brightens
mental horizons clouded by
cerebral arteriosclerosis

Indications:

Arteriosclerosis including cerebral arteriosclerosis, aching or restless limbs, cold hands and feet. Raynaud's disease, intermittent claudication, varicose ulcers, gangrene.

Dosage:

Initially 1 or 2 tablets t.d.s. or q.d.s., or 1 ml. (10 mg.) intramuscularly t.d.s. or q.d.s. For maintenance: 1 tablet b.d. or t.d.s.

Packings and Basic N.H.S. Costs:

Tablets containing 10 mg, isoxsuprine hydrochloride:
Bottles of 50: 8s, 0d, Bottles of 250: 33s, 4d,
1 ml, ampoules containing 10 mg, isoxsuprine
hydrochloride: Boxes of 5: 5s, 4d.

The CROOKES Laboratories Limited, Basingstoke, Hampshire.



....NEW 400-DOSE MEDIHALER

When Riker introduced *Medihaler* it was the world's first pressurised aerosol for inhalation therapy... first with a measured dose... first with a dry spray of premicronised particles for optimum effect in bronchospasm. Now Riker announce another first—the 400-dose *Medihaler*. More compact, yet giving one third more treatment from each unit new *Medihaler* is the result of continuous development and of successful use by doctors the world over. With it the cost per dose is reduced 15 per cent, to less than ½d.

Although each unit lasts so much longer, hygiene is improved because with each prescription the patient receives a complete unit. The empty unit, including adapter is thrown away. In addition, the dustcap completely encloses the mouthpiece.


In bronchospasm, *Medihaler* continues to provide an immediacy and degree of relief unsurpassed by any other method—including injection. Now it is even easier and more convenient for the patient to use.

Basic NHS Prices	Standard Pack (complete unit)
Medihaler-iso (Isoprenaline Sulphate B.P. 4 mg./ml.—0.08 mg. per do	se) 14/4
Medihaler-iso Forte (Isoprenaline Sulphate B.P. 20 mg./ml.—0.4 mg. per do	se) 17/-
Medihaler-epi (Adrenaline Acid Tartrate B.P. 14 mg./ml.—0.28 mg. per do	ose) 14/4

Medihaler, Medihaler-iso, Medihaler-epi: registered trade marks

RIKER LABORATORIES · LOUGHBOROUGH · ENGLAND \

The easy way to run an appointments system —in any general practice



In 1959 the Editor of the MEDICAL WORLD set up a committee of experienced family doctors to prepare a handbook on appointments systems in general practice. All the doctors concerned had practical knowledge in this field.

Lloyd-Hamol Ltd. were so impressed by the potential value of the resulting handbook that they offered to provide it, free of charge, to any doctors who wished to read it, together with appointments diaries for running the system, and specimens of appointments cards and announcements to patients.

Two hundred doctors started the scheme that year and it has proved so successful that today the figure has risen to well over 2,000 practitioners, all of whom are now running a full appointments system for their patients. They practise in every type of area and in all parts of the United Kingdom, and in almost every case the benefits derived from the system, both for doctor and the patients, have been very great.

So far, Lloyd-Hamol Ltd. have done little to publicize the service offered. We now feel, however, that advantages of a well-tried appointments system are so numerous that they should be brought to the notice of all doctors who have not to date introduced one.

The Appointments System handbook will be sent to any doctor who wishes to read it, on application to the address below.

LLOYD-HAMOL LTD.


36-38 BERKELEY SQUARE, LONDON, W.1



Hypon is indicated as the analgesic of choice in cases where prolonged administration is required

HYPON®

CALMIC LIMITED
Crewe Hall, Crewe, Cheshire. Tel: 3251



three patients for 'Aldomet'

the most widely prescribed antihypertensive agent

Many of the advantages of 'Aldomet' stem from the fact that it is not an autonomic blocking agent. As the literature confirms:

"... methyldopa does not usually cause troublesome symptomatic postural hypotension, nor does the blood pressure fall unduly on exertion."

Lancet, 1962, i, 763 (14 Apr)

'Side-effects have not so far been troublesome on long-term follow-up.'

Brit.med.J., 1964, 1, 27 (4 Jan)

ALDOMET

METHYLDOPA

a more normal life for the hypertensive patient

Supplied as 250 mg tablets and, for emergency use, as an injection. Detailed information is available to physicians on request.



Merck Sharp & Dohme Ltd, Hoddesdon, Hertfordshire

In peptic ulcer and functional gastro-intestinal disease, NEW sustained-release anticholinergic PORTYN TEMPLETS inhibit gastric secretion, gastro-intestinal spasm or hypermotility


The sustained-release feature of PORTYN* TEMPLETS* provides the advantages of less frequent dosage and fewer and more mild side effects than with the immediate release form. Only one PORTYN TEMPLET twice daily at 12-hour intervals provides sustained inhibition of gastric secretion and gastro-intestinal hypermotility.

Available in bottles of 15 and 100.

PORTYN TEMPLETS



PORTYN TEMPLETS each contain 35 mg. benzilonium bromide, Parke-Davis, (5 mg. for immediate release and 30 mg. for gradual release)



DQV-K

the quality penicillin V

DQV-K is the accepted contraction for **Distaquaine V-K** potassium penicillin V tablets of 60 mg., 125 mg. and 250 mg., syrup in a new fruit flavour and readyprepared suspension both containing 125 mg. per 5 ml.



Dista Products Limited · Speke · Liverpool 24

'Distaquaine' is a registered trade mark PPH365/B2979



gentle nasal decongestant exceptionally prolonged effect rebound hyperaemia is not a problem


Otrivine is xylometazoline hydrochloride

3 forms	N.H.S. Price
Adult Drops 0.1% 1 fl. oz.	3/4
Adult Spray 0.1% & fl. oz.	3/4
Paediatric Drops 0.05% 10ml.	3/0
(with special safety dropper)	

'FRAGRANT CLOUD'

(From Harry Wheatcroft & Sons,Edwalton, Nottingham.)

From early summer until the first hint of the autumn frosts, these coral-red flowers will float a cloud of fragrance throughout your garden. 'Fragrant Cloud' is a rare combination, it has healthy, vigorous growth, well shaped, free-flowering blooms, and, as its name implies, a heady, intoxicating scent. It's no wonder that 'Fragrant Cloud' was the only rose, out of 249 entries, to win the coveted Gold Medal of the National Rose Society.



Enduronyl® controls high blood pressure and eliminates congestion

- Enduronyl® controls * It depletes peripheral catechol amines
 - * It has a specific hypotensive effect by reducing peripheral resistance
 - * It reduces blood volume
 - * It has a central calming effect


Enduronyl[®]

(Methyclothiazide 5 mg and Deserpidine 0.25 mg)

for hypertension—compensated and decompensated

Dosage: 1 to 2 tablets daily





The international standard for a second . . .

. . . is 9.192.631.770 cycles of the hyperfine resonance of the caesium atom. This, the most recent of the international standards, was adopted only in October 1964. Until then, the standard second had been 1/31,556,925,9747th of the year A.D. 1900, an astronomical measurement based on the daily rotation of the earth. The very high accuracy of the clocks physicists can now build

forced the reconsideration of this standard, and, in fact, it was inaccurate because the earth, instead of turning at an absolutely steady speed, was varying, so that the official second was getting longer and longer. The frequency of the radiation emitted by the caesium atom, on the other hand, is so accurate that even month-by-month changes in the earth's speed of rotation could be detected by atomic clocks.



TRANSVASIN is the standard therapy ... for the relief of rheumatic symptoms

Patients suffering from rheumatic lesions obtain rapid relief with Transvasin. The esters of nicotinic, salicylic and p-aminobenzoic acids penetrate the skin and are concentrated on the

point of injury, producing superficial erythema and deep hyperaemia. Transvasin is a valuable adjuvant in the treatment of osteo- and rheumatoid-arthritis.

LLOYD-**HAMOL** Ltd.

LLOYD-HAMOL LTD. 36-38 BERKELEY SQ., LONDON W.1

Ethyl p-aminobenzoate 2% FORMULA Ethyl nicotinate Tetrahydrofurfuryl salicylate n-Hexyl nicotinate 2% Base to 100%

The N.H.S. price of the 30 g, tube is 2/6 and the cost of an average application of cream is under 1d.

a major discovery for rheumatic disorders

INDOCID

'Indocid' is a new, non-steroid anti-inflammatory agent with analgesic and antipyretic properties, chemically unrelated to the corticosteroids, phenylbutazone-like compounds or salicylates.

Widespread clinical experience has shown that 'Indocid' rapidly relieves pain and inflammation, and allows greater mobility. In osteoarthritis, 'Indocid' is often effective where other agents are inadequate; it has given excellent results in osteoarthritis of the hip. In rheumatoid arthritis, objective evidence of effect may be seen as a measurable reduction of joint swelling. 'Indocid' has also proved effective in ankylosing spondylitis, gout, and certain soft-tissue disorders.

'... the first
non-corticosteroid
agent which produced
a predictable and
measurable reduction
in joint-swelling in
most cases of active
rheumatoid arthritis.'

Brit. med. J., 1963, 2, 965 (19 Oct)



'Indocid' is supplied as 25 mg capsules. Detailed information is available to physicians on request.

> Indocid' was discovered and developed by the Merck Sharp & Dohme Research Laboratories and is made in England by



which hypertensive patients are best treated with 'Esbatal'?

All new patients requiring an adrenergic neurone-blocking agent

- "Bethanidine sulphate is an important and potent hypotensive drug, and by its use the regime of many patients has been simplified and side-effects avoided." J. Irish med. Ass., 1965, 56, 9.
- "While expecting the symptoms of adrenergic neurone blockade, we have been pleasantly surprised by the lack of other side-effects, and especially by the absence of diarrhœa." *Lancet*, 1962, ii, 897.

Male hypertensives

'Esbatal' does not cause impotence, but as with other adrenergic neurone-blocking agents, failure of ejaculation may occur. When this causes concern, withdrawal of 'Esbatal' for 24 hours usually reverses this side-effect. Good control of blood pressure is regained rapidly on resuming treatment. (See *Lancet*, 1964, ii, 659.)

Patients in whom other forms of hypotensive therapy cause diarrhœa

".... a useful alternative to established means of therapy. It seldom causes diarrhoa, and the dose is easily manipulated." Lancet, 1964, ii, 659.

When rapid control of blood pressure is essential

- ".... it has the advantage of more rapid action and excretion so that its dose can be increased more rapidly and the blood pressure brought under control more quickly." *Brit. med. J.*, 1964, ii, 865.
- "Bethanidine sulphate is of particular value for the rapid reduction of high blood pressure " J. Irish med. Ass., 1965, 56, 9.

Rapid action— Simpler management— Minimal side-effects

'Esbatal' BETHANIDINE SULPHATE

Available as 10 mg. and 50 mg. scored peach-coloured tablets

Detailed literature available on request



🛌 Burroughs Wellcome & Co., London

(The Wellcome Foundation Ltd.)

Clinically Unexcelled

Clinical use throughout the world has established CHLOROMYCETIN* (Chloramphenicol B.P. Parke-Davis) as an antibiotic of outstanding efficacy in a wide variety of bacterial, viral and rickettsial infections. CHLOROMYCETIN possesses extremely high anti-microbial activity, crosses tissue barriers readily, diffuses widely and rapidly through nearly all body tissues and fluids, and is well tolerated. It is rapidly absorbed and bacterial resistance is minimal. And because of these notable properties, therapy with CHLOROMYCETIN generally results in prompt response and rapid recovery.

Available in Kapseals* of 250 mg and Palmitate Suspension containing the equivalent of 125 mg. Chloromycetin per 4 ml teaspoonful.

*Trade mark CHLOROMYCETIN



NOW, FROM THE DISCOVERERS OF CHLOROTHIAZIDE

a completely new oral diuretic



extends the reach of diuretic therapy:

- 'Edecrin' is chemically and pharmacologically distinct from all other diuretics in current use.
- □ 'Edecrin' has a wider range of effectiveness than the thiazides and related diuretics.
- □ 'Edecrin' may be used in oedema of all grades of severity, whether acute or chronic.
- 'Edecrin' has proved effective in many cases resistant to all other diuretics.
- □ 'Edecrin' has an extremely rapid onset of action faster than parenteral mercurials.
- □ 'Edecrin' produces a more favourable sodium/potassium excretion ratio than the thiazides.

'Edecrin' is supplied as tablets containing 50 mg ethacrynic acid. Detailed information is available to physicians on request.







Lasix tablets each contain 40 mg. frusemide and Lasix ampoules 20 mg. frusemide in 2 ml. Lasix is a registered trade mark.

not only most powerful but also easily controlled

Not only is Lasix powerful, but it is also unusual among diuretics for its breadth of action, which gives great versatility and a high degree of control. The diuretic effect is proportional to the dose given over a uniquely wide range: the greater the dose, the greater the response. And the effect begins half an hour after an oral dose; is normally over in 4 to 5 hours.


This means that whatever the required effect, Lasix dosage can be adjusted to produce it. The normal dose range is 40 to 120 mg. For special cases, perhaps where other diuretics have failed, higher doses can be used. While these are rarely necessary they will require adequate clinical and laboratory control.

HOECHST



SERVICE IN THE CAUSE OF MEDICINE

Hoechst Pharmaceuticals Limited, Portland House, London SW1



The full hypotensive effect of Rautrax may not be apparent for one to three weeks

after the start of treatment. after the start of treatment. This gradual onset of effect gives the patient time to adjust to new, lower blood-pressure levels. There are no

Rautrax: whole-root Rauwolfia serpentina 50 mg.; hydroflumethiazide 50 mg.; potassium chloride 625 mg.

Basic N.H.S. prices: bottles of 25 tablets 9/10; 100 tablets 37/4; 500 tablets 169/4

sudden sharp drops in pressure.

Squibb

Rautrax trademark

E. R. Squibb and Sons Ltd Regal House Twickenham, Middlesex

RT1-463

IMPORTANT ANNOUNCEMENT

ANP 235 becomes lucidril

CHANGE OF NAME. The central neuroregulatory agent, ANP 235 (BP approved name Meclofenoxate), will henceforth be known as LUCIDRIL*, the name by which it is already known in many continental countries.

PRESENTATION. To facilitate administration a new 300 mg. tablet has been introduced. The normal adult daily dose 900-1200 mg. will, therefore, be 3-4 tablets per diem. The 100 mg. tablet is retained for administration to children. All tablets are now foil packed.

INDICATIONS. LUCIDRIL has a wide range of clinical indications. It has a unique pharmacological action whereby it augments the utilization of glucose by neurons and protects them from anoxia. Great benefit will result from the use of LUCIDRIL in confusional states in the elderly, diminished cerebral activity regardless of aetiology including comas, alcoholism, and mental retardation in children. Herpes Zoster is also an important indication.

A brochure and reference papers are available on request. *Trade name.

LLOYD-ANPHAR LTD., 36/38 BERKELEY SQUARE, LONDON, W.1.
AN ASSOCIATED COMPANY OF LLOYD-HAMOL LTD.



in long-term diuretic treatment

CENTYL-K

the only bendrofluazide with potassium supplement for maintenance therapy in all oedematous states

Centyl-K tablets

bendrofluazide 2.5 mg with pot. chlor. 573 mg in bottles of 100 and 500

Centyl injection

5 mg bendrofluazide per ml in vials of 5 ml

Leo Laboratories Limited
Copenhagen London Dublin

to initiate diuresis Centyl injection replaces mercurials in practice





at home...



at work

Benylin Expectorant* is the ethical preparation of choice for the control of cough, and for rapid relief of nasal and bronchial congestion. Benylets* provide all the advantages of Benylin in an easy-to-carry lozenge form. By prescribing both preparations simultaneously, your treatment continues and your patient is afforded

quicker relief of symptoms.

When you prescribe Benylin — prescribe Benylets.

*Trade mark.

BENYLIN & BENYLETS

PARKE-DAVIS HOUNSLOW LONDON TELEPHONE: HOUNSLOW 2361

We offer all Members

20°/. DISCOUNT

on all branded

FURNITURE. DIVANS, CARPETS AND BLANKETS

DIRECT WAREHOUSE COMPANY LAKEVIEW, EDGWARE, **MIDDLESEX** 26

Telephone: STOnegrove 5097 Dept. G/PCT

BUYING

THEN GET DETAILS OF THE EXCELLENT TERMS OFFERED BY YOUR **PROFESSION'S OWN** HIRE-PURCHASE COMPANY

Send the coupon below to:

THE MEDICAL SICKNESS FINANCE CORPORATION **CAVENDISH SQUARE, LONDON, W.1.**

Telephone: MUSeum 1686

- * Minimum deposit
- ★ Maximum period of repayment
- **★** Payments waived on death

A wholly owned subsidiary of The Medical Sickness Annuity & Life Assurance Society Ltd

Name
Address
Occupation

ACCOMMODATION AT COLLEGE HEADQUARTERS

The College of General Practitioners' Club, to which all members and associates belong, provides temporary residential accommodation for them and their families at 14 Princes Gate, London, S.W.7 (Tel. KNIghtsbridge 6262). There is parking space for cars. Children under the age of 12 years cannot be admitted, and no dogs are allowed.

The charges, including breakfast, are as follows:

For single rooms £2 per night
For double rooms £3 10s. 0d. per night

For a flatlet (bed-sitting room for two, bathroom

and dressing room) £5 per night or £30 per week

For a self-contained flat (double bedroom, sitting room, hall, kitchen and bathroom) £35 per week

Members and associates may, subject to approval, hire the reception rooms for social functions and meetings. The charges for these are:

Long Room 25 guineas for each occasion
Common Room 15 guineas for each occasion.

A service charge of 10 per cent is added to all accounts to cover gratuities to domestic staff. Enquiries should be addressed to the Administrative Secretary, 14 Princes Gate, London, S.W.7. Bookings should be made, wherever possible, well in advance.