
Geigy

Tanderil®

Helps arthritic joints to move again



Tanderil*

Major non-hormonal antirheumatic agent

Freedom from pain Reduction in morning stiffness Improvement in grip strength Decrease in joint swelling Increased mobility

Tanderil® is available as tablets containing 100 mg and as suppositories containing 250 mg of 1-phenyl-2-p-hydroxy-phenyl-3, 5-dioxo-4-n-butylpyrazolidine monohydrate.

Detailed literature describing Tanderil or any other Geigy product will be supplied on request.

Geigy (U.K.) Limited Pharmaceuticals Division Macclesfield, Cheshire

Generations of Students have used and trusted . . .

HUTCHISON'S CLINICAL METHODS

By DONALD HUNTER, C.B.E., M.D., D.Sc., F.R.C.P. and R. R. BOMFORD, C.B.E., D.M., F.R.C.P.

For over 70 years successive editions of Hutchison's Clinical Methods packed with hasic information essential to all who wish to practice the art and science of good medicine, has been used by students and practitioners all over the world, as one of their basic clinical textbooks. For this edition the book has undergone a complete and thorough revision, and much new material has been included.

15th Edition

416 bages 12 pages of coloured blates

12 pages of black'& white plates

and

BACTERIOLOGY AND IMMUNOLOGY FOR STUDENTS OF MEDICINE

By F. S. STEWART, M.D., F.R.C.P.I., F.C. Path., M.R.I.A.

Since the last edition appeared six years ago, the explosive growth in the fields of microbiology and immunology has continued. This has necessarily involved a substantial rewriting of this standard textbook formerly entitled Bigger's Handbook of Bacteriology. Over 40 per cent of the text is entirely new, and all chapters have been extensively revised.

9th Edition

600 bages

78 bhotograbhs

3 colour blates

55s.

BAILLIÈRE, TINDALL AND CASSELL LTD.

7 & 8 Henrietta Street, London W.C.2




Their well-being can he assured . . .

Your family's happiness and comfort depend so much on vou! The best way of ensuring that they will be taken care of, if anything happens to you, is by means of Life Assurance.

For the most favourable terms write to:-

Medical Insurance Agency Ltd. Head Office: B.M.A. HOUSE, TAVISTOCK SQUARE, LONDON, W.C.1

Telephone: 01-387 6031 (9 lines)



Tegretol® Anticonvulsant


Geigy

Widens the scope of anti-epileptic therapy

Tegretol is available as tablets each containing 200 mg 5-carbamoyl-5H-dibenz (b,f) azepine.

Detailed literature will be supplied on request

Geigy (U.K.) Limited Pharmaceuticals Division Macclesfield, Cheshire





A UNIQUE ORAL GEL

WITH STRIKING ANALGESIC AND ANTISEPTIC PROPERTIES

Bonjela has been shown in formal American trials to bring rapid relief from pain in 90% of all cases of oral ulceration, herpes labialis, gingivitis and denture sores. By rapid relief is meant marked alleviation of pain within 2-3 minutes.

Bonjela also promotes healing whenever the aetiology is partly or wholly bacterial. Some ulceration originates from endocrinal, viral or fungal causes which may require other forms of treatment.

Even in these cases where a bacterial element is superimposed on the

original infection Bonjela will reduce the severity of the infection and rapid relief from pain will be obtained.

FORMULA: Choline Salicylate 8.7%, Cetalkonium Chloride 0.01%, Alc. (95%) 39.0%, Menthol 0.057%, Glycer. 4.6%.

INDICATIONS: Herpes labialis, gingivitis, oral ulcerations, cheilosis, denture sores, infant teething disorders.

PRESENTATION: Bonjela is available in 10 g. tubes at a basic N.H.S. cost of 2/8½d. per tube.

LLOYDS' PHARMACEUTICALS LTD. 103 Mount Street, London, W.1. An associate of Lloyd-Hamol Ltd.

Literature and samples are available on request






Like radar, LABSTIX reagent strips will scan, give an early warning and immediate identification about the unknown.


Covering a wide range of five tests (for blood, ketones, glucose, protein and pH) LABSTIX reagent strips provide the most comprehensive urine testing procedure, available in convenient form, for detecting signals of disease.

Like radar, to be effective LABSTIX should be used routinely, day-in and day-out. Insure yourself against missing something—before it is too late.

${f LABSTIX}^*$

gives you early routine protection





your patient appreciates this

He appreciates the significant reduction in the number of attacks.

He appreciates release from the restrictions of his condition.

He appreciates Synadrin 60.

References

Prenylamine in Angina Pectoris Canad. Med. Ass. J. 1963, 89, 248.

Problems in the treatment of Angina Pectoris. Med. Klin. 1966, 61, 352.

Double-blind trial with Synadrin 60 in Angina Pectoris. Zschr. Ther. 1965, 2, 82.

The successful application of Synadrin 60 in the treatment of Angina Pectoris. Wien. Med. Wschr. 1965, 115, 1034.

Clinical investigations of the treatment of Coronary Insufficiency with Synadrin 60. *Med. Klin.* 1964, **35**, 1384.

Full information available on request.

Med. Klin. 1964, **35**, 1384.

SYNADRIN 60 is available as tablets containing 60 mg prenylamine.

Synadrin® 60 is available in many other countries as Segontin® 60

Hoechst Pharmaceuticals Limited Hoechst House Kew Bridge Brentford Middx





so many hypertensive patients do well on Aldonet

Methyldopa BP

- 'Aldomet' offers smooth and even control of blood pressure when the dosage has been established,
- 'Aldomet' usually reduces lying as well as standing blood pressure.
- 'Aldomet' causes much less postural hypotension than other powerful antihypertensive agents.
- 'Aldomet' rarely causes exertional hypotension.
- The dosage of 'Aldomet' is simple, flexible and easily established.

Supplied as tablets containing methyldopa BP equivalent to 250 mg anhydrous methyldopa and, for emergency use, as an injection. Detailed information is available to physicians on request.


In a trial comparing three antihypertensive agents, it was shown that,'...in the patients who could take adequate doses methyldopa provided the best control of blood pressure, with the lowest levels of blood pressure when the patient was supine.'

Brit.med.J.,1968,1,129 (20 Jan)



® denotes registered trademark

Merck Sharp & Dohme Limited Hoddesdon, Hertfordshire Telephone, Hoddesdon 67123



STANDARD The weight of a tennis ball

The International Lawn Tennis Federation lays down that a tennis ball shall be more than 2oz. and less than $2\,\%$ oz. in weight. A fine distinction, you may think. But there is more to come. It must have a uniform outer surface. If there are any seams, they must be stitchless. It must be more than $2\frac{1}{2}''$ and less than $2\frac{5}{8}''$ in diameter, have a bound of more than 53'' and less than 58'' when dropped 100'' upon a concrete base and a deformation of more than .265'' and less than .290'' when subjected to a pressure of 18 lb. applied to each end of the diameter.



The N.H.S. price of the 30 g. tube is 2/6d. and the cost of an average application of cream is under 1d. LLOYD-HAMOL LTD., 103 Mount Street, London, W.1

STANDARD

The Transvasin therapy for rheumatic symptoms

Transvasin as the standard treatment for the relief of rheumatic symptoms is becoming more and more widely accepted every day. Patients suffering from rheumatic lesions obtain rapid relief with this well-proven and economical medication. The esters of nicotinic, salicylic and p-aminobenzoic acids penetrate the skin and are concentrated at the point of injury, producing superficial erythema and deep hyperaemia. Transvasin is a valuable adjuvant in the treatment of osteo and rheumatoid arthritis.

CONTAINS Ethyl Nicotinate 2° Hexyl Nicotinate 2° Tetrahydrofurfuryl Salicylate 14° Ethyl Aminobenzoate 2° Base 10 100°



THE ROYAL COLLEGE OF GENERAL PRACTITIONERS

PUBLICATIONS

Copies of the following publications may be obtained from E. & S. Livingstone, Limited, Teviot Place, Edinburgh.

Reports from General Practice

No. 1. Vocational Training	5s.	
No. 3. Additional Payments for Wide Experience and	i	
Notable Service in General Practice	1s.	
No. 4. General Practice in the New Towns of Britain	6s.	
No. 5. Evidence of the College of General Practi-		
tioners to the Royal Commission on Medical		
Education		6d.
No. 7. Education in Psychology and Psychiatry		6d.
No. 8. General Practice in South-west England		6d.
No. 9. Obstetrics in General Practice		6d.
No. 10. The Practice Nurse	7s.	6d.
Symposia		
Hazards of Middle Age	5s.	
Problems of Sex	6s.	
The Art and the Science	7s.	6d.
Accident Management	6s.	
Nutrition in General Practice	7s.	6d.
Arthritis in General Practice	5s.	
Migraine in General Practice	5s.	
Mental Health and the Family Doctor	5s.	
Rehabilitation	5s.	
The Aetiology of Congenital Anomalies	7s.	6d.
The Early Stages of Chronic Bronchitis	10s.	6d.
Preventive Medicine and General Practice	10s.	6d.
The Quality of Medical Care	5s.	
Health in a Changing Environment		6d.
Clinical Problems of Practice		6d.
Anaemia in General Practice	6s.	
The Art of Listening		6d.
Early Diagnosis		6d.
The Age of Discretion	7s.	6d.
Other Publications		
Training for General Practice (2nd Edition)	4 s.	6d.
Epidemic Winter Vomiting	1s.	
On Undergraduate Education	2s.	
Memorandum for the Guidance of Trainers	1s.	
Guidance for Students visiting a General Practitioner	1s.	
Group Practice, Ancillary Help and Government		
Controls	7s.	6d.