



in contrast with other antihypertensive drugs

VASOLASTINE

is free from side-effects.

It is non-toxic.

In addition it resolves atheromatous deposits, normalises cholesterol and phospholipid values, and restores elasticity of arteries and capillaries.

VASOLASTINE

The MODERN CAUSAL ENZYME THERAPY
for hypertension and cardiovascular diseases



Manufactured by Enzypharm C. V. Holland

*Further information and list of publications
gladly sent on request by the Sole U.K. Distributors:*

F.A.I.R. LABORATORIES LTD., TWICKENHAM MIDDLESEX

FORMULA per ampoule:

Enzymes of the lipid
metabolism 30 mg. ≡ 8000 units
Amin-oxidase 20 mg. ≡ 4000 units
Tyrosinase 20 mg. ≡ 4000 units
Stabilizing solvent to 2 ml.
Presentation:
Boxes of 6, 25 and 100 ampoules.



Butazolidin® Alka mantle tablets

A new approach to the
treatment of patients with
a known gastric sensitivity
to antirheumatic agents

Butazolidin® Alka is formulated as a
Mantle tablet incorporating 100 mg
dried aluminium hydroxide gel B.P. and
150 mg magnesium trisilicate B.P. around
a central core of 100 mg Butazolidin®.

Geigy (U.K.) Limited
Pharmaceuticals Division
Macclesfield, Cheshire

In Bronchitis

SEPTRIN

Trade Mark

**Abolishes pathogens, renders
sputum mucoid and dramatically
decreases sputum volume***

New mode of bactericidal action. An advance on antibiotic therapy. Reduces or eliminates problems of bacterial resistance, disturbance of gut flora and other troublesome side-effects. Exceptionally rapid clinical effect in respiratory tract; urinary tract and other infections caused by a wide range of sensitive organisms – Gram-positive and Gram-negative.

'Septrin' tablets

Each contain trimethoprim 80 mg,
sulphamethoxazole 400 mg.

'Septrin' Paediatric tablets

Each contain trimethoprim 20 mg,
sulphamethoxazole 100 mg.



* Proceedings of the V International Congress of Chemotherapy, Supplement, (A 1-5a/3) 107.
Further information is available on request.

Burroughs Wellcome & Co. (The Wellcome Foundation Ltd.) London

In the elderly, the special problems of constipation...



Constipation in the elderly can present special problems of treatment. Weakened abdominal muscles... the dangers inherent in straining... refusal to co-operate... the presence of pain, to which anal fissures or haemorrhoids may contribute... the need for enemas and suppositories... the risk of faecal impaction, with consequent manual removal...

Dorbanex can simplify management of the patient posing these problems by providing comfortable, reliable and trouble-free evacuation. *Dorbanex* produces a soft but well-formed stool without griping or other unpleasant side-effects, reduces the need for enemas and suppositories, helps to prevent faecal impaction, and makes defaecation easier without increasing faecal incontinence.

...have a straightforward solution

Now available in three forms—as capsules and in two strengths of liquid—*Dorbanex* provides in one preparation a balanced treatment for constipation. By softening the stool and stimulating peristalsis, *Dorbanex* prepares the way for pain-free evacuation.


DORBANEX CAPSULES: each contains 1:8-dihydroxyanthraquinone 25 mg., poloxalkol 200 mg. **DORBANEX LIQUID:** each 5 ml. teaspoonful contains 1:8-dihydroxyanthraquinone 25 mg., poloxalkol 200 mg. **DORBANEX FORTE:** each 5 ml. teaspoonful contains 1:8-dihydroxyanthraquinone 75 mg., poloxalkol 1000 mg. **Dosage DORBANEX CAPSULES:** Adults, one or two capsules. Children, one capsule. **DORBANEX LIQUID:** Adults, one or two teaspoonfuls. Children, half to one teaspoonful, as required. Before or after surgery: two to four teaspoonfuls or two to four capsules. **DORBANEX FORTE:** one teaspoonful, or as directed by the physician. Treatment should be taken at bedtime. **Contra-indications** In common with other gastro-intestinal evacuants, *Dorbanex* should not be given when acute, painful conditions of the abdomen are present, or the cause of constipation is suspected to be obstruction of the small or large intestine. **Side-effects** Dihydroxyanthraquinone may cause a temporary harmless, pink or red colouring of the urine and, in prolonged use or high dosage, of the mucosa of the large intestine. Doses larger than two teaspoonfuls may cause slight gastro-intestinal discomfort in a few patients. **Basic NHS Prices** *Dorbanex* Capsules: bottles of 30 capsules 8/8; bottles of 300 capsules 78/-; *Dorbanex* Liquid: bottles of 60 ml. 4/-; bottles of 500 ml. 30/-; *Dorbanex* Forte: bottles of 60 ml. 6/-; bottles of 500 ml. 45/-. Additional information on request.

Dorbanex prepares the way for pain-free, trouble-free evacuation

Riker Laboratories Loughborough Leicestershire



it's
wrong to
assume
that gorillas
show aggression
towards man



Those old, old stories of explorers being carried off by gorillas were not really true! Although the fully grown gorilla is more than a match for 10 men, man is the one animal from whom the gorilla runs on sight.

An assumption proven incorrect.

Similarly, a negative test for protein in urine, by standard methods, contrary to popular belief, does not exclude the possibility of a significant haematuria. LABSTIX reagent strips avoid the need for assumption, they provide information on the presence or absence of urinary glucose, protein, ketones and blood, together with an indication of urinary pH. This test has now become established as a routine in many wards, clinics and surgeries throughout the United Kingdom.

Facts are more reliable than assumptions.

Let LABSTIX provide the information in 30 seconds.

LABSTIX*

for earlier diagnosis and effective follow-up

Further information is available on request.




Ames Company

Division Miles Laboratories, Ltd.
Stoke Court, Stoke Poges, Slough, Bucks.



**trade mark*



“... the safest oral hypoglycaemic drug
for general use, and it should be tried first”¹



“Three-quarters (2,555 patients or 75.4 per cent) of the
entire group of 3,387 patients achieved satisfactory control”²



“Tolbutamide has been remarkably well
tolerated by almost all patients and side-effects
were observed in less than 1 per cent of the cases”²

References:

1. Today's Drugs, *BMJ*, 1963, 1, 521.
2. *Metabolism*, 1966, 15, 957.

Rastinon® First – decisively

Rastinon tablets contain 0.5G tolbutamide B.P. Full information available on request



Hoechst Pharmaceuticals A Division of Hoechst UK Limited
Hoechst House Kew Bridge Brentford Middlesex

why



so many hypertensive patients do well on Aldomet®

Methyldopa BP

- 'Aldomet' offers smooth and even control of blood pressure when the dosage has been established.
- 'Aldomet' usually reduces lying as well as standing blood pressure.
- 'Aldomet' causes much less postural hypotension than other powerful antihypertensive agents.
- 'Aldomet' rarely causes exertional hypotension.
- The dosage of 'Aldomet' is simple, flexible and easily established.

Supplied as tablets containing methyldopa BP equivalent to 250 mg anhydrous methyldopa and, for emergency use, as an injection. Detailed information is available to physicians on request.


In a trial comparing three antihypertensive agents, it was shown that, '... in the patients who could take adequate doses methyldopa provided the best control of blood pressure, with the lowest levels of blood pressure when the patient was supine.'

Brit.med.J., 1968, 1, 129 (20 Jan)



® denotes registered trademark

Merck Sharp & Dohme Limited
Hoddesdon, Hertfordshire
Telephone, Hoddesdon 67123



GOLF

It was the Scots who brought the game of golf to its greatest flowering, and it is the Royal and Ancient Club of St. Andrews which is supreme arbiter of the game today. It is they who lay down the rules, including the one, for instance, that says that the weight of the ball shall be not greater than 1.620 ounces avoirdupois and the size not less than 1.620 inches in diameter.

The NHS price of the 30 g. tube is 2/6d. plus P.T., and the cost of an average application of cream is under 1d. Literature and samples are available on request.

TRANSVASIN

Patients suffering from rheumatic symptoms obtain rapid relief with this well-proven and economical medication. The esters of nicotinic, salicylic and aminobenzoic acids penetrate the skin and are concentrated at the point of injury, producing superficial erythema and deep hyperaemia. Transvasin is a valuable adjuvant in the treatment of osteo- and rheumatoid arthritis.

CONTAINS

Ethyl Nicotinate	2%
Hexyl Nicotinate	2%
Tetrahydrofurfuryl Salicylate	14%
Ethyl Aminobenzoate	2%
Base	to 100%



**The standard treatment
for rheumatic symptoms**

Lloyd-Hamol Ltd., 103 Mount Street, London, W.1

Transvasin is a Trade Mark



A UNIQUE ORAL GEL WITH STRIKING ANALGESIC AND ANTISEPTIC PROPERTIES

Bonjela has been shown in formal American trials to bring rapid relief from pain in 90% of all cases of oral ulceration, herpes labialis, gingivitis and denture sores. By rapid relief is meant marked alleviation of pain within 2-3 minutes.

Bonjela also promotes healing whenever the aetiology is partly or wholly bacterial. Some ulceration originates from endocrinal, viral or fungal causes which may require other forms of treatment.

Even in these cases where a bacterial element is superimposed on the

original infection Bonjela will reduce the severity of the infection and rapid relief from pain will be obtained.

FORMULA: Choline Salicylate 8.7%, Cetalkonium Chloride 0.01%, Alc. (95%) 39.0%, Menthol 0.057%, Glycer 4.6%.

INDICATIONS: Herpes labialis, gingivitis, oral ulcerations, cheilosis denture sores, infant teething disorders.


PRESENTATION: Bonjela is available in 10 g. tubes at a basic N.H.S. cost of 2/8½d. per tube.

LLOYDS' PHARMACEUTICALS LTD.
103 Mount Street, London, W.1.
An associate of Lloyd-Hamol Ltd.

Literature and samples are available on request



**'Good night'
(less pain)**



**'Good morning'
(more mobility)**



12 to 16 hours relief for many arthritic patients

When night pain and morning stiffness are prominent symptoms in arthritis, 'Indocid' Suppositories can be particularly valuable. They are long-acting, often giving relief for 12 to 16 hours. When inserted at bedtime, 'Indocid' Suppositories frequently give a better night's sleep. Morning stiffness is reduced and the patient has a better start to the working day.

'Indocid' is supplied as 100 mg polyethylene glycol suppositories, and also as 25 mg capsules, and a suspension (25 mg/5 ml).
Detailed information is available to physicians on request.
® denotes registered trademark.

Indocid
Indomethacin
Suppositories



Merck Sharp & Dohme Limited
Hoddesdon, Hertfordshire
Telephone, Hoddesdon 67123

THE ROYAL COLLEGE OF GENERAL PRACTITIONERS

Services provided for members

1. **The Geoffrey Evans Library** provides lists of references and a photocopying service.
Librarian: Miss M. Hammond, 14 Princes Gate, London, S.W.7.
2. **Medical recording service and sound library**
Honorary directors: Drs John and Valerie Graves, O.B.E., Kitts Croft, Writtle, Chelmsford, Essex.
3. **Postgraduate Information Diary**
Information officer: Education Committee of Council, 14 Princes Gate, London, S.W.7.
4. **Practice organization.** Library of architects' plans, photographs, descriptions of practice premises and advice on equipment.
Secretary: Practice Organization Committee, Royal College of General Practitioners, 14 Princes Gate, London, S.W.7.
5. **Awards.** A number of Travelling Fellowships and prizes are in the gift of the College. Information is published periodically in the *College Journal*.
6. **Research Register, and Register of Research Projects**
Honorary registrar: Dr R. M. Griffiths, 4 The Dell, Woodford Wells, Essex.
7. **The Records and Research Advisory Service.** Planning and statistical advice and processing for mechanical analysis of survey results or other records. Advice in planning, in contacting like-minded practitioners, advice in method, on any aspect of research work.
Honorary director: Dr D. L. Crombie, 8 Albert Road, Harborne, Birmingham 17.
8. **Epidemic Observation Unit.** Collates observations of practitioners which might otherwise be lost.
Honorary director: Dr G. I. Watson, Corran, Peaslake, nr. Guildford, Surrey.
9. **Grants.** Grants in aid of research are made from the funds of the Research Foundation. Advice as to other sources of finance for research and help with preparation of applications.
Honorary secretary/treasurer: Dr E. V. Kuenssberg, 15 Warriston Crescent, Edinburgh 3 (or to 14 Princes Gate, London, S.W.7).
10. **College Club.** All members of the College are *ipso facto* members of the club at 14 Princes Gate, S.W.7, where bed and breakfast may be obtained at reasonable rates, and where members may meet friends. Members may, subject to approval, hire rooms for functions and meetings.
Administrative secretary: Mrs E. Phillips, Royal College of General Practitioner, 14 Princes Gate, London, S.W.7.
11. **Journal.** Published monthly and circulated to all members and associates. Subscription to non-members, £4 4s. 0d. per annum. (Single copies 10s. 6d.)
Honorary editor: Dr R. M. S. McConaghey, O.B.E. } Prospect House
Editorial manager: Miss I. Scawn } Newcomen Road, Dartmouth

Publications

Copies of the following publications may be obtained from E. & S. Livingstone, Limited, Teviot Place, Edinburgh.

Reports from General Practice		Mental Health and the Family Doctor	
No. 1. Vocational Training	5s.	Rehabilitation	5s.
No. 3. Additional Payments for Wide Experience and Notable Service in General Practice	1s.	The Aetiology of Congenital Anomalies	7s. 6d.
No. 4. General Practice in the New Towns of Britain	6s.	The Early Stages of Chronic Bronchitis	10s. 6d.
No. 5. Evidence of the College of General Practitioners to the Royal Commission on Medical Education	6s. 6d.	Preventive Medicine in General Practice	10s. 6d.
No. 7. Education in Psychology and Psychiatry	7s. 6d.	The Quality of Medical Care	5s.
No. 8. General Practice in South-west England	7s. 6d.	Health in a Changing Environment	7s. 6d.
No. 9. Obstetrics in General Practice	7s. 6d.	Clinical Problems of Practice	8s. 6d.
		Anaemia in General Practice	6s.
		The Art of Listening	7s. 6d.
		Early Diagnosis	8s. 6d.
		The Age of Discretion	7s. 6d.
		The early detection of imported and endemic disease	7s. 6d.

Symposia

Hazards of Middle Age	5s.
Problems of Sex	6s.
The Art and Science	7s. 6d.
Accident Management	6s.
Nutrition in General Practice	7s. 6d.
Arthritis in General Practice	5s.
Migraine in General Practice	5s.

Other Publications

Training for General Practice (2nd Edition)	4s. 6d.
Epidemic Winter Vomiting	1s.
On Undergraduate Education	2s.
Memorandum for the Guidance of Trainers	1s.
Guidance for Students visiting a General Practitioner	1s.
Group Practice, Ancillary Help and Government Controls	7s. 6d.

All your fees paid, plus £790 a year and allowances, while a Medical student.

As a Medical Cadet in the Royal Army Medical Corps, all your tuition fees are paid by the Army. In addition, you are paid £790 a year, and are eligible for allowances.

You can apply for a Medical Cadetship if you have passed your second M.B. or its equivalent, are under 25, and have a place at a medical school in the United Kingdom or Eire. If you are about to take your second M.B., you may still apply; but your acceptance will be conditional on your passing the examination.

If accepted as a Medical Cadet, you are given the probationary rank of 2nd Lieutenant. On qualification, you are promoted Lieutenant, and are paid as such while you fulfil your pre-registration appointment

—either at a civilian hospital of your own choice, or at certain military hospitals. After being fully registered, you are promoted Captain, and take up your duties in the RAMC.

Your obligation is then for five years of service on a Short Service Commission. During those five years, you will have a good chance of travel, make many friends, have ample leave (six weeks' paid leave a year, with a locum provided by the Army).

During that period you can, if you like, apply for a Regular Commission. Or you can leave the Army for civilian practice, the richer for some valuable experience of General Practice—and for a tax-free gratuity of up to £2,600 according to period of cadetship.

For further information about Medical Cadetships, or to arrange a meeting with an RAMC Officer to answer any questions you may have, write to: Colonel E. L. O. Hood MB, Ministry of Defence (AMD2), Dept. 611, Lansdowne House, Berkeley Square, London W1X 6AA

M.3



Fucidin Ointment

is the only topical antibiotic
which has been shown experimentally
to penetrate intact skin
as effectively as the corticosteroids.*

*This scientific fact makes **Fucidin Ointment**
the logical topical treatment for such common
bacterial infections as boils and carbuncles.*

*Fucidin Ointment may be used alone, or, where indicated,
in combination with systemic antibiotics and/or surgery.*

Fucidin Ointment —the logical topical antibiotic

Fucidin Ointment: sodium fusidate B.P. 2%
in a neutral ointment base. Tubes of 10G & 25G.

**The British Derm. Assoc. Meeting. Aberdeen, July 1968.*

Full information is available on request.

and in infected and non-infected dermatoses – FUCIDIN H OINTMENT

Fucidin H Ointment: sodium fusidate B.P. 2% and hydrocortisone
acetate B.P. 1% in a neutral ointment base.



LEO LABORATORIES LIMITED • HAYES • MIDDLESEX Telephone: 573 6224