# Prevent breakthrough hyperacidity Nulacin


When your patient presents with symptoms of gastric hyperacidity, Nulacin therapy provides continuous neutralisation—essential for the prevention and healing of peptic ulcer since most peptic ulcers arise and recur in the presence of free acid.

The Nulacin tablet is prepared from whole milk combined with dextrins and maltose, together with magnesium trisilicate 230 mg., magnesium oxide 130 mg., calcium carbonate 130 mg., magnesium carbonate 30 mg., Ol. Menth. Pip. q.s.



Full information is available on request to





Nellcome Burroughs Wellcome & Co. (The Wellcome Foundation Ltd.) Dartford, Kent



### Butazolidin Alka

The modern way to prescribe an antirheumatic agent

Unique formulation

Proved effectiveness

**Excellent gastric toleration** 

Pract. 203, 1215, 357-360 (1969)

Formulated as a mantle
of two antacids around a core
of Butazolidin® and antacid,
each tablet contains
Butazolidin (Phenylbutazone B.P.)
100 mg. Magnesium Trisilicate
B.P. 150 mg. Dried Aluminium
ydroxide Gel B.P. 100 mg.

Detailed literature describing any Geigg product will be supplied on sequest.

Geigy (U.K.) Limited Pharmaceuticals Division Macclesfield, Cheshire,



Urticaria is often at its worst at night-time. Pro-Actidil\* taken in the evening, brings relief during the night and with its sustained antihistamine effect goes on working during the next morning and afternoon.

The three-layered Pro-Actidil presentation is effective in the treatment of allergic dermatoses especially if associated with urticaria, angioneurotic oedema, seborrhoeic eczema, pruritus, allergic asthma, and vasomotor rhinitis.


Pro-Actidil tablets each contain 10 mg triprolidine hydrochloride. Also available: Actidil\* tablets, each containing 2.5 mg triprolidine hydrochloride (useful for daytime use and when a short duration of action is required), and as Actidil Elixir containing 2 mg in each 5 ml.

Full information is available on request.

Pro-Actidil sustained relief in urticaria



Burroughs Wellcome & Co.
(The Wellcome Foundation Ltd.) Dartford, Kent.



#### Now is the time for Thean-the bronchodilator for 24-hour control

For high potency and better tolerance in chronic bronchitis and bronchial asthma, **Thean** is a treatment for bronchospasm that offers more than conventional medications. By having very low toxicity, being readily soluble and of neutral pH, **Thean** permits a highly effective therapeutic dosage with minimal risk of side effects. **Thean** can be given with confidence to all bronchitic patients, including those unable to tolerate sympathomimetic drugs. Further information will be sent on request.

#### Thean

Tablets 300 mg. proxyphylline

#### Thean

Ampoules 400 mg. proxyphylline

#### Thean

Suppositories 500 mg. proxyphylline



#### Hippolyte Baillière...

began to sell medical books in Paris in the early nineteenth century. In 1826 he opened a London office and, not long after, published his first medical textbook.

So began the publishing enterprise that was to become Baillière, Tindall & Cassell.

During our long career we have published medical books of many shapes and sizes, including some that are among the classics of medical literature.


Today we have many fine books on our list, and many of our authors are leaders in their fields. Some current authors and their books include:

Goligher · Surgery of the Anus, Rectum & Colon Mayer-Gross, Slater & Roth · Clinical Psychiatry Andrewes & Pereira · Viruses of Vertebrates Scheuer · Liver Biopsy Interpretation Bennett & Elliott · The Physiology & Medicine of Diving & Compressed Air Work

The company has expanded and flourished in the years since its beginning, but we are still constantly on the look-out for new authors. If you would like to write a book on your subject, and you think there is a need for it, why not write to us?

Baillière Tindall & Cassell Ltd

7 & 8 HENRIETTA STREET, COVENT GARDEN, LONDON WC2 · 01-836 3386



## dramatic disappearance of bronchitis pathogens

"...a combination of trimethoprim and sulphamethoxazole provided effective prophylaxis in chronic bronchitis. It dramatically diminished sputum volume, rendered it mucoid and abolished pathogens." 1

This combination, available as SEPTRIN\*, swiftly eradicates a wide range of organisms including the pathogens common in bronchitis—*H. influenzae, Pneumococci* and *Klebsiella* sp.

SEPTRIN has been compared in a trial with ampicillin in treating exacerbations of chronic bronchitis. SEPTRIN produced a significantly greater reduction \*Trade Mark

in sputum volume and of sputum purulence and the trial also showed that SEPTRIN produced a better overall *clinical* response in the bronchitic patients.<sup>3</sup>

- 1. Proceedings of the V International Congress of Chemotherapy (1967) Supplement (A1-5a/3) 107.
- 2. Postgrad med J. (1989) 45 SUPPLEMENT (NOVEMBER) 86 SEPTRIN Tablets, SEPTRIN Paediatric Suspension and SEPTRIN Paediatric Tablets contain trimethoprim and sulphamethoxazole.



Full information available on request.

Burroughs Wellcome & Co., Dartford, Kent,

(The Wellcome Foundation Ltd.)

Septrin-an advance on the antibiotics

# Why give any gastric ulcer time to become chronic?


# Give BIOGASTRONE first and heal it early



Further information available on request from

BERK PHARMACEUTICALS LIMITED GODALMING & SHALFORD, SURREY Made under licence from Biorex Laboratories Ltd. London. Brit, Pat. No. 843,133

Made under licence from Biorex Laboratories Ltd. London. Brit. Pat. No. 843,133 ® Regd. Trade Mark. J.3140



Normax promotes gentle, comfortable evacuation without strain to stool, by a dual action.

First, Normax prevents excessive colonic dehydration and hardening of stools.

Second, Normax acts on the lower bowel stimulating mild peristalsis.

No straining. No griping. Normal bowel function.


Normax\* capsules each contain 60 mg. dioctyl-sodium sulphosuccinate and 50 mg.1-8 dihydroxyanthraquinone and are available in bottles of 30 (basic N.H.S. price 4s. 10d.), and 250 (basic N.H.S. price 26s. 10d.).


Full information is available on request.

Normax means normal bowel function



\*Regd





#### 'I feel on edge and jittery, I can't relax.'

'Tryptizol' can promptly relieve the anxiety and agitation which accompany depression




in anxiety-depressive states

Available as 10 mg, 25 mg, and 50 mg tablets, as a syrup, and as an injection. Detailed information is available to physicians on request.

® denotes registered trademark.



Merck Sharp & Dohme Limited Hoddesdon, Hertfordshire Telephone, Hoddesdon 67123



## A new approach to mouth infections

## Oraldene

Oraldene provides you with an entirely new approach to the local treatment of mouth infections.

Outstanding properties of Oraldene are its wide spectrum of activity and its long lasting effectiveness.

Clinical experience shows that Oraldene has a strong affinity for buccal tissues, and a single rinse provides effective therapy for up to 8 hours<sup>1,2</sup>.

Oraldene has been shown to be effective against bacterial and fungal oral pathogens<sup>3</sup> and this, coupled with its absence of side-effects, makes Oraldene a remarkably effective local treatment

for such conditions as:

**Ginaivitis** 

**Stomatitis** 

Dental ulcers

Aphthous ulcers

Oral thrush in children

Geriatric oral infections

Oraldene is also a useful adjuvent to systemic therapy in tonsillitis and pharyngitis.
It does not produce sensitivity and has a soothing and refreshing effect.

Composition: Hexetidine 0.1% w/v 1,3 bis (β-ethylhexyl) 5 methyl-5-amino hexahydropyrimidine. Full information available from: William R. Warner & Co. Ltd., Eastleigh, Hampshire.

WARNER

Gingivitis, Stomatitis, Dental Ulcers, Geriatric Oral Infections.

REFERENCES: 1. Fosdick, L. D. Symposium North Western University, 23rd October, 1958. 2. Lisanti, V. F. (1958) ibid. 3. Dukes, C. D. (1959) Bact. Proc. p.48 (society report). 4. Fregni, R. (1963) Clin. ter., 27: 46.

### Experience shows

7 out of 8 patients are trouble-free with

from the very first pack



#### FIVE IMPORTANT FACTS:

- **■** contains only 50 mcg oestrogen
- used with satisfaction by more than 100,000 U.K. women
- lowest oestrogenicity of any 'combined' oral contraceptive<sup>2,3</sup>
  - first choice by over 2,000 U.K. doctors
    - four years of U.K. clinical experience

Each Norinyl-1 pack contains 21 tablets of Norethisterone B.P. 1 mg and Mestranol B.P. 50 mcg. Full prescribing information on request.

- 1. Multi-centre clinical trials involving nearly 6,000 patients. Data available to the United States F.D.A., 1968.
- 2. Rudel H.W. & Martinez-Manautou J. Animal and Clinical
- Pharmacologic Techniques in Drug Evaluation (1964).
- Gual C. Amer. J. Obst. Gyn. (1967) 97(4): 443-447.

## YOU'RE RIGHT TO CHOOSE NEGRAM

## (right from the start)

Since Negram was introduced, many investigators have compared its activity with that of other antibacterials. Studies of in vitro sensitivity of common urinary pathogens have consistently shown it to be more effective than any other agent in routine use.

So when you choose Negram as first line treatment for urinary tract infections, nine times out of ten you'll have made the right choice.

#### E. coli

|              | NEGRAM          |
|--------------|-----------------|
|              | Nitrofurantoin  |
|              | Chloramphenicol |
|              | Tetracycline    |
| Ampicillin   |                 |
| Sulphonamide |                 |

#### Proteus


|              |                |                 | Action and the second s | and the second second second |
|--------------|----------------|-----------------|--|------------------------------|
|              |                |                 | NEGRAM   |                              |
|              |                | Chloramphenicol |  |                              |
| <u> </u>     | Ampicillin     |                 |  | Ç.                           |
|              | Nitrofurantoin |                 |  |                              |
| Sulphonamide | Milotatation   |                 |  |                              |
|              |                |                 |  |                              |
| Tetracycline |                |                 |  |                              |

#### Klebsiella/Aerobacter

|                 | NEGRAM |  |
|-----------------|--------|--|
|                 |        |  |
| Nitrofurantoi   | in     |  |
|                 |        |  |
| Chloramphenicol |        |  |
|                 |        |  |
| Tetracycline    |        |  |
|                 |        |  |
| Ampicillin      |        |  |
|                 |        |  |
| Sulphonamide    |        |  |
|                 |        |  |

The figures on these charts show the average findings of the following studies:

## The most fully investigated




## as effective as morphine

## NON-HEDIT

unique amon

- ★ no analgesic tolerance on prolonged use
- \* fewer emetic sequelae than morphine
- \* no clouding of the sensorium

analgesic ever introduced



as effective as pethidine i.m.

g analgesics

- \* constipation and urine retention uncommon
- $oldsymbol{st}$  minimal risk of respiratory depression
- ★ no hypotensive effect

## negram

#### **Negram Tablets**

Formula: Each tablet contains 500 mg. nalidixic acid.

Adult Dosage: For acute infections, 2 tablets q.i.d. for at least 7 days, reducing to 1 tablet q.i.d. for chronic infections.

Basic NHS Cost: (2 q.i.d.) 6/6d. a day. Availability: Containers of 56 tablets.

#### **Negram Suspension**

Formula: Each 5ml. spoonful contains 300 mg. nalidixic acid. Paediatric dosage: 1 ml. Suspension (60 mg.) per kg. body weight (25 mg./lb.) daily for at least 7 days. For children of average weight and development: 1 month-1 year: 2.5 ml. (150 mg.) 2-4 times daily 1-2 years: 5 ml. (300 mg.) 2-3 times daily 2-6 years: 5 ml. (300 mg.) 3-4 times daily 6-14 years: 10 ml. (600 mg.) 2-4 times daily Basic NHS Cost: (5 ml. t.i.d.) 1/10d. per day. Availability: Bottles of 150 ml. and 500 ml.

#### **Side Effects:**

Gastro-intestinal or subjective visual disturbances may occur but are readily reversible on reduction or discontinuation of therapy.

#### **Precautions:**

Use with caution in patients with liver disease and raised intracranial pressure. Patients should avoid excessive exposure to sunlight. Possible risks should be carefully weighed against benefits in the first trimester of pregnancy.

Not recommended for infants less than 1 month old.

### FORTRAL

#### **Fortral Tablets**

Formula: Each Fortral tablet contains 25 mg. pentazocine hydrochloride.

Dosage: Usually 2 tablets (50 mg.) four hourly after meals, adjustable within the range of 1-4 tablets 3-4 hourly. Basic NHS Cost: 100 tablets 33/2d.

#### **Fortral Injection**

Formula: Fortral injection contains 30 mg. pentazocine per ml.

Dosage: 45-60 mg. (1.5-2 ml.) for severe pain; 30 mg. (1 ml.) for moderate pain.

Basic NHS Cost: 10 x 2 ml. ampoules 39/4d., 10 x 1 ml. ampoules 20/8d.

#### **Side Effects**

Dizziness, nausea and vomiting sometimes occur, but tend to decrease after a few doses.

#### **Precautions**

Caution in severe renal, hepatic and respiratory impairment and in patients previously on large doses of narcotics. Ambulatory patients should not drive nor operate machinery. Until further information is available Fortral should be given with caution in the first trimester of pregnancy, to children under 12 and to patients receiving MAO inhibitors.

#### **Contra-indications**

Established respiratory depression. Raised intracranial pressure.

Severe head injury or pathological brain conditions. Convulsive disorders.

Negram and Fortral are registered trade marks

Full information available on request from

The Bayer Products Company Surbiton-upon-Thames, Surrey.



## suppositories in arthritis? Yes...





## **Suppositories**

#### to treat rheumatoid arthritis and osteoarthritis at night

'Indocid' Suppositories provide for most patients:

- rapid relief of pain—the patient can move more easily in bed
- long-lasting relief of pain—from bedtime, quite often till the following afternoon
- a better night's sleep—hypnotics are frequently unnecessary
- less stiffness in the morning—the patient can usually start his routine with greater mobility

Of course, the capsules are the more widely used form of 'Indocid'—
nevertheless, "... those taking their ['Indocid'] capsules religiously every day will sometimes complain that they still have their 'early morning' stiffness in the affected joints. It is here that I think the Indocid suppository inserted at night will usually be of great value..."\*

"... even the most cynical patients will quite often ask for a repeat prescription, to have in reserve for future use."\*

\*GP, 4 Apr 1969, p7

Supplied as 100 mg suppositories, 25 mg capsules, and a suspension (25 mg/5 ml). Detailed information is available to physicians on request.

® denotes registered trademark.



Merck Sharp & Dohme Limited, Hoddesdon Hertfordshire Telephone, Hoddesdon 67123

#### **Publications**

Copies of the following publications may be obtained from E. & S. Livingstone, Limited, Teviot Place, Edinburgh

#### REPORTS FROM GENERAL PRACTICE

| No. 1. Vocational Training                                 | 5s.                |
|--|--------------------|
| No. 3. Additional Payments for Wide Experience and         |                    |
| Notable Service in General Practice                        | 1s.                |
| No. 4. General Practice in the New Towns of Britain        | 6s.                |
| No. 5. Evidence of the College of General Practitioners to |                    |
| the Royal Commission on Medical Education                  | 6s. 6d.            |
| No. 8. General Practice in South-west England              | 7s. 6d.            |
| No. 9. Obstetrics in General Practice                      | 7s. 6d.            |
| No. 10. The Practice Nurse                                 | 7s. 6d.            |
| No. 11. General Practice teaching of undergraduates in     |                    |
| British Medical Schools                                    | 10s. 6d.           |
| No. 12. A study of general practitioners' work-load in     |                    |
| South Wales, 1965-1966                                     | 10s. 6d.           |
| SYMPOSIA   |                    |
| Hazards of Middle Age                                      | 5s.                |
| Problems of Sex  | <i>5</i> s.<br>6s. |
| The Art and Science  | 7s. 6d.            |
| Accident Management  | 6s.                |
| Nutrition in General Practice                              | 7s. 6d.            |
| Arthritis in General Practice                              | 7s. ou.<br>5s.     |
| Migraine in General Practice                               | 5s.                |
| Mental Health and the Family Doctor                        | 5s.                |
| Rehabilitation   | 5s.                |
| The Actiology of Congenital Anomalies                      | 7s. 6d.            |
| The Early Stages of Chronic Bronchitis                     | 10s. 6d.           |
| Preventive Medicine in General Practice                    | 10s. 6d.           |
| The Quality of Medical Care                                | 5s.                |
| Clinical Problems of Practice                              | 8s. 6d.            |
| Anaemia in General Practice                                | 6s.                |
| The Art of Listening                                       | 7s. 6d.            |
| Early Diagnosis  | 8s. 6d.            |
| The Age of Discretion                                      | 7s. 6d.            |
| The early detection of imported and endemic disease        | 7s. 6d.            |
| Society and its general practitioners                      | 10s. 6d.           |
| Adolescence and its problems                               | 10s. 6d.           |
| Psychiatry and general practice                            | 10s. 6d.           |
| The sixth and seventh ages of man                          | 10s. 6d.           |
| Communicable diseases                                      | 10s. 6d.           |
| Rheumatology in general practice                           | 10s. 6d.           |
| OTHER PUBLICATIONS   | * .                |
| Training for General Practice (2nd edition)                | 4s. 6d.            |
| Epidemic Winter Vomiting                                   | 4s. ou.<br>1s.     |
| Memorandum for the Guidance of Trainers                    | 1s.                |
| Group Practice, Ancillary Help and Government Controls     | 7s. 6d.            |
|  |                    |

#### **CLASSIFIED ADVERTISEMENTS**

The charge for this service is 3s. 0d. per line, plus 2s. 6d. if a Box Number is required. Advertisements should be addressed to: The Journal of the Royal College of General Practitioners, Classified Advertisement Department, 104 Gloucester Place, London W1H 4AE.

GENERAL PRACTITIONER—Opportunity to join a multi-specialty group providing comprehensive medical care to university students. In-patient and out-patient facility with complete laboratory and x-ray services. Opportunities for clinical investigation and faculty appointments are available to those who are eligible and interested. Salary \$17,000 per year with advancement. Write the Director, Student Health Services, Infirmary, University of Florida, Gainesville, Florida 32601, U.S.A.

TRAINEE VACANCY, Arundel, Sussex in group practice of three with central surgery, appointment system, full ancillary staff with nurse and health visitor attachment. Cottage hospital, close liaison with consultants and full access all facilities. Active post-graduate centre. Apply Dr Warren, Arundel 3217.

EXPERIENCED PRINCIPAL (aged 43) with good academic background (University Tutor in G.P.) seeks new post away from industrial area in more conducive surroundings (town or country) where he can re-start life with his young family and practice a good standard of medicine. Box 109, The Journal of the Royal College of General Practitioners, 104, Gloucester Place, London W.1.

THIRD PARTNER assistant with view required by two partners in their thirties in a pleasant town north of Leeds. Small N.H.S. list, private practice and a variety of appointments. Full ancillary help. Recently qualified British graduate with obstetrics preferred. Write with full details to Box No. 108, Journal of the Royal College of General Practitioners, 104, Gloucester Place, London W.1.

#### THE EDUCATION FOUNDATION BOARD

This board holds funds for the furtherance of education in and for general practice, and is prepared to consider projects for personal or corporate education or educational research. They must not be such as would be covered by section 63 of the National Health Service Act, or any other similar provision.

Applications should set out the name of the applicant, the aims of the project, the methods to be employed, the estimated cost, the reasons why no other form of funding is likely to be available, and the anticipated commencing and finishing dates of the project. The board will require a typescript report of the results of the work undertaken.

Applications should be typed and sent to the Secretary, Education Foundation Board, The Royal College of General Practitioners, 14, Princes Gate, Hyde Park, London S.W. 7.

#### COLLEGE APPEAL

#### Suggested gifts

The following items are offered for sale in aid of the College appeal fund:

Price per item

s. d.

| College Ties (with all-over owl motif design)       |    |   |
|---|----|---|
| Black 100 per cent Terylene                         | 25 | 0 |
| Green 100 per cent Terylene                         | 25 | 0 |
| Green Pure Silk                                     | 35 | 0 |
| Fellows Ties (with single owl motif)                |    |   |
| Navy Blue Pure Silk (Available to F.R.C.G.P.s only) | 30 | 0 |
| Cuff Links  |    |   |

Gilt Enamel (with single owl motif) (per pair) .. .. 25 0
Postage and packing 1s. 0d. extra (U.K. only)

Orders should be addressed to the Appeal Secretary, The Royal College of General Practitioners, 14 Princes Gate, London, S.W.7.

## Domiciliary Care of the Patient with Cancer

(Proceedings of the Marie Curie Foundation Symposium)

This symposium provided a meeting point for the exchange of ideas concerning the care of patients with cancer, many of whom are nursed at home following discharge from hospitals, thus helping medical nursing and welfare services to envisage a 'total' system of care.

Edited by Ronald W. Raven, O.B.E., T.D., F.R.C.S.

108 pages

25 illustrations

36s net



William Heinemann Medical Books Ltd., 23 Bedford Square, London W.C.I

## intertulle<sup>®</sup> fucidin<sup>®</sup>

### the reasons why



easy to handle



easy to cut to any size or shape



easy to use


and each square is impregnated with FUCIDIN OINTMENT. (2% Sodium Fusidate B.P. in a neutral ointment base) for rapid bactericidal power against staphylococci and streptococci with remarkable freedom from unwanted side-effects.


Presented in special white plastic packs containing 10, 4" x 4" sterile gauze squares, doubly-interleaved for ease of handling.

#### a case in point

Varicose ulcer on leg of 74-year-old man

After 5 week's treatment with Intertulle Fucidin Photographs from Leo Laboratories' Library of British Dermatological Case Histories, 1969.







R Regd. Trade Marks

LEO LABORATORIES LIMITED • HAYES • MIDDLESEX • TELEPHONE: 573 6224