
in urinary infections 'clearly superior' to ampicillin or sulphadimidine



Higher cure rates

These comparative results were obtained in treating hospital in-patients with UTI1, most of whom were infected with entrenched hospital pathogens.

85% cure rate	SEPTRIN*
70% cure rate	ampicillin
40% cure rate	sulphadimidine

(Cure criterion = eradication of original organism 1 week after treatment ended.)

SEPTRIN was "clearly superior to either of the other treatments" and this superior effect was "even more appreciable in the re-treatment of patients who had failed on another drug,"2

SEPTRIN Tablets, SEPTRIN Paediatric Suspension and SEPTRIN Paediatric Tablets contain trimethoprim and sulphamethoxazole.
Full information is available on request.

References: 1. Brit med J, (1969) 1, 541. 2. Leading Article, Brit med J, (1969) 1, 525 *Trade Mark

Clear reasons for using Septrin

Abstracted from a Leading Article in the British Medical Journal²

- ☐ Trimethoprim not only attains therapeutic tissue levels, but is excreted unchanged in high concentration in the urine.
- ☐ All Gram-negative bacilli occurring in the urine, except Pseudomonas aeruginosa, are usually highly sensitive to it.
- ☐ Its minimum inhibitory concentration is extremely low.
- ☐ Trimethoprim and a sulphonamide, by blocking successive stages in the same bacterial metabolic process, exert a strong synergic effect, and this action is bactericidal.



for bacterial infections

an advance on the antibiotics

When Anxiety is more than a bitten nail



TRIPTAFEN-MINOR

calms the patient and regulates associated depression

Bitten nails can be a sign that your patient is worried, nervous, depressed. His confidence is shattered and his life is a muddle. He seeks your advice. He needs the benefit of your understanding—and Triptafen-Minor to calm him.


Triptafen-Minor is ideal for patients with minor neuroses who need treatment – the sort you see so often. It might be the housewife who feels that she can no longer cope. It might be the executive whose job and home worries are too much. It could be a patient who has suffered a recent bereavement.

Whatever the cause, Triptafen-Minor will calm your patient and regulate any associated depression.

Each Triptafen-Minor tablet contains: Amitriptyline hydrochloride BP 10 mg, Perphenazine BP 2 mg.

Full information is available on request.

Triptafen is a Trade Mark of ALLEN & HANBURYS LTD LONDON E2



In hay fever Pro-Actidil Good for up to 24 hours

Each PRO-ACTIDIL* Tablet is specially formulated in 3 layers for prolonged antihistamine action lasting 19-24 hours.
Full information about PRO-ACTIDIL Triprolidine Hydrochloride Tablets is available on request.

Burroughs Wellcome & Co., (The Wellcome Foundation Ltd.) Dartford, Kent.

An advance in the treatment of Parkinson's Disease

Symmetrel® Geigy

"Functional disability, tremor, assessment of limb dexterity, and the timing of walking and of writing, all improved significantly."

Lancet. 1 7641 259 (1970)

Availability
Symmetrel® is available as capsules containing 1-adamantanamine
(Amantadine) hydrochloride 100 mg

Further information will be sent on request. Geigy (U.K.) Limited, Pharmaceuticals Division, Macclesfield, Cheshire

Halt at the flashing light

Flashing lights and visual disturbances, classically early warnings of migraine, are signals for arresting all manifestations of an attack – at onset.

MIGRIL* acts in the minutes between initial warning and the otherwise inevitable full-blown attack.

As well as containing an effective dose of ergotamine (2mg in each tablet), enhanced by coffeine, MIGRIL contains the anti-emetic, cyclizine. Nausea and vomiting, due to migraine itself or to ergotamine, are very effectively relieved.

MIGRIL, taken at the first warning of an attack, provides a specific treatment that the patient will learn to rely on. Full information is available on request.

*Trode Mor

Burroughs Wellcome & Co. (The Wellcome Foundation Ltd) Dartford, Kent.



Migril masters migraine

Whenever oral iron is indicated

FERRODIC

Trada Mar

means effective, well tolerated and palatable treatment

Ferrodic tablets each contain:
Ferrous Iron (as ferrous carbonate) 50 mg
Ascorbic Acid 12.5 mg



Full information is available on request

Ferrodic is a Trade Mark of

ALLEN & HANBURYS LIMITED LONDON E2

DISEASES OF THE NERVOUS SYSTEM

Described for Practitioners and Students

Sir Francis Walshe, MD, DSc, FRCP (Lond.), FRS

1970 Eleventh Edition

416 pages 44 illus. 60s. Post 4s. 6d.

Extract from preface

The steady growth of knowledge since the publication of the tenth edition of this text, in some though not in all aspects of neurological medicine, has increased the difficulty of selection of material for a work designed for a specified circle of readers. This has made the danger of reducing the book to a pocket encyclopaedia, containing too little of too much, more difficult to avert.

Yet it still remains selective, dealing with uncommon diseases only when they exemplify new ways of looking at things that promise fruitful developments in clinical neurology and may close the wide gap in the effective control and treatment of the non-infective diseases of the nervous system that has hitherto been a characteristic of neurological medicine.

The division of the book into two parts: the first dealing with the fundamentals of neurological diagnosis, and the second with an account of the more common nervous disorders, is retained.

Opinions of previous editions

- "This book is now one of the classics of medicine. Clear, concise and authoritative, it bears the hallmark of the author's clinical acumen and wise knowledge of the speciality of which he is the outstanding representative at the present day."—

 Practitioner
- "This book is so well known that it really needs no recommendation. The subtitle of the work is Described for Practitioners and Students, and the emphasis is rightly on the practical aspects of neurology.
- "This book has been called one of the classics of medicine. It can be whole-heartedly recommended."—Middlesex Hospital Journal

Published by

E. & S. LIVINGSTONE

15-17 Teviot Place - Edinburgh



Sitting in comfort with XYLOPROCT

XYLOPROCT is a rapidly effective all-round treatment for haemorrhoids, anal fissures and pruritus ani.

XYLOPROCT is specially formulated for the control, treatment and immediate relief of pain in all types of haemorrhoids.

XYLOPROCT contains:

- *Aluminium Acetate as the antiseptic-astringent.
- *Zinc Oxide for soothing-healing. Hydrocortisone as the antiinflammatory.
- inflammatory.
 *XYLOPROCT for relief of pain.
 Available as SUPPOSITORIES in boxes
 of 10 and OINTMENT in tubes of 15g.

Full information and samples available on request.



Astra Chemicals Ltd. Watford, Herts.

Books from Baillière

GARRAD AND ROSENHEIM

Social Aspects of Clinical Medicine

by Jessie Garrad, B.A., B.Sc., A.I.M.S.W., and Sir Max Rosenheim, K.B.E., M.A., M.D., D.Sc.

Medical students are keenly aware of the importance of this subject, and the study of it is now required in most medical schools. This book presents an examination of the environmental and emotional factors which may affect a patient's disease and the means by which community services can be enlisted in his aid.

184 pages

28s./£1.40

ANDERSON AND ISAACS

Current Achievements in Geriatrics 268 pages 23 illus 45s/£2.25

DRURY

The Medical Secretary's Handbook
2nd edition 326 pages 40 illus 6 plates 40s/£2

ISAACS

An Introduction to Geriatrics
212 pages 16 illus

KING AND NICOL

Venereal Diseases

2nd edition 320 pages 180 illus 4 colour plates 75s/£3.75

Baillière's Handbook of First Aid 6th edition 352 pages 180 illus 20s/£1

MILES AND ROYLANCE **Teaching First Aid**

18s/90p

40s/£2

Baillière Tindall and Cassell 7 & 8 Henrietta Street London W C 2

116 pages



Inderal-80 brings new stability to the hypertensive patient

Smooth and gradual reduction of blood pressure. No postural or exercise hypotension. Round-the-clock control of blood pressure, even whilst supine. Effective in renal and essential hypertension. Minimal side effects. 'Inderal'-80 depends for its effect on a specific pharmacological action. Before using 'Inderal'-80,

it is important that doctors should read the detailed literature for information on dosage, precautions and contraindications. We shall be glad to send you further information.

Inderal-80







PROPRANOLOL HYDROCHLORIDE B.P.

Imperial Chemical Industries Ltd., Pharmaceuticals Division, Alderley Park, Macclesfield, Cheshire.





TO PROBLEM AND PROPERTY OF THE PROPERTY OF THE

when acute peri-articular disorders lead to loss of work or play





is treating pain with simple analgesics enough?

Pain must be relieved of course; but it is the inflammation behind the peri-articular disorder that can delay recovery. This is why 'Indocid' can play such an important part in treatment.

'Indocid' is both analgesic and anti-inflammatory. 'Indocid' may help patients begin graded exercise sooner—and hence speed full recovery.



Supplied as 25 mg and 50 mg capsules, 100 mg suppositories, and a suspension (25 mg/5 ml). Detailed information is available to physicians on request.



NORINYL-1

hormonally 'balanced' oral contraceptive

ideal for every generation of fertile woman

- □ 4 years' U.K. clinical experience
- □ lowest incidence of side-effects because of minimal bio-potency 1,2
- \square used with satisfaction by nearly 200,000 U.K. women
- ☐ first choice by over 2,000 U.K. doctors




References

- 1. Med. Sci. (1967) May p. 37-49
- 2. Brit. Med. J. (1969) 4: 803

Each Norinyl-1 pack contains 21 tablets of Norethisterone B.P. 1mg and Mestranol B.P. 50 mcg







when the colour goes out of eating

'Periactin' can improve the picture

- ☐ clinically proved appetite stimulant
- ☐ increased food intake in most patients
- □ weight gain is usually seen after a week
- □ suitable for children and adults where weight gain is desirable

Periactin Cyproheptadine hydrochloride MSD ®

for patients who lack appetite

Supplied as 4 mg tablets and as a syrup (2 mg/5 ml). Detailed information is available to physicians on request. \circledR denotes registered trademark.



Merck Sharp & Dohme Limited, Hoddesdon, Hertfordshire Telephone, Hoddesdon 67123

YOU'RE RIGHT TO CHOOSE NEGRAM

(right from the start)

Since Negram was introduced, many investigators have compared its activity with that of other antibacterials. Studies of in vitro sensitivity of common urinary pathogens have consistently shown it to be more effective than any other agent in routine use.

So when you choose Negram as first line treatment for urinary tract infections, nine times out of ten you'll have made the right choice.

E. coli

		NEGRAM
		Nitrofurantoin
		Chioramphenicol
	Tetracycline	
A	mpicillin	
Sulphona	mide 4	

Proteus

	NEGRAM
Chloramphenicol	
Ampicillin	
Nitrofurantoin	
Sulphonamide	
Tetracycline	

Klebsiella/Aerobacter

	NEGRAM
Nitrofurantoin	
Chloramphenicol	
Tetracycline	
Ampicillin	
Sulphonamide	

Take the pour pr



Slipped Disc



Maiignancy



Osteoarthritis



Trigeminal Neuralgia

wherever pai for ordinary

Fortral Tablets Dosage: usually 2 tablets (50 mg.) four hourly after meals, adjustable within the range of 1-4 tablets 3-4 hourly. Each Fortral tablet contains 25 mg. pentazocine hydrochloride. Basic NHS cost: 100 tablets 33/2. Fortral Injection Dosage: 45-60 mg. for severe pain; 30 mg. for moderate pain. Fortral Injection contains 60 mg. pentazocine per 2 ml. Basic NHS cost: 10 x 1 ml. ampoules 20/8d. 10 x 2 ml. ampoules 39/4d. Side Effects Dizziness, nausea and vomiting sometimes occur, but tend to decrease after a few doses.

ain out of actice



Coronary



Herpes Zoster



Sprains and Strains



Fractures

n is too great analgesics

Precautions Caution in severe renal, hepatic and respiratory impairment and in patients previously on large doses of narcotics. Ambulatory patients should not drive nor operate machinery. Until further information is available Fortral should be given with caution in the first trimester of pregnancy, to children under 12 and to patients-receiving MAO inhibitors.

Contra-Indications Established respiratory depression, Raised intracranial pressure.

Severe head injury or pathological brain conditions. Convulsive disorders.

Fortral is a registered trade mark. Full information on request from

The Bayer Products Company, Surbiton-upon-Thames, Surrey.



negram

Negram Tablets

Formula: Each tablet contains 500 mg. nalidixic acid.

Adult Dosage: For acute infections, 2 tablets q.i.d. for at least 7 days, reducing to 1 tablet q.i.d. for chronic infections.

Basic NHS Cost: (2 q.i.d.) 5/10d. a day. Availability: Containers of 56 tablets.

Negram Suspension

Formula: Each 5ml. spoonful contains 300 mg. nalidixic acid. Paediatric dosage: 1 ml. Suspension (60 mg.) per kg. body weight (25 mg./lb.) daily for at least 7 days. For children of average weight and development: 1 month-1 year: 2.5 ml. (150 mg.) 2-4 times daily 1-2 years: 5 ml. (300 mg.) 2-3 times daily 2-6 years: 5 ml. (300 mg.) 3-4 times daily 6-14 years: 10 ml. (600 mg.) 2-4 times daily Basic NHS Cost: (5 ml. t.i.d.) 1/10d. per day. Availability: Bottles of 150 ml. and 500 ml.

Side Effects:

Gastro-intestinal or subjective visual disturbances may occur but are readily reversible on reduction or discontinuation of therapy.

Precautions:

Use with caution in patients with liver disease and raised intracranial pressure. Patients should avoid excessive exposure to sunlight. Possible risks should be carefully weighed against benefits in the first trimester of pregnancy.

Not recommended for infants less than 1 month old.

FORTRAL

Fortral Tablets

Formula: Each Fortral tablet contains 25 mg. pentazocine hydrochloride.

Dosage: Usually 2 tablets (50 mg.) four hourly after meals, adjustable within the range of 1-4 tablets 3-4 hourly. Basic NHS Cost: 100 tablets 33/2d.

Fortral Injection

Formula: Fortral injection contains 30 mg. pentazocine per ml.

Dosage: 45-60 mg. (1.5-2 ml.) for severe pain; 30 mg. (1 ml.) for moderate pain.

Basic NHS Cost: 10 x 2 ml. ampoules 39/4d., 10 x 1 ml. ampoules 20/8d.

Side Effects

Dizziness, nausea and vomiting sometimes occur, but tend to decrease after a few doses.

Precautions

Caution in severe renal, hepatic and respiratory impairment and in patients previously on large doses of narcotics. Ambulatory patients should not drive nor operate machinery. Until further information is available Fortral should be given with caution in the first trimester of pregnancy, to children under 12 and to patients receiving MAO inhibitors.

Contra-indications

Established respiratory depression. Raised intracranial pressure.

Severe head injury or pathological brain conditions. Convulsive disorders.

Negram and Fortral are registered trade marks

Full information available on request from

The Bayer Products Company Surbiton-upon-Thames, Surrey.



effective in controlling infection in these circumstances. Erythromycin estolate appeared somewhat more effective. Side effects were minimal with both antibiotics.

Acknowledgements

Our thanks are due to Dr H. St. C. C. Addis for allowing us to study patients under his care, and to Dr A. R. Evans for bacteriological studies, and to Dr G. L. Smith of Eli Lilly & Co., for his co-operation in the trial.

REFERENCES

- MRC Report on Trials of Chemotherapy in early chronic bronchitis. (1966). British Medical Journal. 1, 1317.

 Percival, A., Armstrong, Elizabeth C., and Turner, G. C. (1969). Lancet. 1, 998.
- Sabath, L. D., Gerstein, Deborah A., and Finland, M. (1968). New England Journal of Medicine. 279, 1137.
- Schnoor, T. C. (1939). American Journal of Tropical Medicine. 19, 163
- Tewari, S. N., and Fletcher, R. (1966). British Journal of Clinical Practice. 20, 639.

Publications

Copies of the following publications may be obtained from E. & S. Livingstone, Teviot Place, Edinburgh

Reports from General Practice		Rehabilitation	5s.	
No. 1. Vocational Training	5s.	The Aetiology of Congenital Anomalies	7s.	6d.
No. 3. Additional Payments for Wide	• • • • • • • • • • • • • • • • • • • •	The Early Stages of Chronic Bronchitis	10s.	6d.
Experience and Notable Service	<u> </u>	Preventive Medicine in General Practice	10s.	6d.
in General Practice	1s.	The Quality of Medical Care	5s.	
No. 4. General Practice in the New Towns	1	Clinical Problems of Practice		6d.
of Britain	6s.	Anaemia in General Practice	6s.	
No. 5. Evidence of the College of General	1	The Art of Listening	7s.	6d.
Practitioners to the Royal Com-		Early Diagnosis	8s.	6d.
mission on Medical Education	6s. 6d.	The Age of Discretion	7s.	6d.
No. 6. Implementation of vocational		The early detection of imported and		
training	4s.	endemic disease		6d.
No. 10. The practice nurse	10s.	Society and its general practitioners	10s.	
No. 11. General practice teaching of	!	Adolescence and its problems	10s.	
undergraduates in British Medical		Psychiatry and general practice	10s.	
Schools	10s. 6d.	Sixth and seventh ages of man	10s.	
No. 13. Present state and future needs of	•	Communicable diseases	10s.	
general practice (Second edition)	12s. 0d.	Rheumatology in General Practice	10s.	
Symposia		The management of staff in general practice		
	_	Man, milieu and malady	10s.	
Hazards of Middle Age	5s.	A future in general practice	10s.	6d.
Problems of Sex	6s.	Other Publications		
The Art and Science	7s. 6d.			
Accident Management	6s.	Training for General Practice (2nd Edition	•	6d.
Nutrition in General Practice	7s. 6d.	Epidemic Winter Vomiting	1s.	
Arthritis in General Practice	5s.	Memorandum for the Guidance of Trainers	1s.	
Migraine in General Practice	5s.	Group Practice, Ancillary Help and	_	
Mental health and the Family Doctor	5s.	Government Controls	7s.	6d.

CLASSIFIED ADVERTISEMENTS

The charge for this service is 3s. 0d. per line, plus 2s. 6d. if a Box Number is required. Advertisements should be addressed to: The Journal of the Royal College of General Practitioners, Classified Advertisement Department, 5 Bentinck Street, London W1H 4AE.

TWO YOUNG PARTNERS seek third for Group Practice ideally situated in Nottingham's premier Residential suburb. Full appointment system and open access to all investigational facilities. District Nurse and Health Visitor attached and two secretary/receptionists employed. Purpose-designed premises featured in forthcoming R.C.G.P. exhibition and *Practitioner* article. No capital required but eligibility for Obstetric list and previous general practitioner experience desirable.—Box No. 113, The Journal of the Royal College of General Practitioners, 5 Bentinck Street, London, W.1.

CONFERENCE

'OUT OF HOSPITAL—WHO CARES?"

on Tuesday, September, 22nd, 1970. at Bedford College, London, N.W.I

Chairman: Dr A. S. Gatherer, Medical Officer of Health, County Borough of Reading.

General practitioners are invited to attend this conference for doctors, nurses and social workers concerned with the treatment and care of mentally ill and severely subnormal patients and their families.

Programmes and application forms from the National Association for Mental Health, Social Services Department, 39 Queen Anne Street, London, WIM OAJ. (01-935 1217 ext. 22).

COLLEGE APPEAL

Suggested gifts

The following items are offered for sale in aid of the College appeal fund:

in aid of the College appeal fu	ınd:		
	Price	<i>per it</i> s.	em d.
College Ties			
Green silk (with owl motif) Black 100 per cent Terylen		35	0
owl motif)		25	0
Cuff Links: Rolled gold with gilt ename ing the owl motif	el, bear-	25	0
		per p	
Ash Trays: Tinted blue glass, decorated with the College crest	_		6
Fellows Ties: (with a single ov Navy blue silk (Available to	Fellows		_
only)	• ••	35	0
Postage and packing 1s. 0d. Northern Ireland only)	extra (U	J .K. a	nd

Orders should be addressed to the Appeal Secretary, The Royal College of General Practitioners, 14 Princes Gate, London, S.W.7.

THE EDUCATION FOUNDATION BOARD

This board holds funds for the furtherance of education in and for general practice, and is prepared to consider projects for personal or corporate education or educational research. They must not be such as would be covered by section 63 of the National Health Service Act, or any other similar provision.

Applications should set out the name of the applicant, the aims of the project, the methods to be employed, the estimated cost, the reasons why no other form of funding is likely to be available, and the anticipated commencing and finishing dates of the project. The board will require a typescript report of the results of the work undertaken.

Applications should be typed and sent to the Secretary, Education Foundation Board, The Royal College of General Practitioners, 14, Princes Gate, Hyde Park, London S.W. 7.

The Journal of the Royal College of Physicians of London

This Journal is concerned with the integration of scientific disciplines in the practice of medicine and, by providing a wide ranging commentary on the growing points of medicine.

is an essential complement to the specialised journals.

CONTENTS OF VOLUME 4, No. 4

S. H. Curry	"Drug Distribution and Effect."
M. Gracey and	"Cliffic I Malayeria Di I CV
Charlotte M. Anderson	"Childhood Malabsorptive Disorders of Increasing Importance in Adult Life."
I. H. Mills	"Regulation of Sodium Excretion: Intra-and Extrarenal Mechanisms."
W. R. Thrower	"Agriculture and the Public Health."
Dame Albertine L. Winner	"Death and Dying."
Editor	A. Stuart Mason, M.D., F.R.C.P.
Publication	Published quarterly in October, January, April and July.
Subscription	80/- per annum (including postage).
All enquiries to Editorial Office	Royal College of Physicians of London, 11 St Andrew's Place, Regent's Park, London, N.W.1.

Publishing Office

Pitman Medical and Scientific Publishing Co. Ltd., 31 Fitzroy Square, London, W.1.

intertulle[®] fucidin[®]

the reasons why



easy to handle



easy to cut to any size or shape



easy to use

and each square is impregnated with FUCIDIN OINTMENT (2% Sodium Fusidate B.P. in a neutral ointment base) for rapid bactericidal power against staphylococci and streptococci with remarkable freedom from unwanted side-effects.


Presented in special white plastic packs containing 10, 4" x 4" sterile gauze squares, doubly-interleaved for ease of handling.


a case in point

Varicose ulcer on leg of 74-year-old man

After 5 week's treatment with Intertulle Fucidin

Photographs from Leo Laboratories' Library of British Dermatological Case Histories, 1969.







Regd. Trade Marks

LEO LABORATORIES LIMITED • HAYES • MIDDLESEX • TELEPHONE: 573 6224