The Beecham Penicillins

Beecham research has produced an outstanding range of semi-synthetic penicillins including:

Orbenin*(cloxacillin) Pyopen*(carbenicillin)

Penbritin* (ampicillin) Floxapen* (flucloxacillin) Celbenin* (methicillin) Broxil* (phenethicillin)

Ampiclox*

(ampicillin and cloxacillin)

BRL also produce Maxolon*(metoclopramide) —a modifier of gastric motility.

Full information on all products is available on request.



Beecham Research Laboratories Brentford England



Whatever the Allergy label

PHENERGAN

the general purpose antihistaminic




Effective, Versatile, Reliable


"If I have to nominate any drug as being my favourite, then the honour must go to 'Phenergan', for it goes closer to being a panacea than any other substance I know."

G.P., 6, 7, 1969.

'Phenergan' is a trade mark of May & Baker Ltd for its preparations of promethazine








ETOPHYLATE Syrup is readily accepted by children of all ages. A clinical investigation involving 22 children under the age of 12 months showed that ETOPHYLATE was useful in the relief of infective bronchospasm and devoid of any harmful effects.

Reference: The Practitioner (1968) 200: 833-835

Further information available on reques

DELANDALE LABORATORIES LIMITED. 37 Old Dover Road, Canterbury, Kent.





Jim had a peptic ulcer. His diet was hopelessly monotonous, he couldn't work and he was due for surgery

gery.
On Roter he got almost immediate relief of pain, his ulcer healed quickly, and hospitalisation and surgery were avoided.

Lucky Jim is back at work He'son a normal diet; is enjoying his meals and putting on weight

> Roter: available on E.C.10 in packs of 40 and 120 tablets. Dispensing packs of 360 and 720 tablets.

Roter: Bismuth subnitrate Roter 300 mg., Magnesium carbonate 400 mg. Sodium bicarbonate 200 mg., Cortex rhamni frangula 25 mg.

Citerature and abstracts of clinical tries available on request. F.A.I.R. Laboratories Limited. Twickenham. Middlesex.



Butazolidin Alka

The modern way to prescribe an antirheumatic agent

Unique formulation

Proved effectiveness

Excellent gastric toleration


their previous drug".

Pract. 203, 1215, 357-360 (1969)


Formulated as a mantle
of two antacids around a core
of Butazolidin® and antacid,
each tablet contains
Butazolidin (Phenylbutazone B.P.)
100 mg. Magnesium Trisilicate
B.P. 150 mg. Dried Aluminium
Hydroxide Gel B.P. 100 mg.

Detailed literature describing any Geigy product will be supplied on request.

Geigy Pharmaceuticals, Macclesfield, Cheshire.








Certainly several important risk factors are present: overweight-stress-cigarette smoking, and, frequently found in the presence of the other three - raised blood lipids.


In the treatment of hyperlipidaemia, Atromid-S has been shown to produce remarkably consistent results; it also exerts a beneficial effect on abnormal clotting factors.

for the patient at risk









Actifed and Space age Medicine

The astronaut above is about to be subjected to G-forces in a massive centrifuge. In these situations it was thought that G-forces might restrict astronauts from performing vital tasks. To overcome this, an ingenious device which bounces light into the eye, can be made to actuate a switch or system, whenever the pupil crosses the path of the beam. This sight switch has already been adapted to help paraplegics to help themselves.

In the space programme, medical benefit has been a two-way exchange. Of the medicines taken on Apollo flights, ACTIFED* has been repeatedly chosen as the decongestant. ACTIFED TABLETS and SYRUP contain the decongestant pseudoephedrine and the established antihistamine triprolidine. ACTIFED COMPOUND LINCTUS contains, in addition, codeine. It provides effective and palatable cough control for all age groups.

Full information is available on request.

*Trade Mark
Burroughs Wellcome & Co. (The Wellcome Foundation Ltd.) Dartford, Kent.

For Coughs with Congestion

ACTIFED COMPOUND LINCTUS





Day after day of freedom from asthmatic attack



The unique mode of action of disodium cromoglycate offers the asthmatic patient outstanding benefits. A prophylactic treatment, it prevents the onset of asthmatic episodes – thus minimising long-term damage to lung tissue. Freedom from asthma attacks permits a happier, more normal life for the asthmatic patient.

Published clinical reports confirm that with INTAL COMPOUND, the incidence and severity of acute attacks is considerably reduced - even in patients on conventional therapy such as bronchodilators, antihistamines and corticosteroids. Less time is lost from work or school, exercise tolerance is improved and a consistent clinical comment on disodium cromoglycate has been the total absence of serious side effects.

INTAL COMPOUND - supplied in SPINCAPS (each containing 20 mg disodium cromoglycate, and 0.1 mg Isoprenaline Sulphate BP in powder form, together with an inert carrier). For use in a special insufflator-SPINHALER.

Full information is available on request. The Technical Services Department, Fisons Limited-Pharmaceutical Division, Loughborough, Leicestershire.

There's nothing else like

INTAL Co.

Prophylaxis in asthma



INTAL, INTAL COMPOUND, SPINCAP and SPINHALER are Fisons trade mark

New assisted passage scheme

The formula which really assists and maintains free mucus flow in bronchial conditions—that is new LINCTIFED* EXPECTORANT: LINCTIFED EXPECTORANT: loosens thick tenacious sputum; decreases mucosal congestion; controls irritating, unproductive cough. *Trade mark*

Available in two forms—
LINCTIFED EXPECTORANT and
LINCTIFED EXPECTORANT PAEDIATRIC
containing guaiphenesin, pseudoephedrine,
triprolidine and codeine in a balanced
formulation. LINCTIFED EXPECTORANT
controls cough and clears the way for
easier breathing.
Full information available on request.

LINCTIFED EXPECTORANT LINCTIFED EXPECTORANT PAEDIATRIC





Burrayaba Wellcome & Co. (The Wellcome Foundation Ltd.) Dertford, Kent

a fundamental? Exectment for anxiety!

MEGRIN 10

OSCAPSULE CAPSULE

associated with



the 'fundamental' treatment for anxiety!

Found effective in 90% of patients

In recent G.P. studies improvement was shown in 9 out of 10 patients within 2 weeks. In double-blind comparisons with chlordiazepoxide, Integrin gave relief to nearly 10% more patients.

Gives more <u>profound</u> relief from anxiety

Seventy eight patients, most of whom were already on tranquillo-sedatives, were switched to Integrin. Their average anxiety-rating dropped to less than one third of its previous level.

In another trial, Integrin gave <u>complete</u>—as opposed to partial—relief from anxiety within 4 weeks to as many as 35% of patients: nearly twice the rate of chlordiazepoxide.

Lifts any depressive overlay

In contrast to the tranquillo-sedatives, Integrin actually enhances mood and restores a normal sense of well-being.

◆ Fast action—fast clinical response

Integrin answers the need for a faster acting tranquillizer. Clinical trials show that up to 90% of patients improve within the first week of treatment.

drowsiness not a serious problem	
virtually free from toxicity	
one single strength capsule with simple dosage	
unlike the benzodiazepines, no reports of dependence	
relieves depressive overtones	
no effect on reaction-time at normal dosage	
well tolerated even at 10 times normal dosage	



Indications:

Acute and chronic anxiety states and psychosomatic conditions whether or not accompanied by depressive overlay; tension, apprehension, agitation and sleep disturbances.

Dosage:

The usual adult dose is one 10mg capsule three times daily, with a further capsule at bedtime if required. In certain cases up to 60mg daily may be needed. Should a milder effect be required the midday dose may be omitted.

Side Effects:

Even in high doses, Integrin is well tolerated, with only minor, often transient, side effects. Some patients may experience some drowsiness. Dry mouth and dizziness have also been reported.

Precautions:

Alcohol or other tranquillizers may potentiate the effect of Integrin and patients intending to drive should be made aware of this. Where MAO inhibitors have been used, 2-3 days should elapse before Integrin is started.

No substantiated reports of toxicity have been received, even where patients have taken up to the equivalent of 56 capsules in 24 hours. Neither dependence nor withdrawal symptoms have been reported.

Presentation:

Containers of 100 capsules. Each grey/green capsule contains 10mg oxypertine.

Basic NHS Cost:

One week's treatment (1 capsule 10mg t.d.s.) is 36p

Integrin is a registered trade mark.

Full information is available from

Winthrop Laboratories, Surbiton-upon-Thames, Surrey.



Airst in bronchitis

Septrin* bactericidal to respiratory tract pathogens including *Haemophilus influenzae*.

Septrin quickly reaches effective blood levels and produces an exceptionally rapid clinical response.

Septrin was more effective compared with antibiotics (ampicillin and tetracycline) in reducing sputum purulence and sputum volume.1.2

Septrin first utilisation of antibacterial synergy between trimethoprim and a sulphonamide. *Trade Mark

References


1. Brit med J, (1969), **4**, 470. 2. Postgrad med J, (1969), **45**, Supplement, (November) 91.

For the whole family-SEPTRIN Tablets, SEPTRIN Adult Suspension, SEPTRIN Paediatric Tablets and SEPTRIN Paediatric Suspension containing trimethoprim and sulphamethoxazole.


Full Information is available on request.



Burroughs Wellcome & Co. (The Wellcome Foundation Ltd.) Dartford, Kent.



SEPTRIN eradicates infection by a double blockade of bacterial metabolism.



The low bio-activity of both progestin and oestrogen components 1.2 in Notiny 1-1 produce a combination which, while providing high contraceptive reliability, can reduce the incidence of troublesome side-effects 3 and maintain high usage rates.4

Norinyl-1, which has now been in world-wide use for several years, is used with satisfaction by nearly ‡ million women in Britain alone and prescribed as first choice oral contraceptive by over 2,000 U.K. dectors.

NORINYL-1

provides the confidence you both need.

Norinyl-1 contains Mestranol 50 mcg Norethisterone 1-0 mg.

REFERENCES

- 1. Medical Science (1967) 3. Lance May p. 37-49. 4. Multi
- 2. Contraception (1970)
- 3. Lancet (1967) (2): 612. 4. Multi-centre trial data available

to F.D.A. (1968)

Full prescribing information available on request:



Syntex Pharmaceuticals Ltd. St. Ives House, Maidenhead, Berkshire. British graduate, Edinburgh, 1965. Aged 30. Married. MRCGP, D.Obst.RCOG, Dip FPA. Desires partnership in semi-rural group practice with early parity. Good schools.

Available February, 1972, when service with

Available February, 1972, when service with RAF terminates. Dr R. Bond, Medical Centre, Royal Air Force, Lyneham, Chippenham, Wilts. Telephone: Bradenstoke 381.

RURAL PRACTITIONER, age 50, teaching experience and higher qualifications, seeks partnership in group practice, preferably working from health centre. Family need to return to urban life. Box No. 126, JOURNAL OF THE ROYAL COLLEGE OF GENERAL PRACTITIONERS, 5 Bentinck Street, London, W.1.

PARTNERSHIP following assistantship with view for not more than six months (at £3,500 p.a.) to replace retiring third partner, on or after November 1. Parity in three years. British graduate under 30 with vocational training preferred. Appointment system; rota; full secretarial help; practice nurse; attached LHA staff; general-practitioner maternity hospital; full diagnostic facilities; postgraduate medical centre; health centre within 18 months. Drs J. R. Edwards and J. K. Lloyd, 94 Sidwell Street, Exeter, Devon.

CLASSIFIED ADVERTISEMENTS

The charge for this service is 3s. 0d. per line, plus 2s. 6d. if a Box Number is required. Advertisements should be addressed to: The Journal of the Royal College of General Practitioners, Classified Advertisement Department, 5 Bentinck Street, London W1H 4AE.

Publications

Copies of the following publications may be obtained from E. & S. Livingstone Limited, Teviot Place, Edinburgh.

	Reports from General Practice		The Aetiology of Congenital Abnormalities	37 1
No. 1.	Vocational Training	25p	The Early Stages of Chronic Bronchitis	52½p
No. 3.	Additional Payments for Wide	•	Preventive Medicine in General Practice	52½p
	Experience and Notable Service		The Quality of Medical Care	25p
	in General Practice	5p	Clinical Problems of Practice	42½p
No. 4.	General Practice in the New Towns	•	Anaemia in General Practice	30p
	of Britain	30p	The Art of Listening	37½p
No. 5.	Evidence of the College of General	•	Early Diagnosis	42½p
	Practitioners to the Royal Com-		The Age of Discretion	37½p
	mission on Medical Education	32½p	The early detection of imported and endemic	
No. 8.	General Practice in South-west		disease	37 կ ը
	Wales	37 12 p	Society and its general practitioners	52½p
No. 9.	Obstetrics in General Practice	$37\frac{1}{2}p$	Adolescence and its problems	52½p
No. 10.	The Practice Nurse	37 ½ p	Psychiatry and general practice	52½p
No. 11.	General Practice teaching of under-		Sixth and seventh ages of man	52 1 ₽
	graduates in British Medical Schools	52 1 2p	Communicable diseases	52½p
No. 14.	The Future General Practitioner	25p	The Management of Staff in General Practice	52½p
			Man, Milieu and Malady	52½p
	Symposia		Group Practice	75p
Hazards of Middle Age		25p	Survey of General Practitioners' care of their	
Problems of Sex		30p	Patients in the Hospital Service	30p
The Art and Science		37 1 2p		
Accident Management		30p	04 13 4	
Nutrition in General Practice		37 1 p	Other publications	
Arthritis in General Practice		25p	Training for General Practice (2nd Edition)	22 <u>1</u> p
Migraine in General Practice		25p	Epidemic Winter Vomiting	5p
Mental Health and the Family Doctor		25p	Memorandum for the Guidance of Trainers	5p
Rheumatology in general practice		52 1 2p	Group Practice, Ancillary Help and Govern-	
Rehabilitation		25n	ment Controls	371r

intment

2% sodium fusidate B.P. in a neutral pintment base.

The logical topical antibiotic for primary and secondary bacterial skin infections.

Fucidin Ointment has proved clinically effective in a wide variety of common skin infections. "Its rapid, reliable, bactericidal action against most strains of staphylococcus and streptococcus combined with its good skin tolerance and proven penetration," make it a logical topical treatment wherever infection is present or threatens the skin.

Available in packs of 10g and 25g.

- 1. Clin. Tri. J., (1966) 3: 591
- 2. Brit. J. clin. Pract., (1967) 21: 529 3. Ugeskrift for Laeger, (1966) 128: 204
- 4. Clin. Tri. J. (1966) 3: 529
 5. Paper read to the British
 Dermatological Association Meeting.
 Aberdeen, July 1968.





O LABORATORIES LIMITED, Hayes, Middlesex. Telephone 01-573 6224