

There then followed a section devoted to the executive council. This indicated the range of printed data supplied by the council with which the doctor was expected to become familiar. The data obviously could not be included but it was suggested that acquiring this information and developing a filing system was time well spent during the trainee year. However, the majority of forms supplied by the executive council were displayed. Many of these are self-explanatory, but additional notes as to their nature and use were supplied where necessary. Particular attention was drawn to medical certificates, prescriptions, maternity forms, registration of new patients, temporary residents and patients' notes. Finally, space was given to data relating to local social services.

Both my trainer and the area general practitioner tutor have felt the project worthwhile and made useful comments. It seems reasonable to bring the idea to the notice of as many trainers and potential trainers as possible. Is there a place for a trainee general practitioner working manual?

ALAN R. DUKE
Vocational trainee

23 Welbeck Avenue,
Aylesbury,
Buckinghamshire.

Infectious Mononucleosis

Sir,

I was very interested in Dr K. J. Bolden's study of the effects of corticosteroids in the treatment of infectious mononucleosis (February *Journal*).

I note that "all patients had concurrently a 12 day course of phenoxymethyl penicillin—250 mg q.d.s." Dr Bolden does not state if any of his patients developed an allergic rash in the course of this treatment and if so, was this antibiotic stopped?

I recently had a case in a young woman who was originally misdiagnosed as having a β -haemolytic streptococcal throat infection and who was given a course of ampicillin. On the ninth day she developed a severe rash, and this is apparently a likely complication in 50 per cent of cases given ampicillin.

R. K. McELDERRY

301 Shore Road,
Newtonabbey,
Co. Antrim.

REFERENCE

Bolden, K. J. (1972). *Journal of the Royal College of General Practitioners*, 22, 87-95.

General Practitioners and Social Workers

Sir,

As a social worker who both teaches and works with doctors in training in a family medicine programme, I wish to take issue with some statements in the *Journal*. The report entitled *General practitioners and social services departments*, helps to clarify the problems existing

between the medical and social work professions. The act of discussing problems of communication aids in the process of communication; hence, the article was most helpful.

I was particularly stirred by some remarks attributed to Dr Paul Freeling, general practitioner from Middlesex, which are refuted below:

(1) Patients *can* relate to more than one healer at a time, if they are taught and gently guided into bringing their problems to a team. Recently one cancer patient in our family health centre was introduced by her doctor to the social worker. A three way discussion about her physical, social and severe emotional problems ensued. Subsequently, she reached out for help alternatively either to the doctor or social worker, or as they were available. This kind of situation is common in our family health centre. We demonstrate to patients that any member of the team is interested and willing to be of help, while some members will have special skills to offer, such as, marital counselling by the social worker, health education by the nurse, and medical management by the physician. Patients learn to utilise all team members as a result of a casual, instructive approach.

(2) We have found great utility in formulating psychosocial-physiological diagnoses and treatment plans by use of inter-disciplinary team conferences. Families' total problems are elicited and discussed in an open, informal manner by physician, social worker, nurse and senior physician. Consensus is reached as to what plans should be undertaken, and which team member shall accomplish which task.

(3) Patients are helped immeasurably by having all resources "behind the same door", contrary to Dr Freeling's belief. Easy accessibility is one of the hallmarks of comprehensive family health care. When the patient's needs are kept foremost, and not the professional's need for self-esteem and status, close co-operation and good patient care are the result of such propinquity.

(4) In listing the resources of social workers, Dr Freeling may have been only facetious or simply uninformed in excluding case-work and counselling skills, knowledge of individual and family dynamics, knowledge of community resources of all kinds, to name only a few.

(5) In our experience, screening for emotional illness in patients and families although not always successful and certainly difficult to measure objectively, does work when patients and families are helped to look at potential trouble areas, and to accept counselling before a crisis has occurred; such as in pre-marital counselling, child management and discipline difficulties, and in many other common family developmental stages.

We advocate many of the issues raised: (a) common areas of training for all health professionals, and opportunities to study and work together so that knowledge of the skills and