

ment which, to quote our Royal President "sets out clearly and concisely just what general practice is all about"<sup>4</sup>.

There is, of course, no harm in all this, *provided the statements of Council reflect accurately the consensus of opinion of the college membership*. But do they? Does our Council submit its policy statements to critical scrutiny by the Faculties, and by the membership, before they receive the *imprimatur* of the College? Are we, in fact, sufficiently democratic? I do not think so.

As a College, we are unique in our faculty organisation. It is time now, I submit, to put this unique structure to better use. Much more power should be delegated from the centre to the Faculties; they should play an active part in formulating and monitoring college policy. It will, of course, take longer to reach decisions; this is a penalty paid by all democratic institutions, and it is infinitely preferable to the cabalistic rule of authoritarian regimes.

Secondly, let us consider the world of general practice outside our own tight little college membership. When the present criteria for membership were established, the distribution of that membership within the general-practice community was crystallised. We badly need to map out this distribution throughout the country. We shall find lush areas where a high proportion of the general practitioners are members of the College, and wide expanses of desert where the college representation is virtually nil. I believe that one of the top priorities of our College should be to fertilise these deserts.

For better or for worse, the time may come when a young doctor aspiring to the position of a principal in general practice will have to gain his M.R.C.G.P., in much the same way as a hospital consultant is now required to show some higher qualification as evidence of his training. If and when this happens, within a few decades *all* general practitioners will be members of the College, and the deserts I have spoken of will disappear. Can we afford to wait that long? I think not.

I believe we were mistaken to tighten our criteria when we did, so excluding permanently from membership very many general practitioners of high quality, and leaving the distribution of our membership permanently unbalanced. It is perhaps too much to ask that the policy be wholly reversed. But we are not going to tempt many of these experienced non-collegiate colleagues of ours, now in their forties and fifties, to submit themselves to examination before seeking entry to the College.

I would like to suggest that they be offered some special form of associate membership, granting them a status within the College not too far removed from that of a fully established member. If this idea be rejected, then I believe we should address ourselves to rectifying the maldistribution of our membership in some other way. We can-

not afford to continue to ignore this thorny problem.

CYRIL HART

Goldthorne,  
Stilton,  
Peterborough.

#### REFERENCES

- 1 Hunt, J. H. (1973). *Journal of the Royal College of General Practitioners*, **23**, 5-20.
- 2 Royal College of General Practitioners (1971). *Journal of the Royal College of General Practitioners*, **21**, 485-486.
- 3 Royal College of General Practitioners (1972). *Journal of the Royal College of General Practitioners*, **22**, 593-595.
- 4 Philip, H.R.H. The Prince (1973). *Journal of the Royal College of General Practitioners*, **23**, 65.

#### PRESCRIBING IN GENERAL PRACTICE

Sir,

Since the paper on the prescribing of barbiturates and their substitution by nitrazepam was written, further developments have taken place.

Six patients of the original 116 are no longer with us, four having died and two having moved away, but none of our patients are now on more than 10 mg of nitrazepam. None of them need added chlorpromazine, and of the 110 patients remaining under review, 54 are on no sedation, the others now being equally divided between a nightly dose of 5 mg and 10 mg. This represents a reduction in hypnotic use of 62 per cent.

F. O. WELLS

49 Christchurch Street,  
Ipswich IP4 2DF.

#### REFERENCE

- Will, F. O. (1973). *Journal of the Royal College of General Practitioners*, **23**, 164-70.

#### CHAPERONES

Sir,

Members of the Medical Defence Union have been advised that accusations of impropriety are made with such unpredictability, and such frequency, that examinations of females should always be chaperoned.

Many general practitioners, none the less, examine without. In the other scale are the points that many women prefer no third party present, that examinations, troublesome (by reason of disturbing staff), are that much more likely not to be done at all, and that the burden on staff is impossible to accept.

A consecutive series of 530 consultations of unaccompanied post pubertal women was analysed. The policy of the doctor was to examine where medically indicated.

Examinations with a chaperonable element were scored:

1. Genital: vaginal and rectal