

in the practice, a part-time general practitioner will attract allowances from the family practitioner committee of at least £4,185 per year, plus £700 per year for out-of-hours cover. She may also be eligible for a vocational trainee allowance of £855 per year. Thus any partnership in a non-restricted area can reduce its workload without incurring any additional expense by employing a part-time partner at a salary between £4,000 and £6,000 a year. Unfortunately, salaried partnership is not recognized by the FPC nor in common law. The salaried partner is still liable for the debts, tax, negligence, and acts of omission of the other partners. Thus, the part-time partner must ensure that she has a comprehensive contract, specifying hours of work, a stated percentage share of profits, an agreed minimum salary, and also insist that money should be retained in the partnership account for tax purposes.

The additional problem facing full-time women partners is that of maternity leave. Most partnerships have an agreement that, if one of the partners is absent owing to sickness, he shall, at his own expense, provide a locum for the practice after a certain specified period of absence. Failing this, he shall pay the remaining partners at the prevailing BMA rates for locums. No doubt a similar clause could be included to cover maternity leave.

But now the snags appear. First, the FPC will only pay sickness payments of up to £125 per week provided that a locum has been employed from outside the practice. (Maternity leave payments rules are very similar but are only paid for a maximum of 13 weeks.)

Secondly, reliable locums are hard to find. Thus, the sick or pregnant partner may have to pay the remaining partners at BMA rates which are at least £200 per week, without any reimbursement from the FPC.

Thirdly, if locums are not available, the additional workload for the remaining partners can be quite considerable, especially in a small partnership. In a society progressing gradually towards a 35-hour working week, the retention of free time and the non-interruption of holiday plans are more important to many doctors than an increase in salary.

Finally, most general practitioners are paying tax above the basic level. An additional income for locum duties will be lost to the tax man. It should be noted that locum rates are still below the average weekly earnings of most general practitioners.

I regretfully concluded that our female colleagues are liable to be exploited or suffer discrimination. How many general practitioners looking for a replacement partner in a practice whose

circumstances do not necessarily require a woman partner would choose a female replacement? Probably the answer is inversely proportional to the number of good male applicants. As the number of women graduates increases, there is an urgent need for the profession to negotiate better maternity leave payments with the Government.

K. C. HARVEY

Medical Centre  
Hay-on-Wye  
via Hereford.

## ROYAL COLLEGE OF PHYSICIANS OF IRELAND

Sir,

In your Medical News section (January *Journal*, p. 59) you quote Professor T. Counihan as saying that the recent dinner given by the regional vocational training committee at the Royal College of Physicians was the first occasion in history at which the College has welcomed representatives from the Royal College of General Practitioners.

The Chairman of Irish Council of the Royal College of General Practitioners has been a guest at the annual St Luke's day dinner of the Royal College of Physicians almost since the inception of Irish Council.

J. S. MCCORMICK  
*Professor of Community Health  
Trinity College*

196 Pearse Street  
Dublin 2.

## AMSPAR/RCGP WORKING PARTY

Sir,

At its meeting on 12 December 1979, the joint Working Party of AMSPAR and the RCGP agreed the following statement:

"The Joint Working Party of AMSPAR and RCGP is supporting the establishment of courses for middle management in eight centres commencing in September 1980. The Joint Working Party is to monitor and evaluate these courses which are designed for practice administrators/managers and intending administrators/managers."

The Working Party believes that such co-operation between our two organizations represents an important advance in the improvement of training facilities for ancillary staff.

S. O. OLIVER  
*Chairman,*

*Joint AMSPAR/RCGP Working Party*  
14 Princes Gate  
Hyde Park  
London SW7 1PU.

## SINGLE-HANDED PRACTICES

Sir,

The Rural Practice Sub-Committee of the BMA's General Medical Services Committee has asked me to prepare a report on the difficulties experienced by doctors who take over single-handed practices. I would be most grateful if any doctor who has experienced any such difficulty would kindly get in touch with me.

The report will contain no detail by which a doctor or practice could be identified. It will be used to try and influence Departmental policy towards more effective continuity of patient care when single-handed practices change hands.

B. D. MORGAN WILLIAMS  
10 St Michaels Road  
Claverdon  
Warwick CV35 8NT.

## GENERAL PRACTITIONERS AND CLINICAL PSYCHOLOGISTS

Sir,

We welcomed the section devoted to behavioural problems in general practice in the June 1979 issue of the *Journal*. It was particularly gratifying to note that two of the articles were written by clinical psychologists who reported close co-operation with general practitioners in the spirit of the role envisaged for us by Trethowan (1977).

We had developed a clinical psychology unit in a Newham district health centre (Hallam and Liddell, 1978) which provided a new service in an area in great need of psychiatric provision. However, after two years of operation, the number of referrals from general practitioners was lower than that from psychiatrists, social services, or self-referred clients. With the aim of expanding our service, we systematically attempted to contact every general practitioner listed in Newham. Half were contacted by telephone and half by pre-arranged visit during a three-month period. In this way we were able to discuss our service with 69 per cent of those contacted. Of the 31 per cent not contacted, 11 per cent no longer practised in Newham, five per cent were on sick or maternity leave, it was impossible to find a mutually satisfactory time for 10 per cent of the sample, and only five per cent actually refused to spare us the time. Six months after the beginning of the survey, general practitioners' referrals constituted the largest proportion of our patients.

We feel that the interchange of ideas