

before the smear is repeated.<sup>2</sup> This course of action although perhaps unwittingly dealing with the identifiable pathogens already mentioned, does not provide the opportunity to isolate, treat and contact trace chlamydial and/or gonococcal infection.

Another approach would be to follow the advice of Wilson's team and consider risk factors in deciding in whom to suspect an associated sexually transmitted disease: those aged under 25 years or those with a recent change in sexual partner.<sup>1</sup> Robinson and colleagues suggest this approach would be a suitable compromise in general practice as routine swabbing of women with inflammatory smears may not be cost effective.<sup>3</sup> It is also suggested that consideration should be given to a more aggressive policy for referring young women with a mildly abnormal smear to a genitourinary clinic for screening for sexually transmitted diseases.

Thus, although Wilkinson (review, August *Journal*, p.336) suggests high vaginal swabs for the follow up of persistent inflammatory smears, we urge the use of an additional investigation — endocervical swabbing.

RICHARD MENDELSON

Department of Genitourinary Medicine  
Birmingham General Hospital  
Steelhouse Lane  
Birmingham B4 6NH

MELANIE C MANN

The Birmingham and Midland Hospital  
for Women  
Showell Green Lane  
Sparkhill  
Birmingham B11 4HL

#### References

1. Wilson JD, Robinson A-J, Kinghorn SA, Hicks DA. Implications of inflammatory changes on cervical cytology. *BMJ* 1990; **300**: 638-640.
2. Kelly BA, Black AS. The inflammatory cervical smear: a study in general practice. *Br J Gen Pract* 1990; **40**: 238-240.
3. Robinson AJ, Mercey DE, Preston M, Bingham JS. Inflammatory cytology, infection and intraepithelial neoplasia. *Int J Sex Trans Dis* 1992; **3**: 123-124.

Sir,

Dr Clare Wilkinson must be congratulated on her review article (August *Journal*, p.336), and her extensive perusal of the literature. I certainly agree with her advice to repeat smears at three yearly intervals, and to take at least one smear from women aged over 65 years.

I would add two further recommendations. After the first smear, a second smear should be taken one year later, and three-yearly surveillance should then be started. Smears should be taken as soon

as patients are known to be sexually active, even before the age of 20 years. In my practice of 2200 patients abnormal smears needing treatment were detected in three patients under 20 years of age over a five year period.

GUS PLAUT

233 Boroughbridge Road  
York YO2 6AY

### Emergency contraception

Sir,

With reference to Dr Fluxman's letter on postcoital contraception (September *Journal*, p.394), I am writing to correct the point made about the Family Planning Association's leaflets on contraception. All the current Family Planning Association leaflets on contraceptive methods contain information about emergency contraception. The newest leaflet on male and female condoms contains a specific boxed section on emergency contraception, and this clearer presentation will be available in all future revisions.

The Family Planning Association is aware through its nationwide enquiry service (approximately 200 000 enquiries each year), that there is considerable confusion and misinformation among both consumers and professionals about emergency contraception. The Family Planning Association is currently working on a number of new initiatives to address this.

TONI BELFIELD

Family Planning Association  
27-35 Mortimer Street  
London W1N 7RJ

### Contraception clinic

Sir,

A child was waiting to cross the road, but kept standing without crossing. Somebody asked what she was waiting for and she replied 'My mummy told me to let a car pass by and then cross, looking right and left all the time. I have not seen any car pass by so how can I cross the road?'

A similar example of the literal interpretation of language occurred with a patient attending a special contraception clinic for young people asking for termination of a pregnancy, which had resulted from a burst condom. I asked if she knew about postcoital contraception. She replied that she had heard about it but did not know how to go about obtaining it. On asking if she had consulted her own general practitioner she replied that she could not as he knew her family well. I

then asked whether she had considered attending a family planning clinic. She replied 'A family planning clinic is for married people to plan a family. I am not married and I have nothing to plan yet. I will not have a family and my boyfriend also feels this way. There are too many people on this earth so it is silly to go on producing more children. I feel very angry that it is not clear that one can go to a family planning clinic for contraception when one does not need to plan a family.'

Family planning clinics should therefore be renamed contraception clinics so that it is quite clear what they provide.

R P YADAVA

Merton Surgery  
Merton Street  
Longton  
Stoke-on-Trent ST3 1LG

### Telephone consultations

Sir,

The practice of telephoning patients on the evening prior to their appointment, as advocated by Stainer (letter, August *Journal*, p.348), is unacceptable owing to the resulting breach of confidentiality.

A patient has the right to confidential treatment, and this applies to the fact that a consultation takes place as well as to the contents of that consultation. Once a telephone call has been made to the patient's home, that confidentiality is broken, as there is no control over who answers the telephone. Domestic telephones are often placed in living rooms or halls, and privacy is unlikely. Even if the patient declines to talk for fear of being overheard, there is no way this can be explained without in itself causing the patient to be compromised.

The timing and unexpectedness of the calls places the patient at a disadvantage. The early evening is a time of maximum chaos in most family homes, and the patient is also denied the time to mentally rehearse the presentation of his or her problem that the journey to the surgery or wait once there usually provides.

Even if none of these considerations applies, patients may well be left with the impression that their family doctor dislikes seeing them so much that he or she would rather spend the evening on the telephone trying to dissuade patients from coming, rather than spending the day actually attending them.

ANDREW GREEN

Hedon Group Practice  
Market Hill House  
Hedon  
Hull HU12 8JD