

a six-week intrauterine pregnancy and the second an asymptomatic ovarian cyst (7 cm diameter). Using a visual analogue scale to record the degree of discomfort caused by screening procedures, a bimanual vaginal examination did not cause more distress than breast examination or venesection. Ninety four per cent of women felt that a vaginal examination reassured them that all was well and 94.6% felt that it should be performed routinely, although 22.6% felt that it might put other women off attending.

In conclusion, it would appear that these women felt reassured that they had a healthy pelvis if a vaginal examination was performed at the same time as a cervical smear. As a group they were not unduly distressed by the examination and, as pathology may be detected, the opportunity to examine the pelvis digitally at cervical screening should not be missed. Either general practitioners should resume the practice of smear taking and combine this with a bimanual examination, or nurses responsible for cervical screening should be trained in performing the additional examination.

A MICHÈLE ARNOT

Huddersfield Royal Infirmary
Lindley
Huddersfield
West Yorkshire HD3 3EA

References

1. Department of Health and the Welsh Office. *General practice in the National Health Service. A new contract*. London: HMSO, 1989.
2. Osofsky HJ. Women's reactions to pelvic examination. *Obstet Gynecol* 1967; **30**: 146-151.
3. Haar E, Halitsky V, Stricker G. Patients' attitudes towards gynecologic examination and to gynaecologists. *Med Care* 1977; **15**: 787-795.

Alternative contracts in the NHS

Sir,

A letter from Julian Tudor Hart¹ quotes from an article by Professor Maynard, Dr Marinker and me, published in the *British Medical Journal* in 1986, about the possibility of alternative contracts in the National Health Service.² Unfortunately, Tudor Hart has misrepresented the sense of that paper.

First, in terms of historical accuracy, it was Enthoven who first proposed a market approach.³ His monograph was published the year before our article and we as professionals were examining what was then a new idea. The article to which Tudor Hart refers was the third in a series and its title raised as a question whether alternative contracts were viable.

In the first paragraph the exact words

were: 'Here we examine five further options.' This makes it clear that the authors were in no way at that time advocating a managed NHS market; on the contrary, our exact words were: 'We conclude that the development of a good practice allowance, along the lines described in our previous paper⁴ represents the best choice at this time for the public, the government, and the profession.' Furthermore we repeated this point in the final paragraph headed 'What option?': 'The opportunity exists to reform the present contract by adopting some variant of our own interpretation of the government's desire for a good practice allowance.'

'It is not impossible that a system similar to HMOs [health maintenance organizations] may become the pattern for a future national health service. Implementing a good practice allowance now would give us invaluable experience in setting standards for primary health care, monitoring performance, and reviewing and maintaining progress. If this does not happen it is likely that some stronger and more radical medicine will be administered to the system of general practice in the UK.'

In hindsight it now seems clear that had the profession felt able to accept a good practice allowance which the Royal College of General Practitioners had proposed in 1985⁵ and the government had offered the profession in 1986,⁶ it would inevitably have had to be negotiated with the representatives of general practice. This would have been a more professional arrangement than the subsequent contract, which was imposed on the profession in 1990 and indeed formed the 'stronger and more radical medicine' which we feared in 1986.

DENIS PEREIRA GRAY

Institute of General Practice
Postgraduate Medical School
Barrack Road
Exeter EX2 5DW

References

1. Hart JT. Burnout or in to battle? [letter]. *Br J Gen Pract* 1994; **44**: 96.
2. Maynard A, Marinker M, Gray DP. The doctor, the patient and their contract. III. Alternative contracts: are they viable? *BMJ* 1986; **292**: 1438-1440.
3. Enthoven AC. *Reflections on the management of the National Health Service*. London: Nuffield Provincial Hospitals Trust, 1985.
4. Gray DP, Marinker M, Maynard A. The doctor, the patient and their contract. II. A good practice allowance: is it feasible? *BMJ* 1986; **292**: 1374-1376.
5. Royal College of General Practitioners. *Quality in general practice. Policy statement 2*. London: RCGP, 1985.
6. Secretaries of State for Social Services, Wales, Northern Ireland and Scotland. *Primary health care. An agenda for discussion. Cm 9771*. London: HMSO, 1986: 14.

Patient involvement in medical education

Sir,

Jones and Oswald conclude that patients' attitudes are unlikely to present an appreciable barrier to increased medical student learning in general practice (letters, April *Journal*, p.184).

Patients in a recent qualitative study undertaken in our department were enthusiastic about involvement in medical education and some believed they could play a more active role, particularly in improving students' communication skills. All of the 26 patients interviewed expressed a willingness to take part in consultations where a student was present. A subsequent questionnaire study based on our results, carried out by medical students with 180 patients in Leeds confirmed our results (unpublished results). Both studies showed that patients perceive benefits for themselves and have few reservations.

Patients perceived the advantages of having a student present in the consultation as: helping to educate future doctors; learning more about their condition from the discussion between doctor and student; receiving a more thorough check up; interest in seeing the students' approach; and having someone uninvolved in their care to talk to (particularly if students visit patients at home). They saw the disadvantages as: potential embarrassment if the patient has a personal problem; feeling uncomfortable at being watched by the student; upsetting if the student appears uninterested; and the consultation may take longer.

Older patients (mostly those who had retired) in particular were enthusiastic about a more active involvement in student education. They had strong views about what students should learn, feeling that good listening, questioning and explaining skills were of paramount importance, along with the right attitude towards the patient (kind, respectful, not patronizing). This suggests that patients, in common with other potential community teachers,¹ would like a say in the sort of doctor that medical schools produce.

Thus, potential teachers, including health visitors, general practitioners, nurses and complementary practitioners would need to be involved in developing a curriculum which reflected the priorities, needs and ethos of primary care, rather than being expected to teach topics simply 'transferred' from the hospital. Realizing the potential for patient involvement in medical education clearly requires further work to identify the range of issues patients wish to see addressed, and appropriate teaching and learning methods.