

receiving regular prescriptions for digoxin, amiodarone, flecainide, verapamil or procainamide; patients with 'atrial' in their notes summary; patients with 'transient ischaemic attack' in their notes summary; and patients receiving regular prescriptions for warfarin or phenindione (Dindevan®, Goldshield).

From these data it was possible to define a group of patients with probable non-rheumatic atrial fibrillation and to determine their anticoagulation status. The groups with atrial fibrillation who were not receiving anticoagulation were further divided into three therapeutic groups as defined by Sweeney and colleagues:<sup>1</sup> patients aged less than 60 years with no complicating factors; patients aged less than 75 years without hypertension, congestive heart failure, thromboembolism or diabetes; and patients not in the two previous groups (the at risk group).

Each practice partner was given a list of patients in the at risk group and asked to decide, with the aid of patients' computer records, whether or not the patient was still in atrial fibrillation and if so whether to include or exclude the patient from being offered anticoagulation. Suggested reasons for exclusion included unsteady gait, tendency to falls, dementia, poor compliance, poor eyesight, excess alcohol consumption and active peptic ulceration (including maintenance on acid suppressing drugs).

Results revealed 224 patients with non-rheumatic atrial fibrillation (16 cases per 1000 patients) with an age range of 32-92 years and a mean age of 76.6 years. Of these 224 patients, 75 (33%) were receiving anticoagulation, with individual general practitioner rates ranging from 23% to 53%. Of the 149 patients not anticoagulated, six were aged less than 60 years with no complicating factors, 26 were aged less than 75 years with no additional risk factors and 117 were in the at risk group. Overall, practice partners included 59 of the 117 patients (50%) for consideration for anticoagulation, with inclusion rates for individual partners ranging from 17% to 86%.

This preliminary study indicates that with computerized records an audit of at risk patients is possible but this is time consuming (the audit took up to 20 hours of doctor time). The study produced a group of over 100 patients considered to be at risk and yet individual practitioner assessments as to which patients should be offered anticoagulation varied greatly despite having received the same suggested exclusion criteria. The resulting group of 59 eligible at risk patients would not all be expected to take up the offer of anticoagulation but even if half were to do so

this would represent a substantial workload in terms of surgery time and practice nurse time.

Despite the commendable effort of Sweeney and colleagues to define an at risk group of patients, the variation in individual practitioner's decision making may still prove to be an insurmountable hurdle in the effort to deliver care to a group in need.

D A SEAMARK

The Surgery  
Marlpits Road, Honiton EX14 8DD

### Reference

1. Sweeney KG, Gray DJP, Steele R, Evans P. Use of warfarin in non-rheumatic atrial fibrillation: a commentary from general practice. *Br J Gen Pract* 1995; **45**: 153-158.

## Urban community hospitals

Sir,

In their letter (June *Journal*, p.326) Hamilton and Round came to conclusions about the potential usage of an urban community hospital which do not appear to have been supported by the results of their study. Having found that 53% of the local general practitioners said that they would probably use such a hospital (82% for respite care, 75% for social reasons, 79% for elderly acute medical patients, 77% for observation, assessment and simple investigation, 56% for early hospital discharges following surgery, 59% for terminal care, and 56% for early hospital medical discharges), the authors conclude that 'an urban community hospital would provide services not now available, rather than being an alternative to district general hospital admission'.

A different conclusion from their study could have been that about half of the local doctors said that they would use an urban community hospital for a wide range of patients, many of whom currently occupy beds inappropriately in the local district general hospital.

RICHARD J COOK

Galen Lodge  
Eastfield Road  
Ross-on-Wye, Herefordshire  
HR9 5AN

Sir,

Cook agrees with the conclusion of our letter (June *Journal*, p.326) that an urban community hospital would generate a reduction in work for the local district general hospital. Where we do not agree is in the magnitude of the effect. We believe it would be small.

Admissions to community hospitals come from three main sources: early district general hospital discharges, direct admissions which would otherwise have gone to the district general hospital, or patients who would never have been sent to hospital at all. Our study shows general practitioner preference for this last group (respite care and social admissions), and least interest in the first group.

The rate of 'inappropriate' admissions to a district general hospital, usually quoted as around 15%, has always been hospital-defined, and is approximately halved when general practitioners' opinions are sought.<sup>1</sup> Such a false positive rate for emergency hospital admission of under 10% is excellent, especially when one considers the potential consequences of failing to admit a patient to hospital when admission is needed. The myth that there is a large pool of inappropriate admissions that can be redirected to a community hospital does not stand up to examination.

Nonetheless, an urban community hospital may still have a role — it was supported by 49% of general practitioners, in some cases strongly. It cannot be seen, however, as a quick-fix solution for rising medical admissions to district general hospitals.

WILLIAM HAMILTON

Barnfield Hill Surgery  
13 Barnfield Hill, Exeter EX1 1SR

ALISON ROUND

Department of Public Health  
Exeter and North Devon Health Authority  
Exeter EX1 1PQ

### Reference

1. Coast J, Inglis A, Morgan K, et al. *The hospital admissions study. Final report.* Bristol: Health Care Evaluation Unit, University of Bristol, 1994.

## Clinical guidelines

Sir,

The paper by Conroy and Shannon (July *Journal*, p.371) was a masterly review of the pitfalls that await the implementation of clinical guidelines.

We recently conducted a critical review of clinical guidelines, assessing the evidence to see whether guidelines have had any statistically significant effect on the outcomes of patient conditions in primary care, as opposed to merely changing the process by which family doctors deliver care to their patients (Worrall G, Chaulk P. Hope or experience? A critical appraisal of the effects of clinical guidelines on patient outcomes in primary care, *J Fam Pract* 1995; under review). We