

## REFERENCES

1. Horn CR, Cochrane GM. Colour coding for bronchodilator inhalers. *Lancet* 1986; **1(8473)**: 165.
2. Minerva. *BMJ* 1992; **305(6853)**: 594.
3. Maxwell D. Distinguishing inhalers to aid blind people. *BMJ* 1992; **305(6863)**: 1226.
4. Partridge M. Coloured inhalers. *BMJ* 1992; **305(6858)**: 890.
5. Jones KP. Guidelines on the management of asthma. *Thorax* 1993; **48(10)**: 1050.

DOI: 10.3399/bjgp10X515449

## Steroid treatment cards: patient safety remains at risk

The *British National Formulary* advises that all patients prescribed corticosteroids for more than 3 weeks are issued with a steroid treatment card that they carry on their person at all times.<sup>1</sup> An audit was carried out to determine if these guidelines are being met in patients receiving palliative care at Willow Wood Hospice. A similar survey carried out by Zeppetella in 1998, and published in a letter to this journal, revealed that 43% of patients referred to St Joseph's Hospice, London, did not have a steroid treatment card,<sup>2</sup> and I was interested to determine if practice had now improved.

Seventeen patients were included in my audit; all were prescribed corticosteroids orally, either dexamethasone or prednisolone, which they had been taking for over 3 weeks. Nine patients (53%) did have a card; eight (47%) did not. Out of those with a card, only two (22%) admitted to carrying it at all times.

Steroids are prescribed for many indications in palliative care, including the relief of non-specific symptoms, such as anorexia, fatigue, and cachexia syndrome.<sup>3</sup> However, their long-term usage is associated with serious side effects, such as Cushing's syndrome and an increased susceptibility to infection. Prolonged use may cause suppression of the hypothalamic-pituitary-adrenal axis resulting in adrenal insufficiency with risk of adrenal crisis, cardiovascular collapse, or even death in the event that steroids

are stopped suddenly, including during anaesthesia.<sup>4</sup> The steroid treatment card acts as a reminder to patients about the potential side effects and dangers of abrupt cessation, as well as a warning to healthcare professionals about patients' medication status: ultimately it aims to ensure patient safety.

This audit demonstrated that the British National Formulary guidelines are not being met in 2010: almost 50% of patients did not have a steroid treatment card and of those with a card, few always carried it. These findings reflect those of Zeppetella's survey, suggesting that this may be a long-standing, widespread problem. Steroid treatment cards should be distributed to and carried by patients on long-term steroids: this issue needs to be recognised and steps taken to improve clinical practice in this area.

### Elizabeth Rusby,

Fourth-year medical student, Low Winsley House, Burnt Yates, Harrogate, HG3 3ES.

E-mail:

Elizabeth.Rusby@student.manchester.ac.uk

## REFERENCES

1. Martin J. *British National Formulary* 59. London: BMJ Group, 2010.
2. Zeppetella G. Steroid treatment cards. *Br J Gen Pract* 1998; **48(432)**: 1436.
3. Hardy JR, Rees E, Ling J, et al. A prospective survey of the use of dexamethasone on a palliative care unit. *Palliat Med* 2001; **15(1)**: 3–8.
4. Jabbour SA. Steroids and the surgical patient. *Med Clin North Am* 2001; **85(5)**: 1311–1317.

DOI: 10.3399/bjgp10X515458

## Patients' perceptions of hypertension management: a concern for healthcare practitioners

Patients' attitudes towards medication have a major impact on disease management, but this is often neglected. A large body of literature has been published from developed nations on the

evaluation of non-adherence in chronic disease management and in patients with hypertension in particular.<sup>1–2</sup> However, there is a paucity of data in the context of hypertensive patients' perspective from developing countries, to gather in-depth information towards hypertension management.

In the context of developing nations like Pakistan, health-seeking behaviour always occurs in the context of medical pluralism, where the patient uses different systems of healing. Patients make independent assessments regarding the use of modern medicines. Orthodox medications do not have such dominance over the population as seen in the western world.<sup>3–4</sup> In addition, indigenous healing systems, particularly hikmat (treatment with herbs) and spiritual healing, are quite prominent in these populations. There is no legal or official acknowledgment as far as the spiritual healing is concerned; but it is still the treatment of choice for the majority of the population. Insufficient information and understanding emerges as major barriers associated with all aspects of medicine use. Knowledge about the disease and drugs used for its treatment is apparently poor and in return produces a great impact on patients' adherence to medication. Another issue that lies ahead for health practitioners is that discussions of the advantages and disadvantages of drugs in ways that are relevant for individual patients is rarely seen in practice.

What is needed at this point is a complete revision of the practices that are currently being employed in the healthcare system. Individuals must be educated not only on the risk factors, presenting features, and complications of hypertension, but also about the benefits of medications on treatment outcomes and quality of life. Patients may need to be educated about the differences between curing hypertension and treating it with medications. Efforts to educate the public about lifestyle modifications in the prevention of hypertension and other cardiovascular diseases may also be of great importance. Advancement in