prescribing in primary care continues to elicit debate and regular research contributions.^{1,2} I carried out a single practice survey of depression management at a practice in south east London. The surgery's electronic records were searched for patients who had been treated for a diagnosis of depression, with or without anxiety between the dates of 1 January 2000 to 21 February 2007. These subject-records were then searched for aspects of management that had been put in place with demographic details and management recorded. A total of 91 patients were returned from the search population by the electronic records database search engine, this amounted to 4.5% of the 2031 registered patients. The ratio of females to males in the cases was 2:1. The majority of patients were between the ages of 30 and 49 years, and more than three-quarters of the sample were of either of black or white ethnicity, with Asians making up 7% of the cohort. In terms of management, most (86%) of the patients had been treated with an antidepressant at some point. Interestingly, the most commonly prescribed agent was dothiepin, a tricyclic, which was prescribed in 27 of the cases.

This finding was interesting in relation to the NICE guidelines on management of depression, whose recommendations at the time of the study were that dothiepin should be commenced only by mental health professionals. The next most commonly used drug was the SSRI, paroxetine, used for 23 cases. Other agents used, in order of reducing frequency were: fluoxetine, venlafaxine, citalopram, mirtazepine, sertraline, and escitalopram. Of four patients on venlafaxine, two patients were found to have been in contact with secondary care, with two patients having no electronic record of secondary service contact. Thirty-six of the 91 patients received some form of further care after their initial point of contact with the GP, 19 receiving an outpatient psychiatric referral, eight being referred for counseling, and the rest being referred either to a CRISIS team, PCT psychotherapy services, Victim Support, Gamblers Anonymous, or an outpatient

liaison psychiatry clinic. Overall, it appears that the pattern of UK antidepressant prescribing, even in relatively well resourced areas, remains patchy.

Vishal Bhavsar,

*Academic Clinical Fellow, 16 Colin Court, Streatham Hill, London, SW16 1LJ.
E-mail: vishal.2.bhavsar@kcl.ac.uk*

REFERENCES

1. Cameron IM, Lawton K, Reid IC. Appropriateness of antidepressant prescribing: an observational study in a Scottish primary-care setting. *Br J Gen Pract* 2009; 59(566): 644–649.
2. Morrison J, Anderson MJ, Sutton M, *et al.* Factors influencing variation in prescribing of antidepressants by general practices in Scotland. *Br J Gen Pract* 2009; 59(559): e25–31.

DOI: 10.3399/bjgp10X539353

Academic general practice

I really appreciated the Back Pages paper on academic general practice.¹ It emphasises not only the history but the importance of academic general practice. John Howie warns that the increase in renaming general practice as primary care could mean that general practice may not be a GP-led service in the future. Also, I would strongly underline what Howie writes, 'academic general practice has contributed significantly to the evolving understanding of the relationships between medicine and society'.

I think that this understanding is a key political point to spread worldwide of the importance of teaching in undergraduate academic general practice. National and international GP organisations should help political developments in other countries, at least the European ones.

Howie mentioned some international organisations also naming the Leeuwenhorst Group. This became EURACT (European Academy of Teachers of Family Medicine). EURACT is really the most active network (and academy with a legal body) in the WONCA context. During these years EURACT worked with national representatives from 40 different European countries to write key basilar documents

for family medicine.² They are: the European Definition, the EURACT Educational Agenda, the EURACT Performance Agenda, the Statement on Selection of Teachers and Practices, and the Checklist for Course Organizers. EURACT Educational Agenda, with its core competences, describes the content and the way for teaching and learning in a general practice context so effectively that it was used by the RCGP in its official documents and is used in other European countries. This is not the same in all countries and I think it is a duty for the strong national and international academies and organisations of family medicine to spread academic general practice in each country with the obvious positive consequences on family medicine and the population.^{3,4,5}

As the EURACT Basic Medical Education Committee, we are now at the point of publishing some research, based on the Delphi process, a minimal teaching core curriculum in general practice which will hopefully be useful in introducing a minimal homogeneity in programmes, but mainly to help less established countries open general practice departments and courses for all the students in medicine.⁶

Francesco Carelli,

*Professor of Family Medicine, University of Milan, EURACT Council Director of Communications, Chair BME Committee.
E-mail: carfra@tin.it*

REFERENCES

1. Howie J. Academic general practice: reflections on a 60-year journey. *Br J Gen Pract* 2010; 60(577): 620–623.
2. <http://www.euract.org> (accessed 8 Oct 2010).
3. Carelli F. The current status of family medicine in Europe. *LJPC* April 2009: 1–2.
4. Basak O, Yaphe J, Spiegel W, *et al.* Early clinical exposure in medical curricula across Europe: an overview. *Eur J Gen Pract* 2009; 15(1): 4–10.
5. Soler JK, Carelli F, Lionis C, Yaman H. The wind of change. After the European definition: orienting undergraduate medical education towards general practice/family medicine. *Eur J Gen Pract* 2007; 13(4): 248–251.
6. EURACT BME Committee. *Family medicine/general practice undergraduate teaching in Europe: a Delphi study to determine a minimal curriculum*. Oral presentation 499. WONCA Europe Conference in Malaga, October 2010.

DOI: 10.3399/bjgp10X539362