9. Little P, Dorward M, Warner G, *et al*. Importance of patient pressure and perceived pressure and perceived medical need for investigations, referral, and prescribing in primary care: nested observational study. *BMJ* 2004; **328**(7437): 444.

DOI: 10.3399/bjgp12X656757

One-dimensional learning: detrimental to the Situational Judgement Test?

Medical students acquire knowledge, apply theory, and develop skills during their undergraduate training. At medical school, formal teaching sessions tend to focus on the communication between doctor and patient, often with only passing reference to more complex everyday scenarios in which doctors find themselves. There are many situations and diverse interactions demanding effective communication in which the future clinician may be undertrained. These could be with, but are not limited to, members of the patient's family or other healthcare professionals and through different media such as by telephone, email, or with legal personnel. The nature of medical training regarding communication is too one-dimensional. The importance of multiprofessional teamwork is emphasised throughout our training, but opportunities to consolidate this theory in real-life scenarios are all too rare. As a newly qualified doctor or when later holding a position of increased responsibility, a lack of experience of real-life professional interactions may contribute to difficulties with colleagues and lead to increased stress in the workplace. For example it would be helpful to provide strategies for confrontation resolution when in challenging situations. At the other end of the spectrum, introducing students to the art of communicating a referral to another healthcare professional would be beneficial. The focus during training needs to broaden.

Difficult decisions may have to be made in parallel with communicating in unfamiliar settings. Thus the lack of experience of communicating in these situations may affect the development of clinical decision-making skills; skills that are indispensable to a competent clinician. For instance, when faced with the uncertainty that often permeates general practice, it is imperative to both arrive at and communicate a well-

reasoned, decisive plan of action. This can be complicated further if the setting of the communication is foreign to the clinician. GPs familiar with personal and emotional experiences of these situations will be better equipped to deal with the challenges posed by time, language, cultural, and sensitive issues, and the patient's agenda, while still establishing rapport and trust. The lack of exposure to such scenarios, whether real life or simulated, will surely be to the detriment of future skills as a clinician and also in the short term potentially, in achievement in the Situational Judgement Test.

Jason K Ferdjani,

Brighton and Sussex Medical School, BSMS Teaching Building, University of Sussex, Brighton, East Sussex, BN1 9PX.
E-mail: J.Ferdjani1@uni.bsms.ac.uk

Amour BU Patel,

Brighton and Sussex Medical School, BSMS Teaching Building, University of Sussex, Brighton, East Sussex, BN1 9PX.

Soram RU Patel,

Brighton and Sussex Medical School, BSMS Teaching Building, University of Sussex, Brighton, East Sussex, BN1 9PX.

DOI: 10.3399/bjgp12X656766

Introducing an old teaching tool to general practice

The term 'doctor' comes from Latin for 'to teach'. In general practice, this can involve teaching patients, medical students, trainees and other health professionals. As part of my medical school rotations, I recently shadowed a doctor who was particularly keen to teach both his patients and medical students. I thought his method of drawing out diagrams for patients was particularly helpful to a patients' understanding of what was happening in their bodies. However, when drawing on paper, it is necessary to show the patient what is being drawn in the correct orientation. This involves positioning oneself next to the patient, that can potentially make some patients uncomfortable. This can be compounded by the doctor inviting medical student(s) to also learn from the drawings at the same time as

the patient, creating an unintentional huddle mid-consultation.

I suggest that doctors who are keen to teach in this way consider investing in a whiteboard. Whiteboards have largely replaced traditional chalk and blackboards. Marker pens are used to write on the smooth surfaces of whiteboards, making them easy to erase. A quick search on a popular shopping website revealed that a 60 x 40 centimetre whiteboard could be bought for as little as £5. An investment in a whiteboard may therefore be a way to reduce paper costs in the long term. They may also be more environmentally-friendly than using paper. Importantly, as they can be placed on a part of the wall visible to all, they can be used to teach students and patients simultaneously without unintentional invasion of personal space.

However, careful thought regarding placement of the whiteboard is necessary to avoid introducing a lecturer-student dimension to the already complicated patient-doctor relationship. This may be the case if the doctor has to stand up to write on the board. I suggest placing the board in such a way that the doctor can write on it while sitting down so they remain on the same level as the patient.

Nigel Mabvuure,

Brighton and Sussex Medical School, Phase 2, Audrey Emerton Building, Eastern Road, Brighton, BN2 5BE.
E-mail: N.Mabvuure1@uni.bsms.ac.uk

DOI: 10.3399/bjgp12X656775

Are GPs unsympathetic to infertile women?

Are GPs unsympathetic to infertile women? The National Infertility Awareness Campaign (NIAC) published results from their most recent patient survey on 27th August 2012, based on responses from 456 patients collected in 2011. While 78% of responders indicated their GP was sympathetic and helpful, almost 50% also said they felt their GP lacked the necessary knowledge of infertility and treatment options available to provide an effective service. Clare Lewis-Jones, Chief Executive of NIAC, suggested this could explain why some patients (22%) with fertility issues perceive their GPs as unsympathetic.¹

Dr Clare Gerada, chair of the Royal