Kahneman's central point is that we use two different kinds of thinking. 'System 1' thinking works fast, using intuitive mental short-cuts. 'System 2' thinking is slower, rational, and deliberative. In his very readable book² Kahneman demonstrates that System 1 is what people use most of the time, even highly-educated, numerate people like the students who volunteered for his experiments: they were unaware of jumping to conclusions, and shocked when later shown the errors this had led them to make. When given increasingly complex tasks that forced them to switch into using System 2 thinking, they reported that this felt like hard work, and Kahneman observed highly-consistent physiological changes associated with this feeling.

Explaining things to people and helping them make choices is central to our job as GPs. It is high time we stopped acting as if people were likely to respond by engaging in System 2 thinking, or to be comfortable when given tasks that require it. For example, the whole enterprise of communicating risk and facilitating shared decision-making is aimed at an imaginary patient who weighs up the facts and her doctor's advice alongside her individual preferences in order to compute a rational choice. We should acknowledge that this patient is a very rare bird, and begin working out how best to interact with System 1 thinkers, as Gardner *et al* suggest.

While we are about it, our own learning and teaching would be improved by recognising how seldom we use System 2 thinking ourselves, especially within 10-minute consultations. Healthy (consultation) habits are what keep us safe most of the time, but the drawbacks of intuitive short-cuts explain many of our mistakes, mistakes that often seem surprising when examined retrospectively using System 2 thinking. We should all read Kahneman!

Louisa Polak,

GP and PhD Student, North Hill Medical Group, Colchester, CO1 1DZ.

E-mail: louisa.polaki@lshtm.ac.uk

REFERENCES

1. Gardner B, Lally P, Wardle J. Making health habitual: the psychology of 'habit-formation' and general practice. *Br J Gen Pract* 2012; DOI: 10.3399/bjgp12X659466.
2. Kahneman D. *Thinking, fast and slow*. London: Penguin, 2011.

DOI: 10.3399/bjgp13X662966

Questionnaire survey of the effect of calorie labelling on consumers' cold drink purchases

In their systematic review of the health impacts of neurolinguistic programming (NLP), Sturt and colleagues found no evidence of benefit in 'weight challenged adults'.¹ As GPs know only too well, obesity is a major health problem in the UK. The UK Department of Health recently launched the Responsibility Deal programme in which organisations pledged to provide calorie labelling on their products.² In September 2012, for a medical student project, we conducted a confidential questionnaire survey of customers sitting in Eddie Wilson's café at St. George's, University of London, to investigate whether their choices of cold drinks were influenced by calorie labelling.

The response rate was 92% (92/100). The mean age of participants was 25 years (range 18 to 54) and 60% were female. They describe their ethnicity as white 55%, Asian 27%, black 11%, or other ethnic group 7%. On average the 37% (34/92) of responders who said they read the calorie content chose a lower calorie drink than those who didn't: mean (SD) calorie content 62.3 kcal (85.7) versus 103.2 kcal (83.7), $P = 0.03$. The 26% (24/92) of people who said they were influenced by the labelling chose even lower calorie options, often diet drinks or water: mean 27.3 kcal (55.3). On average women chose lower calorie drinks than men: 71.5 kcal (83.2) versus 112.8 kcal (85.5), $P = 0.02$. However there were no differences between the calorie content of drinks in people with BMI ≤ 25 and >25 (based on reported height and weight).

A recent study suggested that in New York, US, a policy of banning super-sized sugar-sweetened drinks (>16 fl oz) in fast-food restaurants could reduce calorie consumption per consumer by 63 kcal (95% CI = 61 to 66).³ Similarly in England, enforcing a minimum price per unit of alcohol could help to reduce alcohol consumption. Around one-third of responders from our health education institution said they read the calorie labelling and were influenced in selecting a lower calorie drink. In contrast to NLP, where evidence is of limited quantity and quality,¹ it is possible that changes in government policy could contribute towards tackling the obesity epidemic.

Christianne Dominise,

Medical Student, Population Health Sciences and Education, St George's, University of London, London, SW17 0RE.
E-mail: m0901443@sgul.ac.uk

Pippa Oakeshott,

Reader in General Practice, Population Health Sciences and Education, St George's, University of London, London, SW17 0RE.

REFERENCES

1. Sturt J, Ali S, Robertson W, *et al*. Neurolinguistic programming: a systematic review of the effects on health outcomes. *Br J Gen Pract* 2012; **62(604)**: 580-581.
2. Dumanovsky T, Huang CY, Nonas CA, *et al*. Changes in energy content of lunchtime purchases from fast food restaurants after introduction of calorie labelling: cross sectional customer surveys. *BMJ* 2011; **343**: d4464.
3. Elbel B, Cantor J, Mijanovich T. Potential Effect of the New York City Policy Regarding Sugared Beverages. *New Engl J Med* 2012; **367(7)**: 680-681.

DOI: 10.3399/bjgp13X662975

Not just another primary care workforce crisis

It seems that the general practice workforce is being managed by insurance companies.

A colleague, sick with clinical depression following marital breakdown and disruption within her practise, is subject to certain GMC undertakings that limit future practice. She has met their requirements to sit and pass the RCGP clinical examinations and has successfully applied for a supervised GP Registrar post and a place on an Induction and Refresher (I&R) scheme, all supported by her last GP Dean, her future Dean, and the director of the training practise that she will be joining. She can now proceed with her refreshed career, but a place on the PCT performers list is conditional on evidence of professional indemnity.

And here is the rub; the prospective employer has a group indemnity but their insurance company, incorrectly, gave advice that an individual application would have to be made. This was refused, and a complaint was followed by a review 1 month later, and the application was again refused. Two further insurance companies refused cover

outright despite being in receipt of GMC approved PDP plans and references from her new employer. A fourth company has agreed cover at the cost of £16 000 per annum.

The first three companies have been anonymously approached to provide their assessment criteria for return to practice applicants and copies of their policies for those on I&R schemes. None have yet responded.

She has now been waiting a month for a comment from the GMC on this position; their continuing delay in responding perhaps indicates that they do not have a policy of dialogue with medical indemnity companies, or to their own customer service commitment, that promises a reply within 10 days.

The country is desperately short of doctors experienced in primary care and particularly short of mature doctors who have met the most recent and rigorous standards for practice through their professional body, the RCGP, and its Returners Scheme.

Surely it cannot be true that medical indemnity companies have more influence than the GMC on return to practise for the UK's GP workforce.

The Author can be contacted via the BJGP office

This letter was sent to the General Medical Council, the Medical Defence Union and the Medical Protection Society all of whom declined to comment.

DOI: 10.3399/bjgp13X662984

As the government launches its consultation on medical performers list (MPL) regulations, it seems timely to review the assessment of those returning to UK general practice after a period away from clinical practice. An issue that causes considerable controversy.¹

The evidence of specific patterns of deterioration of clinical skills after absences from the workplace is thin, and not directly applicable to primary care.² Practitioners vary both in their baseline clinical knowledge, and in the rate of deterioration during their time away.

Severn Deanery launched a 'Returners' Scheme' in 2007.

Entry is restricted to those out of practice in excess of 2 years. It includes a structured interview with a senior GP educator, and a national computer-marked knowledge test and simulated surgery. Each is standard set by experienced GP trainers following international best practice. Applicants are

Table 1. Severn Deanery induction and refresher scheme activity — 2007–2012

	Time away from UK general practice, years		
	2–5	5–10	>10
Total Number Assessed	17	17	13
Competent	5	1	0
Shortened scheme	7	6	1
Full Scheme	4	9	4
Fail	1	1	8

MPL = medical performers list. Competent = passed comfortably with good scores in all domains. Letter of support for return to MPL. Shortened scheme = borderline result. Short, typically 6–13 week, attachment to training practice with workplace based assessments (WPBAs) before letter of support for return to MPL. Full scheme = weak scores with extensive learning needs. Full training programme (6 months) and comprehensive WPBAs before letter of support for return to MPL. Fail = learning needs beyond the scheme's capacity and patient safety standards for supervised practice.

required to achieve a minimum entry score before progressing (Table 1).

A recent review of the scheme provides some evidence of the educational value of the programme.³

Primary care organisations are responsible for ensuring that the competency of practitioners admitted to their MPL. Our data suggests that many are not, after 2 years away from the workforce. This proportion increases with additional time away.

Jim Morison,

Severn Deanery, Deanery House, Vantage Office Park, Old Gloucestershire Road, Hambrook, Bristol, BS16 1GW.

Bill Irish,

Severn Deanery, Deanery House, Vantage Office Park, Old Gloucestershire Road, Hambrook, Bristol, BS16 1GW.
E-mail: Bill.Irish@southwest.nhs.uk

Paul Main,

Severn Deanery, Deanery House, Vantage Office Park, Old Gloucestershire Road, Hambrook, Bristol, BS16 1GW.

REFERENCES

1. Berger D. Not just another primary care workforce crisis. [letter]. *Br J Gen Pract* 2012; **63**(598): 236.
2. Bonrath EM, Weber BK, Fritz M, et al. Laparoscopic simulation training: testing for skill acquisition and retention. *Surgery* 2012; **152**(1): 12–20.
3. Kahol K, Ashby A, Smith M, Ferrara JJ. Quantitative evaluation of retention of surgical skills learned in simulation. *J Surg Educ* 2010; **67**(6): 421–426.

4. Morison J, Main P, Irish B, Curtis A. GPs' experience of the Returner (Induction & Refresher) Scheme in Severn: a qualitative study. *Educ Prim Care* 2012; **23**(4): 255–262.

DOI: 10.3399/bjgp13X662993

Work and health reforms unattainable?

The recent editorial encouraging the integration of occupational health and primary care¹ was inspirational but seems a long way from any reality in England.

When Dame Carole Black reported her findings and proposals on work and health it was difficult to resist the rhetoric. Work is good for you in determining your self and worth as long as the work is 'good'.

Charles Handy² describes the characteristics of organisations and individuals in organisations using four dimensions, four Greek Gods as their embodiment, and how the cultures they represent interact and have impact upon individuals within the organisation. He describes how social values and expectation have changed from the post war acceptance of authoritarian structures/organisations (The God Apollo). Increasingly we all want to be valued and have a share in the 'fruits' of our collective effort (The God Dionysius).

In the context of health and work, he describes how the great efficiency driver of Apollo, the bureaucratic organisation, so devalues the members that they resort to disruptive behaviour and demonstrate ill health. In Apollonian organisations, where members have little control or discretion about the work, the impact cannot be good,