All letters are subject to editing and may be shortened. General letters can be sent to bjgpdisc@rcgp.org.uk (please include your postal address for publication), and letters responding directly to *BJGP* articles can be submitted online via **eLetters**. We regret we cannot notify authors regarding publication. For submission instructions visit: bjgp.org/letters

Editor's choice

Bad medicine: resilience

I read Dr Spence's polemic against resilience support and training with a mixture of sadness and disappointment.¹ I am sad that he had a poor experience of training. It is terrible that he felt no support or worthwhile mentoring from the old style 'firms' who trained him. This is very different from the experience I had, in the same country and era, which was enhanced by many excellent role models and at its best created a real feeling of belonging.

I am disappointed at the offhand dismissal of a whole area of psychological theory and a simplistic conclusion that ignores the complexities of the interactions between life, work, and the individual.

Clearly improvements in the workload, working environment, and organisational culture of health care are vital in protecting staff from stress. Having said that, bad things will still happen in the quietest, cleanest, and best-run health centres and hospitals. Patients die unexpectedly and make complaints no matter how good your job, home life, and upbringing. It is vital that we make efforts to allow doctors to handle these events, which are an unavoidable part and parcel of our lives.

Resilience isn't hardness, stiff upper lips, cynicism, or emotional blunting. It's talking, openness, maturity, and professionalism. Good medicine.

Gregory C Jones,
Consultant/Associate Postgraduate
Dean, NHS Education Scotland.
E-mail: g.jones3@nhs.net

REFERENCE

1. Spence D. Bad medicine: resilience. *Br J Gen Pract* 2016; **66**(653): 621. DOI: <https://doi.org/10.3399/bjgp16X688201>.

DOI: <https://doi.org/10.3399/bjgp17X688957>

The value of clinical examination

The article 'Bad medicine: clinical examination' might have made me chuckle at the imaginative use of Harry Potter references but it did not persuade me to change my clinical practice.¹ The author seems to equate disdain for the perceived hierarchical teaching and comment from medical schools and royal colleges as proof that clinical examination has no relevance.

He suggests that we 'embrace technology' and cites how CT and MRI have blown away the pomposity of consultants; both these things may have merit but they do not mean that examination is without value. He is right that it is dangerous to rely on clinical examination. But a good clinician will take a full history and enhance that with appropriate examination; there is no absolute reliance. As a GP, I don't have access to investigations during a consultation; examination helps to determine whether an illness can be managed in primary care or needs to be referred for further investigation or secondary care consultation and whether such referral should be routine or urgent.

He again highlights intimate examination such as pelvic examination (bimanual examination +/- visualisation of the cervix).² The evidence that this examination is of no benefit to asymptomatic women is clear, but to suggest that pelvic examination has no place in the management of symptomatic women is cause for concern. There has been little research conducted into the role of pelvic examination in primary care so evidence is limited, but what evidence there is supports the use of pelvic examination prior to referral.³ There is nothing illogical about using pelvic examination to determine if a patient's postcoital bleeding is caused by a cervical ectropion or polyp or a possible cervical cancer.

Pauline Williams,
GP/CSO Clinical Academic Training Fellow,
University of Aberdeen.
E-mail: pauline.williams@abdn.ac.uk

REFERENCES

1. Spence D. Bad medicine: clinical examination. *Br J Gen Pract* 2016; **67**(654): 31. DOI: <https://doi.org/10.3399/bjgp17X688693>.

[org/10.3399/bjgp17X688693](https://doi.org/10.3399/bjgp17X688693).

2. Spence D. Bad medicine: gynaecological examinations. *BMJ* 2011; **342**: d1342.
3. Vandborg MP, Christensen RD, Kragstrup J, *et al*. Reasons for diagnostic delay in gynecological malignancies. *Int J Gynecol Cancer* 2011; **21**(6): 967-974.

DOI: <https://doi.org/10.3399/bjgp17X688981>

Bad medicine: clinical examination

Des Spence questions the value of clinical examination and claims that '... the most common defence is that doctors need these "skills" to work in low-income countries'.¹ This is untrue. Practitioners, when making a clinical decision with patients, need to incorporate all the tools available to them, including the patient's background, a focused history, and a clinical examination. The thought processes of many junior hospital doctors are echoed by Dr Spence. As a result departments become saturated, tests beget tests, and decisions spiral ever upwards. One of the first lessons of a new GP registrar is to 'unlearn' this mindset.

A reassuring history and *negative* examination allow us to use time. Conversely, a worrying history especially with demonstration of hard physical signs guides us to the next stage. The skill then becomes 'when to do what'. Patients have a right to expect their GP to recognise serious illness: the 'rare but important' if you like. In adults the acute physicians now use a generic National Early Warning Score to spot such cases. Guess what? The parameters are clinical.² Please don't throw your stethoscopes and thermometers away yet. When the chips are down you may one day need them.

Michael Houghton,
GP, Leagram, Nr Preston, Lancashire.
E-mail: mike.houghton@nhs.net

REFERENCES

1. Spence D. Bad medicine: clinical examination. *Br*

J Gen Pract 2016; **67(654)**: 31. DOI: <https://doi.org/10.3399/bjgp17X688693>.

2. Royal College of Physicians. *National Early Warning Score (NEWS): standardising the assessment of acute illness severity in the NHS*. Report of a working party. London: RCP, 2012.

DOI: <https://doi.org/10.3399/bjgp17X688969>

Colchicine in overdose

We read with interest the letter from Lloyd warning of the danger of colchicine overdose¹ and agree that this can have devastating consequences. However, the majority of overdoses involve analgesics, antidepressants, hypnotics, anxiolytics, and antipsychotic medications whereas colchicine overdose appears to be an uncommon occurrence.²⁻⁴ Colchicine is an effective and useful treatment for both acute attacks of gout and prophylaxis against acute attacks when commencing urate-lowering therapies such as allopurinol, particularly in the many patients who are intolerant of or have contraindications to non-steroidal anti-inflammatory drugs.^{5,6} Although we concur with Lloyd that assessment of mood and risk of overdose should be considered when prescribing any medication, we urge prescribers not to abandon an effective treatment for this excruciatingly painful and frequently poorly managed condition.

Edward Roddy,

Clinical Senior Lecturer in Rheumatology and Consultant Rheumatologist, Research Institute of Primary Care and Health Sciences, Keele University.
E-mail: e.rodgy@keele.ac.uk

Christian D Mallen,

NIHR Research Professor in General Practice, Research Institute of Primary Care and Health Sciences, Keele University.

REFERENCES

1. Lloyd G. Colchicine in overdose. *Br J Gen Pract* 2016; **66(653)**: 605. DOI: <https://doi.org/10.3399/bjgp16X688333>.
2. Prescott K, Stratton R, Freyer A, *et al*. Detailed analyses of self-poisoning episodes presenting to a large regional teaching hospital in the UK. *Br J Clin Pharmacol* 2009; **68(2)**: 260-268.
3. Doak MW, Nixon AC, Lupton DJ, Waring WS. Self-poisoning in older adults: patterns of drug ingestion and clinical outcomes. *Age Ageing* 2009;

38(4): 407-411.

4. Armstrong TM, Davies MS, Kitching G, Waring WS. Comparative drug dose and drug combinations in patients that present to hospital due to self-poisoning. *Basic Clin Pharmacol Toxicol* 2012; **111(5)**: 356-360.
5. Jordan KM, Cameron JS, Snaith M, *et al*. British Society for Rheumatology and British Health Professionals in Rheumatology guideline for the management of gout. *Rheumatology (Oxford)* 2007; **46(8)**: 1372-1374.
6. Richette P, Doherty M, Pascual E, *et al*. 2016 updated EULAR evidence-based recommendations for the management of gout. *Ann Rheum Dis* 2017; **76(1)**: 29-42. DOI: [10.1136/annrheumdis-2016-209707](https://doi.org/10.1136/annrheumdis-2016-209707). Epub 2016 Jul 25.

DOI: <https://doi.org/10.3399/bjgp17X688993>

Barriers to lifestyle changes in people with diabetes

The study on the impediments to good diabetes control by Elliott *et al*¹ corresponds with our findings from an interview study of patients with diabetes from a deprived background in the northeast of England. We undertook semi-structured interviews with eight patients with established type 2 diabetes from one general practice and explored their perceived barriers to making lifestyle changes. The incidence of diabetes in the northeast of England is relatively high and is known to be inversely related to a low socioeconomic status,²⁻⁴ but there is little research exploring the reasons for the link with deprivation. Our results highlighted four common themes: dietary education, motivation to change, family support, and comorbidities.

Patient education was the most commonly reported barrier, related to poor dietary education and misperceptions gained during childhood on healthy foods. Participants struggled to find the motivation to change their lifestyles. This fluctuated in some participants, who reported returning to bad habits during prolonged periods of no medical contact. Interestingly, the level of perceived support a patient felt they had seemed to correlate with their overall motivation. Participants who were in a relationship commented on how their partner acted as a source of continual encouragement to help them continue with the lifestyle regimen they had been given. However, following a healthy lifestyle was challenging for family members of our

patients, especially those who were part of a big family with children. Surprisingly, financial barriers were felt to be less of an issue than might have been expected: most participants suggested that the cost of leading a healthier lifestyle was balanced by the cost of unhealthy habits such as 'takeaways'. Finally, the participants' overall health and the consequential impact of their comorbidities were highlighted as an important barrier to following a healthy lifestyle.

Harley Fuller,

Medical Student, Newcastle University.

Hugh Alberti,

GP, Subdean for Primary and Community Care, Newcastle University.

REFERENCES

1. Elliott AJ, Harris F, Laird SG. Patients' beliefs on the impediments to good diabetes control: a mixed methods study of patients in general practice. *Br J Gen Pract* 2016; **66(653)**: e919-e919. DOI: <https://doi.org/10.3399/bjgp16X687589>.
2. Connolly V, Unwin N, Sherriff P, *et al*. Diabetes prevalence and socioeconomic status: a population based study showing increased prevalence of type 2 diabetes mellitus in deprived areas. *J Epidemiol Community Health* 2000; **54(3)**: 173-177.
3. Robbins JM, Vaccarino V, Zhang H, Kasl SV. Socioeconomic status and diagnosed diabetes incidence. *Diabetes Res Clin Pract* 2005; **68(3)**: 230-236.
4. Ricci-Cabello I, Ruiz-Pérez I, Olry de Labry-Lima A, Márquez-Calderón S. Do social inequalities exist in terms of the prevention, diagnosis, treatment, control and monitoring of diabetes? A systematic review. *Health Soc Care Community* 2010; **18(6)**: 572-587.

DOI: <https://doi.org/10.3399/bjgp17X689005>

The recipe for general practice

The production process for fresh GPs from our specialty training is like the process for making sausages: a pleasing end product, but we distract ourselves from thinking about the content too much.^{1,2}

The difficulties of the right mix of hospital training placements is that, for general practice, every hospital-based speciality is both relevant, and yet irrelevant. But why hospital-based training at all? No GP programme director has yet been troubled