Anna Cheshire, Damien Ridge, John Hughes, David Peters, Maria Panagioti, Chantal Simon and George Lewith

Influences on GP coping and resilience:

a qualitative study in primary care

Abstract

Background

'Neoliberal' work policies, austerity, NHS restructuring, and increased GP consultation rates provide the backdrop against increasing reports of GP burnout and an impending shortage of GPs.

Aim

To explore GPs' experiences of workplace challenges and stresses, and their coping strategies, particularly focusing on understanding the impact of recent NHS workplace change.

Design and setting

Study design was qualitative, with data collected from two focus groups and seven one-to-one telephone interviews.

Method

Focus groups and one-to-one telephone interviews explored the experiences of GPs currently practising in England, recruited through convenience sampling. Data were collected using a semi-structured interview approach and analysed using thematic analysis.

Results

There were 22 GP participants recruited: focus groups ($n = 15$) and interviews ($n = 7$). Interviewees understood GPs to be under intense and historically unprecedented pressures, which were tied to the contexts in which they work, with important moral implications for 'good' doctoring. Many reported that being a full-time GP was too stressful: work-related stress led to mood changes, sleep disruption, increases in anxiety, and tensions with loved ones. Some had subsequently sought ways to downsize their clinical workload. Workplace change resulted in little time for the things that helped GP resilience: a good work-life balance and better contact with colleagues. Although some GPs were coping better than others, GPs acknowledged that there was only so much an individual GP could do to manage their stress, given the external work issues they faced.

Conclusion

GPs experience their emotional lives and stresses as being meaningfully shaped by NHS factors. To support GPs to provide effective care, resilience building should move beyond the individual to include systemic work issues.

Keywords

coping skills; general practitioners; political factors; primary health care; professional burnout; psychological resilience.

INTRODUCTION

Contemporary general practice

Post-1970s, Western governments have pursued 'neoliberal' policies^{1,2} prioritising employee productivity and flexibility, alongside increasing job insecurity and unpaid overtime.^{3,4} Subsequently, there was commodification of NHS health care, increasingly conceptualised as a competitive marketplace.⁵ Cost, value, and privatisation were prioritised,⁶ creating variation in access to health care.⁷ NHS policies have reduced GPs' autonomy too.⁸ The 2004 General Medical Services (GMS) contract, although initially helpful,⁹ increased the authority of primary care trusts¹⁰ and created a market for resources.⁹ The cycle of performance management, monitoring, and competition for scarcer resources has resulted in NHS savings,¹¹ but created a focus on cost-effective health care; competition for funding against quality standards; increased primary care workload for conditions previously managed in secondary care;¹² and growing responsibility for delivering quality with fewer resources.^{13,14}

As well as neoliberalism, austerity, and NHS restructuring, GPs face cultural

changes with increased patient demand and expertise,^{12,15} and new technologies to administer assessments and clinical care.⁸ Although these trends may add to quality and patient-centredness,¹⁶ they also present challenges for the profession.⁸

Links to GP wellbeing

Research suggests that uncertainty at work contributes to distress and dysfunction among healthy adults,¹⁷ although links between rapid change and worker wellbeing are unclear.^{18,19} GPs in the UK face cognitively and emotionally challenging environments with high workloads and long hours.²⁰ Although many cope successfully,²¹ reports of distress and burnout, and related negative job performance, are increasing.^{18,19,22} Burnout is a descriptive measure (rather than a clinical diagnosis) of feelings of emotional exhaustion, depersonalisation, and a diminished sense of personal accomplishment primarily driven by workplace stressors.^{18,22} One in three GPs from the UK and Europe are experiencing burnout;²² GPs in the UK report lower levels of work satisfaction than other Western countries,²³ where one in six GPs report unmanageable distress.²⁴ Doctor

A Cheshire, PhD, research fellow; **D Ridge**, PhD, professor and head of psychology; University of Westminster, Department of Psychology, London.

J Hughes, PhD, senior research coordinator, Royal London Hospital for Integrated Medicine UCLH NHS Trust, London. **D Peters**, MD, FRCGP, clinical director, University of Westminster Polyclinic and Centre for Resilience, University of Westminster, London. **M Panagioti**, PhD, research fellow, Institute of Population Health, Centre for Primary Care, Manchester. **C Simon**, PhD, MA, MRCGP, GP partner, the Banks and Bearwood Medical Centres, Bournemouth; and medical director for professional development, Royal College of General Practitioners, London.

G Lewith, DM, FRCGP, MRCGP, professor of health research, Primary Care, University of Southampton, Aldermoor Health Centre,

Southampton; and visiting professor, University of Westminster Centre for Resilience, London.

Address for correspondence

Damien Ridge, University of Westminster, Department of Psychology, 115 New Cavendish Street, London W1W 6UW, UK.

E-mail: d.ridge@westminster.ac.uk

Submitted: 19 September 2016; **Editor's response:** 14 October 2016; **final acceptance:** 14 November 2016.

©British Journal of General Practice

This is the full-length article (published online 9 May 2017) of an abridged version published in print. Cite this article as: **Br J Gen Pract 2017; DOI: <https://doi.org/10.3399/bjgp17X690893>**

How this fits in

Primary care is currently facing unprecedented challenges including increasing GP burnout and staff shortages. GPs report being under intense and historically unparalleled pressures that are tied to the contexts in which they currently work. In the intense, micromanaged, competitive NHS 'marketplace' the GP study participants were conscious of the potential damage to relationships (to self and others) the current system engenders. At the individual level, resilience training may be of benefit to GPs, yet an exclusive focus on improving individual coping risks sidestepping the systemic challenges shaping primary care.

distress increases professional inefficiency, depression, and alcohol and substance misuse,²⁵⁻²⁷ and is linked to medication errors, lower patient satisfaction, and non-adherence to treatment.²⁸⁻³⁰ Stress also reduces productivity,³¹ increasing absenteeism, presenteeism, job turnover, clinical errors, and early retirement.^{32,33} GP vacancy rates in the UK are at their highest ever levels,³⁴ with more GPs planning to leave.³⁵ The desire to leave general practice is linked to high stress levels,²³ and government plans for routine 7-day GP working will create further challenges for practices.³⁶

In the context of these complex changes in the GP workplace, and their links to wider political changes, this study used a qualitative interpretative approach in order to address the question: what are GPs' experiences of contemporary workplace challenges and stresses, and their preferred coping strategies?

METHOD

Design

Focus groups allowed GP discussions about their experiences of the current 'austerity NHS'. GPs are busy,¹² therefore flexible telephone interviews (covering the same topics) were offered to those GPs unable to attend a focus group. The interview topic guide was further informed by themes emerging from the group discussions.³⁷

Participants and recruitment

Recruitment packs including participant information sheets were made available to GPs at a resilience talk delivered at the Royal College of General Practitioners (RCGP) 2015 Annual Conference in Glasgow.

Additionally, a study flyer was placed on the RCGP website and sent to local RCGP faculties and medical committees. The authors used their extensive network of primary care contacts, targeting GP gatekeepers, asking them to distribute the flyer to their contacts, and by means of snowballing, with those recruited asked to contact colleagues about participating in the study.

The inclusion criterion was: currently practising as a GP in England. GPs who expressed an interest were emailed a participant information sheet and consent form, and invited to a focus group in London or Bournemouth, or a telephone interview. Participants received no financial reimbursement for participation.

Data collection

A semi-structured approach to data collection was adopted. Topics covered current sources of GP stress, coping strategies, and barriers or facilitators to successful coping. Focus groups lasted 37 and 77 minutes, interviews between 35 and 65 minutes, and all were conducted by an experienced qualitative researcher. Discussions in focus groups developed easily and, once the facilitator raised a topic, minimal facilitation was required. Focus groups allowed debate and drawing out of issues, whereas phone interviews explored underlying issues and in-depth individual experiences.³¹ The point of data saturation (when no new themes of interest were emerging)³⁸ was debated between the first authors, and determined to be 22 participants. Interviews and groups were recorded and transcribed verbatim; transcripts were checked for accuracy and anonymised.

Analysis

A constructivist epistemological approach was adopted. The study was approached from the position that '*data do not provide a window on reality, rather, the "discovered" reality arises from the interactive process and its temporal, cultural, and structural contexts*'.³⁹ Data were analysed inductively,⁴⁰ using thematic analysis.⁴¹ Two researchers immersed themselves in the data, repeatedly reading the transcripts to understand participants' experiences. Key issues, concepts, and themes arising from the data were identified and debated, creating a draft coding framework that was discussed with the research team, to construct the final conceptual framework. Transcripts were coded and explored in NVivo 11 and findings were written up into

Table 1. Participant demographics

	N	%
Age, years		
20–29	1	4.5
30–39	10	45.5
40–49	3	13.6
50–59	6	27.3
Missing	2	9.1
Sex		
Female	18	81.8
Male	4	18.2
Type of GP		
Salaried	10	45.5
Partner	6	27.3
Trainee	4	18.2
Locum	1	4.5
Rapid response	1	4.5
Working hours		
Full time	14	63.6
Part time	8	36.4
Type of practice		
Urban	16	72.7
Rural	5	22.7
Mixed	1	4.5

a draft, which was then debated and refined by all authors.

RESULTS

Twenty-two GPs participated in the study (January to March 2016): two focus groups (Bournemouth, $n = 8$; London, $n = 7$) and seven telephone interviews. A wide demographic was recruited in terms of age, sex, type of GP, practice type, and working hours (Table 1).

The findings explore core workplace challenges discussed by GPs, their experiences of workplace stress, and how they are coping. These are presented around the following broad themes: work intensification and morality issues; intensification and patient complexities; GP coping, work–life balance, and downsizing (see Box 1 for summary).

Work intensification and morality: ‘It’s becoming very Big Brother’

Interviewees felt GPs to be under intense and historically unprecedented pressures, suggesting these issues were tied to work contexts, with important moral implications. NHS factors were considered particularly important in understanding GP stress. Participants stated that linking the Quality and Outcomes Framework (QOF) directly to GP funding was detrimental to GP wellbeing and patient-centred care. In particular, some participants argued that QOF made unreasonable demands on a 10-minute consultation, and had more to do with allocating funding than good use of evidence-based practice:

‘To be a good GP you need to think about these things, but in order to obtain your funding you need to run through your little hoops. So it might be that your computer’s saying, needs this check and that check and this check. And you think, oh no, I haven’t got time to do all that.’ (FG1F) (focus group [FG] or interview [I] participant, sex [M/F], age in years, and working hours full time [FT]/part time [PT] — if available)

Participants agreed their human connection with patients was important, and worried that they were less able to connect because of conflicting time demands. The imperative to collect and record patient indicators during consultations, and the need to work within a new NHS marketplace for resources, reportedly risked undermining the ‘art’ of medicine and ‘good’ care by impinging on GPs’ ability to authentically engage with patients. Primary care was said to be functioning in more detached ways, *‘like secondary care’*, while being micromanaged and professionally deskilled. Care Quality Commission (CQC) inspections also were seen as increasing the intensity — and external scrutiny — of GP work:

‘It’s becoming very Big Brother, the whole of the NHS and especially general practice about what we have to do in order to earn money and look after people. It’s becoming more, obviously guidelines and protocols are really useful and we need those, especially as things become more complex. But that’s squeezing out just the relational aspect of general practice, which is a lot of the time what people need and where help and healing really happens I think.’ (I24F, 36, PT)

‘You want me to put them on a tablet just to get the money in? This is not what I signed myself up for ... So the other obstacle is trying to get these other GPs on board who have moved onto the “other side”, which is, we must get QOF, we must get QOF, we must get QOF.’ (I26M, 45, PT)

A perceived negative portrayal of GPs by the UK media and politicians particularly concerned participants: the public prestige and/or esteem of the profession seemed to be under threat, affecting morale. Others felt the coverage negatively influenced the patients’ trust in GPs, which could in turn affect consultations and their sense of being a ‘good’ doctor:

‘Negative portrayal of the profession in the press I think, is a strong demotivator. It

Box 1. Participant themes and subthemes

Theme	Subthemes within narratives
Work intensification and morality: <i>‘It’s becoming very Big Brother’</i>	<ul style="list-style-type: none">• NHS factors affecting GP stress• Quality and Outcomes Framework• Primary care as an NHS ‘marketplace’• Care Quality Commission inspections• Maintaining the ‘human’ connection with patients• Portrayal of GPs by the UK media and politicians• Patient empowerment, complaints culture, and defensive practice
Intensification and patient complexities	<ul style="list-style-type: none">• Competing demands• Pressure of time• Complexity of workload• Resources• Personal patient lists• The 10-minute consultation• The proposed 7-day working week
GP coping, work–life balance, and downsizing	<ul style="list-style-type: none">• GP coping• Impacts of GP stress• Work–life balance• Coping strategies• Limitations of resilience

actually impacts quite significantly on the doctor–patient relationship as well ... When they walk into the consulting room, that wariness is very, very difficult to overcome, to gain a rapport. (I25M, 38, PT)

GPs endorsed patient empowerment and favoured patients making formal complaints where medical care was genuinely compromised. However, there was a perception that trivial complaints were being normalised within a ‘complaints culture’. Consequently, doctors practised more defensively by increasing their paper trails or making referrals they felt were probably unwarranted:

‘I feel the need increasingly to be defensive about the way that one practises because of the increased levels of complaints and increased empowerment really, the complaints that one receives, this takes an immense amount of time and emotional energy to respond to.’ (I25M, 38, PT)

Intensification and patient complexities

GPs tried in various ways to convey the competing demands that they had to balance in their work, emphasising the unrelenting ‘pressure of time’:

‘The thing that is most stressful I think is the relentless nature of it a lot of the time. It seems a lot of the time like a constant race against the clock from half eight in the morning or before to seven at night.’ (FG1M)

GPs saw consultations with patients as their ‘bread and butter’. But day-to-day administrative duties, phone calls to patients, and ‘surprise work’ (for example, working with social services or paramedics on behalf of specific patients) inevitably increased their workload. GPs perceived that the complexity of their work was increasing without additional resource. An ageing population needing more consultations was presenting with more complex multimorbidities. Additionally, GPs were taking on the management of chronic conditions previously referred to secondary care. They were concerned about being out of their depth or ‘set up to fail’:

‘Patients are getting much more complex and they’re getting much more demanding.’ (FG1F)

‘It’s what secondary care is putting back into primary care, it’s very complex because they’re reducing mental health services and pushing patients back on to us, we have to

deal with. We are not properly trained to deal with them, they are taking a lot more time. We try to refer them, they get bounced back at us.’ (FG1F)

Changes to practice management, such as GPs not having their own personal lists, were thought to contribute to a lack of continuity of care, with less ability to develop long-term patient relationships. This tended to reduce the efficiency and effectiveness of consultations, because GPs had to ‘go back to square one’ with unfamiliar patients:

‘I don’t have a list ... I may never have met them [patient] and not know anything about the issue, but in order to safeguard them I do have to go back to square one to find out where we are, before we then decide how we move forward. So it’s not like a follow-up, it’s like a new consultation which takes more time.’ (I14F, 57, FT)

Ten minutes for each patient was unanimously perceived as inadequate for treating increasingly empowered patients with complex issues; clinics routinely ran late, and GPs often felt unable to take adequate working breaks. They universally worked longer hours than contracted, and were uneasy about the impact of the proposed government 7-day working week:

‘There are mornings where I think, oh dear, is this going to be a two-wee day or a one-wee day?’ (FG1F)

‘We were all thinking, we can’t cope with 5 days at the moment, how on earth are they going to get us doing 7 days!’ (FG1M)

GP coping, work–life balance, and downsizing

Some GPs are coping better than others, although reports of coping often came with caveats (for example, working longer hours to cope):

‘Sometimes you just stay late or you come in another day and you do your extra work. I find actually I’d rather do that than cope with that feeling of being overwhelmed really.’ (FG1F)

‘Well I think some people innately can always look at the cup half full can’t they, and I probably have that personality or I wouldn’t have survived this long.’ (I3F, 59, FT)

However, most GPs were adamant that being a full-time GP was now ‘too stressful’. For some, their current role was perceived

as undermining their ability to function effectively, or even safely. Cognitively stressed GPs felt unable to handle the levels of incoming information, and were worried they might make errors. Work-related stress led to changes in mood, disruptions to sleep patterns, and increases in anxiety:

'I felt unsafe in my practice because my head was too full and there was too much going on and I didn't feel I was able to think clearly, rationally.' (I24F, 36, PT)

'You're constantly worried and the more stressed you are the more worried that, "oh my God, I'm definitely missing something now".' (FG1F)

Many participants said that being a full-time GP was incompatible with an adequate work-life balance. Female GPs with children experienced this issue most acutely. Childcare forced some GPs to 'down tools' earlier, but this then meant working overtime to catch up. Even if partners and families were supportive, many GPs still lamented the stress they experienced, with limited time for loved ones:

'Then there's the stress, we're talking about stressors, the stress of not knowing that you're not going to get home for bed time with the kids or to have a bit of a row with the other half because you've been late leaving yet again.' (I25M, 38, PT)

GPs also highlighted the lack of time to pursue hobbies or leisure activities. Yet a good work-life balance was widely considered to increase GPs' resilience and better equip them to deal with the stresses associated with their role:

'I think an understanding that you have a right to have a life outside your job ... And that actually the richer that is probably the more resilient a GP you could be I think.' (FG1F)

All participants spoke about the strategies they employed to mitigate work stress. These included meditation and or mindfulness, stress management techniques, taking regular exercise, and eating well. Participants also adapted practical aspects of their day-to-day working routine in an effort to ease their workload and/or make their work more efficient:

'I think one of my ways of coping is to do blocks of work in chunks, so different modalities of work. So say results, so try

and not switch over to documents say, to go through my block of blood results before I move on to the next thing like phone calls.' (FG1F)

'And mindfulness isn't just meditating, it's lots of other things as well. And that just helps me, it helps keep me in check.' (I26M, 45, PT)

'I mean just lifestyle things, making sure you've slept enough and you haven't drunk too much alcohol the night before, and that you eat regularly and you exercise, all of those things.' (I24F, 36, PT)

Other participants reported focusing on what they enjoyed about their job, which helped with their stress, and included helping patients and having a supportive and friendly practice. Space for meeting and debriefing with the team was perceived as having a positive impact on stress levels, however increasing work demands meant GPs had less time to connect with colleagues:

'There's been nothing official, but I have felt supported, appreciated, and been able to talk about stuff when I needed to with my colleagues, but that is changing or has changed a lot. Because one of the biggest differences from when I started in the practice where I'm in now, is that I now arrive at work half an hour early to get all the login and look at my path links and things like that, so I go in and I shut my door. And then I can possibly be in there for about 11 hours without much opportunity to talk to anybody.' (I14F, 57, FT)

'We all love our job, we work for the patients, that's why we do this job' (FG1F)

Participants emphasised that no matter how good they were at coping with external factors and work stressors, there was only so much that individual GPs could do to cope. Many factors, such as 10-minute consultations and QOF, were considered unlikely to change imminently. A number of participants had implemented far-reaching changes, having come to a gradual awareness (or a defining moment) of the limits of their ability to cope working full-time. Participants talked about colleagues who had left the profession. A surprisingly high number of participants had reduced their working hours or changed their role (for example, to salaried or locum) to enable them to downsize, better cope, and regain some work-life balance:

'Since qualifying I've really been doing a regular six-session week, so that means 3 days, I've been doing some locum shifts as and when. But it just means that if I know that those 3 days I'm always in work 12 hours at least, and then I have a long drive home at the moment, so actually that's 36 hours a week so that's full-time work. That's really helped me. At the moment I'm now going to a regular yoga class, I've got a bit fitter, I can see my friends ... Which is not great for the GP workforce on the whole because if it was more manageable I could do 5 days, no problem, but because it's not manageable.' (FG1F)

'I'd say even among my cohort of ST3s, I would say probably half of us are thinking of just working part time.' (FG2F)

'What you've got to be careful to do is not ignore the fact that actually, maybe, for most of us, we are not coping with the stressors because there's too much stress, not because we're not resilient enough. And therefore if you don't solve the root cause you get nowhere.' (FG2M)

DISCUSSION

Summary

GP work is shaped by the policy agendas affecting other public services. The profound impact of increasing workloads, and increasing demands on GPs, needs highlighting. The development of the NHS as a marketplace, with increased regulation but under-resourcing, is perceived as detracting from quality patient-centred care. The moral implications of work intensification include disconnection from patients and fears about bad doctoring. GPs increasingly feel the need to practise more defensively, while at the same time they may be too busy to connect with their peers to adequately debrief.

Some GPs had found ways to cope with conflicts between work and home life, but others simply looked to reduce their workload. Concurrently an ageing patient population, increasing patient contact, relocation of secondary care services into the community, and rising public expectations have intensified GP workload. Participants acknowledged the need to build personal resilience, but also recognised that ultimately organisational change is needed to improve their wellbeing and job satisfaction.

Strengths and limitations

The study sample was from a range of demographics, practices, and roles, but

with a larger number of females, more salaried GPs than partners, and more participants from urban practices,^{42,43} yet the proportions of full- and part-time GPs were consistent with national figures.⁴² The sample size ($n = 22$) is adequate for this type of qualitative study, and the data reached saturation.⁴⁴

The sampling methods may have attracted GPs who were interested in resilience and had time to participate. These GPs may be coping better than others. Nevertheless, a number of the participants reported that they were not — or had not previously been — coping well. GPs who had left the profession were not interviewed, nor were there questions focusing specifically on the positive aspects of GPs' careers.^{45,46} However, interviews and focus groups provided a complementary combination of data collection methods.

Comparison with existing literature

Others have described similar pressures on GPs,^{12,14} including dissatisfaction with appointment length,²³ increasing patient expectation and demand,^{12,45} negative media portrayals, and a reduced ability to practise patient-centred care.⁴⁵ Although the proportion of NHS funding for primary care has declined, consultation rates and workload have increased dramatically,^{12,14} thus creating a 'feeling of crisis' in primary care.¹² Research has identified similar trends across Europe.⁴⁷

A contradiction between government rhetoric that puts patients '*at the centre of everything the NHS does*'⁴⁸ and current policy was identified. The NHS as a 'marketplace' places a limit on quality of care.⁴⁹ Previous research has found that neoliberal ideas encourage a '*specific way of being, acting, and understanding the world, which is highly individualistic*';⁵⁰ the authors found that NHS marketisation promotes a kind of GP individualism — or atomisation — where GPs feel increasingly removed from meaningful interactions with patients and colleagues, even loved ones. Although some doctors cope better in the new NHS than others,⁵¹ there is a growing recognition that primary care is at breaking point.^{12,14} Early indicators of change may be government proposals to cut bureaucracy⁵² and abandon the QOF,⁵³ and the British Medical Association's call to increase GP consultation time.⁵⁴

Implications for research and practice

Resilient doctors are better at caring for others, and are less likely to commit errors, get sick, or leave practice.^{55,56} Systemic

changes to the work environment^{55,57,58} and educational and organisational interventions that could increase physician resilience are key to more effective primary care. Resilience training should be a preventive strategy rather than a response to overwhelming systemic problems.⁵¹ Mindful self-compassion, optimism, adaptability and prioritisation, teamwork, supportive relationships, and job-related satisfaction all enhance resilience.^{21,59–60} However, resilience building requires a synergy between personal characteristics, management support, teamwork, and

workplace/social buffers.⁶¹ The study findings suggest that NHS organisational factors and wider government policy impact directly on GPs' emotional lives in predictable ways. Further research should address how organisational change can best promote both individual and systemic resilience.⁵⁶ It is important to be wary though of how neoliberal ideas could turn resilience building into a project of the self, with a focus on individual choice and the pursuit of individual wellbeing, while neglecting the systemic realities facing GPs in the NHS.^{50,62}

Funding

The study received funding from the Westminster Centre for Resilience and the Royal College of General Practitioners.

Ethical approval

Ethical approval for the study was obtained through the University of Westminster Ethics Committee. NHS ethical approval was not required for this study.

Provenance

Freely submitted; externally peer reviewed.

Competing interests

The authors have declared no competing interests.

Acknowledgements

The authors thank the Royal College of General Practitioners and Dragana Milosevic for supporting the study and providing funding; all those who helped with recruitment, particularly Chris Manning, and the practices who hosted the focus groups and the staff who helped organise them; and all the participants who gave up their time to take part in the study. Finally, a special thanks to George Lewith, who passed away during the writing of the article, for all his enthusiasm and work on this project.

Discuss this article

Contribute and read comments about this article: bjgp.org/letters

REFERENCES

- Rose N. *Governing the soul: the shaping of the private self*. London: Taylor & Francis/Routledge, 1990.
- Illouz E. *Cold intimacies: the making of emotional capitalism*. Cambridge: Polity, 2007.
- Bauman Z. *Liquid modernity*. Cambridge: Polity Press, 2000.
- Burchell B, Ladipo D, Wilkinson F. *Job insecurity and work intensification*. London: Routledge, 2005.
- Greenhalgh T. Future-proofing relationship-based care: a priority for general practice. *Br J Gen Pract* 2014; DOI: <https://doi.org/10.3399/bjgp14X682357>.
- Pollock AM. *NHS plc: the privatisation of our health care*. London: Verso, 2004.
- Bambra C, Fox D, Scott-Samuel A. Towards a politics of health. *Health Promot Int* 2005; **20**(2): 187–193.
- Harrison S. Co-optation, commodification and the medical model: governing UK medicine since 1991. *Public Adm* 2009; **87**(2): 184–197.
- Russell J, Greenhalgh T, Burnett A, *et al*. 'No decisions about us without us'? Individual healthcare rationing in a fiscal ice age. *BMJ* 2011; **342**: d3279.
- Grant S, Huby G, Watkins F, *et al*. The impact of pay-for-performance on professional boundaries in UK general practice: an ethnographic study. *Social Health Illn* 2009; **31**(2): 229–245.
- Grant S, Ring A, Gabbay M, *et al*. Soft governance, re-stratification and the 2004 general medical services contract: the case of UK primary care organisations and general practice teams. *Social Health Illn* 2015; **37**(1): 30–51.
- Baird B, Charles A, Honeyman M, *et al*. *Understanding pressures in general practice*. London: King's Fund, 2016.
- Lionis C, Petelos E. The impact of the financial crisis on the quality of care in primary care: an issue that requires prompt attention. *Qual Prim Care* 2013; **21**(5): 269–273.
- Primary Care Workforce Commission. *The future of primary care: creating teams for tomorrow*. Leeds: Health Education England, 2015.
- Hobbs FD, Bankhead C, Mukhtar T, *et al*. Clinical workload in UK primary care: a retrospective analysis of 100 million consultations in England, 2007–14. *Lancet* 2016; **387**(10035): 2323–2330.
- Lupton D. Doctors on the medical profession. *Social Health Illn* 1997; **19**(4): 480–497.
- Melchior M, Caspi A, Milne B, *et al*. Work stress precipitates depression and anxiety in young, working women and men. *Psychol Med* 2007; **37**(8): 1119–1129.
- Maslach C, Schaufeli WB, Leiter MP. Job burnout. *Ann Rev Psychology* 2001; **52**: 397–422. Epub 2001/01/10.
- Mind. Mind finds 'Worrying' levels of stress among primary care staff. 2016. <http://www.mind.org.uk/news-campaigns/news/mind-finds-worrying-levels-of-stress-among-primary-care-staff/#.V8WFBqKbEXI> (accessed 20 Apr 2017).
- Bell BG, Reeves D, Marsden K, Avery A. Safety climate in English general practices: workload pressures may compromise safety. *J Eval Clin Pract* 2016; DOI: 10.1111/jep.12437.
- Zwack J, Schweitzer J. If every fifth physician is affected by burnout, what about the other four? Resilience strategies of experienced physicians. *Acad Med* 2013; **88**(3): 382–389.
- Soler JK, Yaman H, Esteve M, *et al*. Burnout in European family doctors: the EGPRN study. *Fam Pract* 2008; **25**(4): 245–265.
- Martin S, Davies E, Gershlick B. *Under pressure: what the Commonwealth Fund's 2015 international survey of general practitioners means for the UK*. London: Health Foundation, 2016.
- ICM Unlimited, British Medical Association. *British Medical Association national survey of GPs: the future of general practice 2015*. London: BMA, 2015.
- Dyrbye LN, West CP, Satele D, *et al*. Burnout among U.S. medical students, residents, and early career physicians relative to the general U.S. population. *Acad Med* 2014; **89**(3): 443–451.
- Embrico N, Hraiech S, Azoulay E, *et al*. Symptoms of depression in ICU physicians. *Ann Intensive Care* 2012; **2**(1): 34.
- Pedersen AF, Sørensen JK, Bruun NH, *et al*. Risky alcohol use in Danish physicians: associated with alexithymia and burnout? *Alcohol Drug Depend* 2016; **160**: 119–126.
- Halbesleben JRB. Patient reciprocity and physician burnout: what do patients bring to the patient-physician relationship? *Health Serv Manage Res* 2006; **19**(4): 215–222.
- Walocha E, Tomaszewski KA, Wilczek-Ruzyczka E, *et al*. Empathy and burnout among physicians of different specialities. *Folia Med Cracov* 2013; **53**(2): 35–42.
- Anagnostopoulos F, Liolios E, Persefonis G, *et al*. Physician burnout and patient satisfaction with consultation in primary health care settings: evidence of relationships from a one-with-many design. *J Clin Psychol Med Settings* 2012; **19**(4): 401–410.
- Stokes D, Bergin R. Methodology or methodolatry: an evaluation of focus groups and depth interviews. *Qual Mark Res* 2006; **9**(2): 26–36.
- Linzer M, Visser MRM, Oort FJ, *et al*. Predicting and preventing physician burnout: results from the United States and the Netherlands. *Am J Med* 2001; **111**(2): 170–175.
- Dewa CS, Loong D, Bonato S, *et al*. How does burnout affect physician productivity? A systematic literature review. *BMC Health Serv Res* 2014; **14**: 325.
- Kaffash J. GP vacancy rates at highest recorded with one in eight positions unfilled. *Pulse* 2016. <http://www.pulsetoday.co.uk/your-practice/practice-topics/employment/gp-vacancy-rates-at-highest-recorded-with-one-in-eight-positions-unfilled/20031836.article> (accessed 20 Apr 2017).
- Hann M, McDonald J, Checkland K, *et al*. *Seventh national GP worklife survey*. Manchester: University of Manchester, 2013.
- Twaddell I. BMA votes to resist plans for a non-urgent seven-day NHS service. *Pulse* 2016. <http://www.pulsetoday.co.uk/hot-topics/seven-day-gp-access/bma-votes-to-resist-plans-for-a-non-urgent-seven-day-nhs-service/20032117.fullarticle> (accessed 20 Apr 2017).
- Minichiello V, Aroni R, Hays T. *In-depth interviewing: principles, techniques, analysis*. Melbourne: Longman Cheshire, 1995.
- Morse JM. The significance of saturation. *Qual Health Res* 1995; **5**(2): 147–149.
- Charmaz K. Grounded theory: objectivist and constructivist methods. In: Denzin N, Lincoln Y, eds. *Handbook of qualitative research*. 2nd edn. Thousand Oaks, CA: Sage, 2000.
- Bowling A. *Research methods in health: investigating health and health services*. Buckingham: Open University Press, 1997.
- Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006; **3**(2): 77–101.
- NHS Digital. *General and Personal Medical Services, England September 2015–September 2016*. Surrey: Health and Social Care Information Centre, 2016.
- Commonwealth Fund. *2015 Commonwealth Fund international survey of primary care doctors in 10 nations*. New York, NY: Commonwealth Fund, 2015.
- O'Reilly M, Parker N. 'Unsatisfactory saturation': a critical exploration of the notion of saturated sample sizes in qualitative research. *Qual Health Res* 2012; **13**(2): 190–197.
- Doran N, Fox F, Rodham K, *et al*. Lost to the NHS: a mixed methods study of why GPs leave practice early in England. *Br J Gen Pract* 2016; DOI: <http://doi.org/10.3399/bjgp16X683425>.
- Spooner S. Unfashionable tales: narratives about what is [still] great in NHS general practice. *Br J Gen Pract* 2016; DOI: <http://doi.org/10.3399/bjgp16X683401>.
- Lippert ML, Reventlow S, Kousgaard MB. The uses and implications of standards in general practice consultations. *Health* 2017; **21**(1): 3–20.
- Department of Health. Complaints figures published by the Information Centre. DH, 2010. http://webarchive.nationalarchives.gov.uk/20130107105354/http://www.dh.gov.uk/en/MediaCentre/Statements/DH_118993 (accessed 20 Apr 2017).
- Gilbert BJ, Maruthappu M, Leaver L, *et al*. Morality in the NHS marketplace. *Lancet* 2013; **382**(9910): 2065–2066.
- Cabanis E. Rekindling individualism, consuming emotions: constructing 'psytizens' in the age of happiness. *Cult Psychol* 2016; **22**(3): 467–480.
- Lee FJ, Stewart M, Brown JB. Stress, burnout, and strategies for reducing them: what's the situation among Canadian family physicians? *Can Fam Physician* 2008; **54**(2): 234–235.
- Matthews-King A. GP payment system to be reformed to end 'unacceptable' burden on practice time. *Pulse* 2016. <http://www.pulsetoday.co.uk/hot-topics/general-practice-forward-view/gp-payment-system-to-be-reformed-to-end-unacceptable-burden-on-practice-time/20031654.article> (accessed 20 Apr 2017).
- Matthews-King A. QOF has reached the end of its useful life, says NHS chief. *Pulse* 2016. <http://www.pulsetoday.co.uk/your-practice/qof/qof-has-reached-the-end-of-its-useful-life-says-nhs-chief/20033054.article> (accessed 20 Apr 2017).
- Siddique H. GP appointments should be five minutes longer, says BMA. *Guardian* 2016; **28 Aug**: <https://www.theguardian.com/society/2016/aug/28/>

- doctor-appointments-15-minutes-bma-overweight-population.
55. Epstein RM. Realizing Engel's biopsychosocial vision: resilience, compassion, and quality of care. *Int J Psychiatry Med* 2014; **47(4)**: 275–287.
 56. Lown M, Lewith G, Simon C, Peters D. Resilience: what is it, why do we need it, and can it help us? *Br J Gen Pract* 2015; DOI: <http://doi.org/10.3399/bjgp15X687133>.
 57. Lentz V, Montgomery AJ, Georganta K, *et al*. Constructing the health care system in Greece: responsibility and powerlessness. *Br J Health Psych* 2014; **19(1)**: 219–230.
 58. Linzer M, Poplau S, Grossman E, *et al*. A cluster randomized trial of interventions to improve work conditions and clinician burnout in primary care: results from the Healthy Work Place (HWP) Study. *J Gen Intern Med* 2015; **30(8)**: 1105–1111.
 59. Robertson HD, Elliott AM, Burton C, *et al*. Resilience of primary healthcare professionals: a systematic review. *Br J Gen Pract* 2016; DOI: <https://doi.org/10.3399/bjgp16X685261>.
 60. Lemaire JB, Wallace JE. Not all coping strategies are created equal: a mixed methods study exploring physicians' self reported coping strategies. *BMC Health Serv Res* 2010; DOI: 10.1186/1472-6963-10-208.
 61. Matheson C, Robertson HD, Elliott AM, *et al*. Resilience of primary healthcare professionals working in challenging environments: a focus group study. *Br J Gen Pract* 2016; DOI: <https://doi.org/10.3399/bjgp16X685285>.
 62. Holley D, Santos P, Cook J, *et al*. 'Cascades, torrents & drowning' in information: seeking help in the contemporary general practitioner practice in the UK. *Interact Learn Environ* 2016; **24(5)**: 954–967.