Does aerobic exercise reduce postpartum depressive symptoms?

a systematic review and meta-analysis

Abstract

Background

There is currently no specific guidance on the role of exercise in managing postpartum depression in the UK and US, and international guidance is inconsistent.

Aim

To assess the effectiveness of aerobic exercise on postpartum depressive symptoms.

Design and setting

Systematic review and meta-analysis. There was no restriction to study site or setting.

Method

The databases MEDLINE, EMBASE, Cochrane Library, PsycINFO, SportDiscus, Clinical Trials.gov, and the World Health Organization International Clinical Trials Registry Platform were searched. Titles and abstracts, then full-text articles, were screened against inclusion criteria: RCTs measuring depressive symptoms in mothers ≤ 1 year postpartum; and interventions designed to increase aerobic exercise compared with usual care or other comparators. Included studies were assessed using the Cochrane Collaboration's risk of bias tool. Meta-analysis was conducted. Pre-planned subgroup analyses explored heterogeneity.

Results

Thirteen RCTs were included, with 1734 eligible participants. Exercise significantly reduced depressive symptoms when all trials were combined (standardised mean difference -0.44 ; 95% confidence interval $= -0.75$ to -0.12). Exploration of heterogeneity did not find significant differences in effect size between women with possible depression and in general postpartum populations; exercise only and exercise with co-interventions; and group exercise and exercise counselling.

Conclusion

This systematic review provides support for the effectiveness of exercise in reducing postpartum depressive symptoms. Group exercise, participant-chosen exercise, and exercise with co-interventions all may be effective interventions. These results should be interpreted with caution because of substantial heterogeneity and risk of bias.

Keywords

depression; depression, postpartum; exercise; postnatal; review, systematic.

INTRODUCTION

Postpartum depression is a global mental health issue, annually affecting 13 million women worldwide.¹⁻³ Postpartum depression can lead to thoughts of self-harm and/or harm of the child,⁴ and negatively affect children's development.^{5,6} There can be reluctance by women to take antidepressants postnatally, particularly if they are breastfeeding.^{7,8} Cognitive behavioural therapy can be effective but waiting lists are often long in the UK.⁹ Exercise can have a moderate effect in reducing general adult depression (standardised mean difference [SMD] -0.62 , 95% confidence interval [CI] $= -0.81$ to -0.42).¹⁰

Practitioner-led group exercise is recommended by the UK National Institute for Health and Care Excellence (NICE) for persistent subthreshold depressive symptoms and mild-to-moderate depression.¹¹ However, women in the postpartum period have particular physical, practical, and psychological challenges to exercising.^{12,13}

A 2009 review provided some support for exercise as an adjunctive treatment for postpartum depressive symptoms.¹⁴ Despite its challenges, exercise provides another intervention option when many mothers are reluctant to seek treatment and/or take antidepressants.⁸

Exercise also offers improved fitness and weight loss, and opportunities for social interaction at a time when women experience decreased exercise and weight retention.¹⁵⁻¹⁷

Aim and objectives

The aim and objectives of this study were to undertake a meta-analysis of randomised controlled trials (RCTs) investigating the effect of aerobic exercise interventions, compared with usual care, on depressive symptoms in women up to 1 year postpartum. The study aimed to explore the effect of exercise on depressive symptoms in three subgroup analyses:

- women with possible depression and general postpartum populations;
- interventions based only on exercise and those with co-interventions; and
- interventions providing structured group exercise and those supporting participant choice of exercise.

METHOD

The review was registered on PROSPERO: CRD42016047656.

Data sources and searches

The Cochrane Collaboration's guidance was followed on reviews of interventions¹⁸ and PRISMA guidelines.¹⁹ The following bibliographic databases were searched electronically for eligible trials: MEDLINE, EMBASE, the Cochrane Library, PsycINFO, SportDiscus, and Clinical Trials.gov. The World Health Organization (WHO) International Clinical Trials Registry Platform was searched for trials in progress. Searches were conducted for grey literature in Open Grey (<http://opengrey.org>).

RV Pritchett, PhD, research fellow, **AJ Daley**, PhD, reader, behavioural medicine, Institute of Applied Health Research; the Murray Learning Centre; **K Jolly**, PhD, MFPH, professor of public health and primary care, Institute of Applied Health Research, Public Health Building, University of Birmingham, Birmingham.

Address for correspondence

Kate Jolly, Institute of Applied Health Research, Public Health Building, University of Birmingham,

Edgbaston, Birmingham, B15 2TT, UK.

E-mail: c.b.jolly@bham.ac.uk

Submitted: 19 January 2017; **Editor's response:** 27 March 2017; **final acceptance:** 5 April 2017.

©British Journal of General Practice

This is the full-length article (published online 29 Aug 2017) of an abridged version published in print. Cite this version as: **Br J Gen Pract** 2017; DOI: <https://doi.org/10.3399/bjgp17X692525>

How this fits in

UK clinical guidance recommends psychological therapy and antidepressants for postnatal depression. However, women can be reluctant to take antidepressants postnatally and the availability of psychological therapies is often limited. Exercise is effective in treating depression in the general adult population; however, there are particular challenges to be overcome in the postnatal period. This systematic review found that exercise is an effective intervention for reducing postnatal depressive symptoms.

eu/) and OAlster Worldcat (<http://oalster.worldcat.org/>).

Searches were conducted on a range of psychological outcomes; only the results for depression are presented. The following terms were searched as text words and MeSH terms where applicable: exercise, physical activity, postpartum, postnatal, mother, birth, perinatal, depression, anxiety, self-efficacy, quality of life, mother and infant bonding, and child development (the MEDLINE search strategy is available from authors on request). Searches were unrestricted by date or language and undertaken up to September 2016. Bibliographies of eligible studies and reviews were searched for additional trials, and authors contacted if required. Two researchers independently reviewed titles, abstracts, and full-text articles. Eligibility discrepancies were discussed with a third reviewer.

Study selection

Inclusion criteria were:

- populations of mothers <1 year postpartum;
- interventions designed to increase aerobic exercise (activity causing increased heart rate, respiratory rate, and sweating), including those with co-interventions such as social support or nutrition elements;
- comparator groups receiving no care or any form of usual care;
- depressive symptoms measured by questionnaire or diagnostic interview; and
- RCTs.

Exclusion criteria were trials comparing two types of exercise.

Data extraction and quality assessment

Data describing the population, intervention,

and findings were extracted independently by two researchers using standardised, piloted, extraction spreadsheets. Reviewers were not blinded to authors, institution, or journal. Authors were contacted if clarification was required.

Two researchers independently applied the Cochrane Collaboration's risk of bias tool¹⁸ in relation to sequence generation, allocation concealment, blinding of outcome assessors, incomplete outcome data, selective outcome reporting, and other sources of bias.¹⁸ Blinding of personnel and participants to group allocation was not assessed because of the impracticalities of this in exercise trials. Each study was allocated an overall risk of bias: with a low risk of bias being assigned if the risk of bias was deemed to be low in all subdomains; an unclear risk of bias if the risk was considered unclear in one or more subdomains; and a high risk of bias if the risk was considered high in one or more subdomains.¹⁸ Publication bias was assessed by visually inspecting a funnel plot of the SMDs for depressive symptoms.

Data synthesis and analysis

Populations indicating possible depression on screening questionnaires or in the clinical judgement of a health professional are referred to as 'depressed' postpartum populations.

Meta-analyses and meta-regressions were conducted using Review Manager 5.3. An SMD¹⁸ was calculated for continuous outcomes of depressive symptoms. If more than one measure for depression was reported, the primary continuous outcome measure was used. The final point of follow-up was used because the aim was to report long-term outcomes. A weighted mean difference (WMD) was calculated for all trials using the Edinburgh Postnatal Depression Scale (EPDS) to assess depressive symptoms.²⁰ Where the standard deviation of the difference in score was not reported, this was calculated. A correlation of 0.6 was used in the assumptions to estimate standard deviations of differences, based on a previous sample of postpartum women.²¹ A random effects model was used throughout because of known clinical heterogeneity between populations and methodological heterogeneity between interventions.

An initial pooled SMD difference was calculated with 95% CI, with all trials included. Statistical significance was defined as having a 95% CI that did not include zero. Clinical heterogeneity was explored (by qualitatively comparing characteristics

between trials) and statistical heterogeneity using χ^2 tests of heterogeneity and the I^2 statistic.

Three a priori subgroup analyses were performed, investigating whether the effectiveness of exercise in reducing postpartum depressive symptoms varied in relation to the population of women (women with possible depression or general postpartum populations); the presence of co-interventions (exercise only or exercise with co-interventions such as diet or social support); and the context of exercise (exercise groups or participant choice; participant choice often consisted of exercise counselling interventions in which participants typically elected to exercise alone, often by walking). For each hypothesis, the trials were categorised into subgroups; and a random effects meta-regression model was used to determine the effects of exercise in each subgroup and the significance of differences between subgroups.

RESULTS

Trial selection

Of the 9165 records identified after the removal of duplicates, 9043 were excluded after reading titles and abstracts. A further 109 full-text records were excluded based

on the eligibility criteria. Thirteen trials were included in the meta-analysis (Figure 1).

Trial characteristics

All 13 RCTs had been peer reviewed and published as journal articles,^{22-24,26-33} a PhD thesis,²⁵ or abstracts.³⁴ Four trials were conducted in the US,^{25,27,29,31} three in England, UK,^{28,33,34} two in Australia,^{22,26} one in Canada,²³ one in India,³⁰ one in Japan,³² and one in Taiwan.²⁴

The total population of the combined studies was 1734 eligible participants, with follow-up data provided by 1307 participants who were included in the primary meta-analysis. Seven trials recruited participants with possible depression,^{22-25,28,33,34} of which three required participants to score above a threshold on the EPDS questionnaire (from ≥ 10 to >12),^{26,27,32} and two required a diagnosis of depression (International Classification of Diseases version 10 [ICD-10]³³ or Diagnostic and Statistical Manual of Mental Disorders fourth edition [DSM-IV] criteria).³⁴ Despite not reporting baseline depression threshold criteria, two trials reported mean baseline depression scores indicating depressed populations (EPDS scores of 18.9–19.8,²⁵ Beck Depression Inventory [BDI] 15.8–16.9).²⁷ These trials were therefore considered to have ‘depressed’ postpartum populations. Six trials recruited general postpartum populations (Table 1).^{26,27,29-32}

Intervention characteristics

All trials included interventions designed to increase exercise levels. In addition, two trial interventions also aimed to improve diet and encourage a healthy lifestyle,^{24,27} two also provided social support,^{22,31} and one also provided education on postpartum issues.²⁶ Six trials had group exercise interventions,^{22,26,30-32,34} seven trials had interventions in which exercise counselling was provided and the participant was free to choose their own form of exercise (Table 1).^{23-25,27-29,33} The interventions ranged from 4 weeks’ duration^{30,32} to 6 months.^{24,29,33} Most aimed to achieve 30 minutes of moderate activity three to five times weekly,^{22,23,25,27-29,32-34} although some consisted of once-weekly group exercise.^{26,30} Eight of the studies reported adherence to the intended intervention.^{22,23,25,28,29,32-34}

Risk of bias in included studies and publication bias

Six trials were considered at unclear risk^{22,23,28-30,34} and seven at high risk of bias.^{24-27,31-33} Principal factors introducing a risk of bias were a lack of intention-to-

Figure 1. PRISMA flow chart. The 40 records in the top right-hand box are additional papers found.

Table 1. Trial characteristics

Study	Country	Population	N	Intervention	Comparator	Outcomes	Result
Armstrong 2003 ²²	Queensland, Australia	Depressed population	20	Group exercise programme (pram walks) with social co-intervention 12 weeks	Phone support at 6 weeks post-randomisation	EPDS and DASS 6 and 12 weeks	Intervention group depression improved significantly compared with control
Daley 2008 ²⁸	Birmingham, UK	Sedentary population Depressed population	38	Exercise counselling 12 weeks	Usual care Exercise consultation of trial	EPDS 24 weeks	No significant difference in EPDS between groups
Robichaud 2008 ²⁵	Tennessee, US	Depressed population	48	Exercise programme (home-based walking) 6 weeks	Usual activity Intervention after trial	EPDS 8 weeks	No significant difference in EPDS between groups
Da Costa 2009 ²³	Montreal, Canada	Sedentary population Depressed population	88	Exercise counselling 12 weeks	Usual care	EPDS and 17-item HAM-D 3 and 6 months	No significant change in EPDS from baseline to follow-up
Norman 2010 ²⁶	Melbourne, Australia	General postpartum population	161	Group exercise programme with co-intervention (weekly educational session on postpartum issues) 8 weeks	Education on topics unrelated to diet or physical activity Booklet of exercises, list of gyms and community facilities	EPDS 8 and 12 weeks	Significant decrease in EPDS in intervention compared with comparator
Huang 2011 ²⁴	Northern Taiwan	Depressed population	160	Exercise counselling with dietary co-intervention Postpartum group: 6 months	'Standard care' discussions with nurses and leaflets on unrelated topics	BDI 6 months	Significant increase in BDI in intervention and comparator groups
Surkan 2012 ²⁷	Northeast US	General postpartum population	679	Exercise counselling with dietary co-intervention 12 months	Usual WIC care: nutrition assessment, education, food vouchers	CES-D Mean follow-up 13.3 months (SD 3.7)	Modest decrease in CES-D in intervention compared with control. Difference was significant in fully adjusted analysis
Haruna 2013 ³²	Tokyo, Japan	General postpartum population	110	Group exercise programme 4 weeks	No details provided Intervention at end of trial	EPDS 2 and 4 months	No significant difference in EPDS between groups
Thiruppathi 2014 ³⁰	Nellore, India	General postpartum population	45	Group exercise programme 4 weeks	Healthcare education programme. Written materials on nutrition, baby care, posture	EPDS 12 weeks	Significant decrease in intervention group from baseline to follow-up. No significant change in comparator group
Keller 2014 ³¹	A southwestern city, US	General postpartum population	139	Exercise counselling with group exercise (walking) and a co-intervention of 'promotoras' (peer support) 12 weeks	Attention control. Monthly newsletters and weekly phone calls on unrelated topics	EPDS 6 and 12 months	No significant difference between groups
Lewis 2014 ²⁹	Minnesota, US	General postpartum population	130	Exercise counselling 6 months	Counsellor delivered telephone sessions on stress management, sleep, nutrition	SCID-1 at 6 months only PHQ-9 EPDS at 6 months only	Significant decrease in PHQ-9 and EPDS in intervention compared with control

... continued

Table 1 continued. Trial characteristics

Booth 2015 ³⁴	Stoke-on-Trent, UK	Depressed population	22	Group exercise programme (pram walk or facility based) 12 weeks	One discussion about their goals for managing postpartum depression	EPDS 3 and 6 months	No significant difference in EPDS between groups
Daley 2015 ³³	Birmingham, UK	Depressed population	94	Exercise counselling 6 months	Usual care Leaflet on healthy postpartum lifestyle	EPDS 6 and 12 months	Decrease in EPDS in intervention versus comparator, only significant at 6 months not at 12 months

BDI = The Beck Depression Inventory. CES-D = The Center for Epidemiologic Studies Depression Scale. DASS = The Depression Anxiety Stress Scale. EPDS = The Edinburgh Postnatal Depression Scale. HAM-D = The Hamilton Rating Scale for Depression. PHQ-9 = The Patient Health Questionnaire-9. SMD = Standardized Mean Difference. SD = standard deviation. WIC = Special Supplemental Nutrition Program for Women, Infants, and Children.

treat analyses (some studies excluded non-adherent participants), a lack of clarity on selective outcome reporting, a lack of robust sequence generation and concealment of randomisation procedures, and unclear blinding of those conducting outcome assessments and analyses (details of the risk of bias in included studies are available from the authors on request).

The funnel plot was visually inspected for the main analysis including all trials. There appeared to be a lack of smaller studies with results indicating an increase in depression in exercise intervention groups compared with comparator groups. This may be an indication of publication bias (details available from the authors on request).

Provision of additional data

Authors of four trials provided additional data for this review: DaCosta *et al.*,²³ Surkan *et al.*,²⁷ Lewis *et al.*,²⁹ and Boath *et al.*³⁴

Data analysis

Overall effect of exercise on depression scores. Exercise interventions significantly reduced depressive symptoms (SMD -0.44, 95% CI = -0.75 to -0.12, $n = 1307$, $I^2 = 85\%$, 13 trials) (Figure 2).²²⁻³⁴ The WMD was -1.54 EPDS units, 95% CI = -2.97 to -0.12, $n = 652$, $I^2 = 87\%$.^{22,23,25,26,28,30-34}

Population. Exercise interventions had a significant effect in reducing depressive symptoms in 'depressed' postpartum populations (SMD -0.32, 95% CI = -0.63 to -0.00), $I^2 = 55\%$ (Table 2),^{22-25,28,33,34} and in general postpartum populations (-0.57, 95% CI = -1.12 to -0.02, $I^2 = 92\%$) (Table 2).^{26,27,29-32} The effect of exercise interventions in the 'depressed' and general postpartum populations was not significantly different (test for subgroup differences $\chi^2 = 0.62$, $P = 0.43$, $I^2 = 0\%$) [further details available from the authors on request].

Intervention type. Exercise-only interventions had a non-significant effect in reducing depressive symptoms (SMD -0.56, 95% CI = -1.13 to 0.01, $I^2 = 89\%$).^{23,25,28-30,32-34} Exercise with co-interventions had a significant effect on reducing depressive symptoms (-0.35, 95% CI = -0.66 to -0.04, $I^2 = 72\%$).^{22,24,26,27,31} The effect of exercise-only interventions and exercise co-interventions on depressive symptoms was not significantly different (test for subgroup differences: $\chi^2 = 0.41$, $P = 0.52$, $I^2 = 0\%$) (Table 2) [further details available from the authors on request].

Exercise context. Group exercise

interventions had a significant effect in reducing depressive symptoms (SMD difference -1.10, 95% CI = -1.99 to -0.21, $I^2 = 93\%$).^{22,26,30-32,34} Participant choice interventions such as exercise counselling with personal choice of exercise (often exercise alone) had a significant effect in reducing depressive symptoms (-0.20, 95% CI = -0.33 to -0.06, $I^2 = 0\%$).^{23-25,27-29,33} The effects of group exercise and participant choice exercise on depressive symptoms were not significantly different (test for subgroup differences: $\chi^2 = 3.89$, $P = 0.05$, $I^2 = 74\%$) (Table 2) [further details available from the authors on request].

DISCUSSION

Summary

There is no specific guidance on the role of exercise in management of perinatal mental health in the UK or the US.^{7,35} This systematic review provides support for exercise as an effective treatment for reducing postpartum depressive symptoms, whether or not women meet robust criteria for postpartum depression. Additionally, characteristics of exercise interventions have been explored that are most likely to be effective for the postpartum population.

Strengths and limitations

Caution should be taken when interpreting the present results, as a substantial level of heterogeneity was present. Heterogeneity was present in the design of the trials, including exercise-only interventions and those also promoting a healthy diet or social support among peers. Exercise contexts also varied between structured, group-based exercise and tailored exercise counselling with participant choice of exercise. A random effects model was used in analyses to account for this variation, and potential causes of heterogeneity were explored in subgroup analyses; however, there were insufficient numbers of trials to explore this heterogeneity in depth. The last follow-up point was used because the focus of the present study was on long-term outcomes; the median duration of follow-up was 6 months from recruitment (IQR 3.6 months).

The methodological quality of several of the included trials was low. Exclusion of non-adherent participants, insufficiently robust sequence generation, and unclear blinding of outcome assessors resulted in an increased risk of bias. Only two included trials in 'depressed' populations recruited women with a diagnosis of depression (ICD-10 or DSM-IV);^{33,34} the remainder used screening questionnaires such as the

Figure 2. Meta-analysis of the effect of exercise on depressive symptoms (standardised mean difference).
CI = confidence interval. SD = standard deviation.

EPDS²⁰ to indicate women with possible depression. The EPDS has the advantage of being relatively short and simple to complete, but has been reported to have considerable heterogeneity in sensitivity and specificity for detecting postpartum depression across different settings and populations.³⁶

There was a lack of assessment of exercise duration and intensity in included studies; of those that did provide this information, only two included an objective measure of exercise.^{31,33} Exercise intensity can be difficult to determine accurately by self-report. For future research, objective accelerometry would allow for greater accuracy and more meaningful comparison between intervention types.

This review has several strengths. Recommended methods were followed; searching was systematic and not limited by language of publication. The review was restricted to RCTs to reduce the potential for

confounding. The effectiveness of exercise was explored in different populations, intervention types, and exercise contexts, which has not been previously attempted.

Comparison with existing literature

The evidence in this field has increased substantially since the review by Daley *et al* in 2009.¹⁴ Although the primary effect size in this review is smaller than that reported by Daley *et al* (SMD -0.44, 95% CI = -0.75 to -0.12 rather than -0.81, 95% CI = -1.53 to -0.10¹⁴), the CI surrounding the effect is narrower here, indicating more precision.

In this review, the effect of exercise in depressed postpartum populations was smaller than that reported in a recent review of exercise in general adult depressed populations (SMD -0.62, 95% CI = -0.81 to -0.42),¹⁰ indicating that there may be differences in the way general adult populations and postpartum populations

Table 2. Subgroup analyses: population, intervention type, and exercise context

Category	Standardised mean difference	95% CI	P-value	I ²	Number of trials	Number of participants
Depressed postpartum populations	-0.32	-0.63 to -0.00	0.05	55%	7	416
General postpartum populations	-0.57	-1.12 to -0.02	0.04	92%	6	891
Exercise-only interventions	-0.56	-1.13 to 0.01	0.05	89%	8	528
Exercise co-interventions	-0.35	-0.66 to -0.04	0.03	72%	5	779
Group exercise interventions	-1.10	-1.99 to -0.21	0.02	93%	6	406
Participant choice exercise interventions	-0.20	-0.33 to -0.06	0.003	0%	7	901

respond to exercise or differing abilities to adhere to an exercise regime. Women in the postnatal period have the time constraints of new parenthood and barriers to exercise such as fatigue. These factors could be explored by qualitative research with postnatal women.

The effect of exercise on postpartum depressive symptoms reported in this review compared favourably with that reported in a review of low-intensity psychological interventions (such as online cognitive behavioural therapy and self-help literature) versus usual care for depression in the general population (SMD -0.42, 95% CI = -0.55 to -0.29).³⁷ Despite some temporary changes in the composition of breastmilk following maximal exercise, the literature does not suggest any detrimental effects of moderate levels of aerobic exercise,³⁸ as evaluated by the trials in this review. Such interventions are often among the first treatments offered to individuals presenting with depression.¹¹ The effect reported here of exercise on postpartum depressive symptoms is also comparable with those reported for antidepressants in depressed adult populations (SMD -0.49, 95% CI = -0.67 to -0.32 for tricyclic antidepressants and -0.24, 95% CI = -0.35 to -0.12 for selective serotonin reuptake inhibitors compared with placebo).³⁹

Exercise interventions also offer the

additional health benefits of improved physical fitness and weight loss, outcomes particularly relevant to postpartum women, who often experience decreased exercise levels and excess weight.¹⁵⁻¹⁷ The finding that both group exercise and participant choice of exercise (often solitary exercise) may be effective in reducing postpartum depressive symptoms is also noteworthy. Current guidance from NICE on the treatment of postpartum mental illness refers clinicians to exercise advice for the general adult population, recommending only group exercise,¹¹ which may not be the most feasible or acceptable option for women after having a baby.⁴⁰

Implications for practice

National and international guidance on the use of exercise to reduce depressive symptoms in the postpartum period is lacking. Although acknowledging considerable uncertainty about the present findings because of substantial heterogeneity, given the high prevalence of postpartum depression and the potential for exercise to be a low-cost, freely available intervention, aerobic exercise should be considered as a management option for postpartum women with depressive symptoms and as a potential preventative measure more generally in postpartum women.

Funding

This study presents independent research funded by the University of Birmingham, the National Institute for Health Research (NIHR) School for Primary Care Research, and the Collaboration for Leadership in Applied Health Research and Care West Midlands. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health.

Ethical approval

Not applicable.

Provenance

Freely submitted; externally peer reviewed.

Competing interests

All authors declare no support from any organisation for the submitted work; there are no financial relationships with any organisations that might have an interest in the submitted work. Ruth Victoria Pritchett (née Blamey) reports that she is an author of one of the trials in this review, Amanda J Daley reports that she is an author of two of the trials included in this review, and Kate Jolly reports that she is an author of one of the trials included in this review.

Discuss this article

Contribute and read comments about this article: bjgp.org/letters

REFERENCES

- Gavin NI, Gaynes BN, Lohr KN, *et al*. Perinatal depression: a systematic review of prevalence and incidence. *Obstet Gynecol* 2005; **106**(5 Pt 1): 1071–1083.
- Bener A, Gerber LM, Sheikh J. Prevalence of psychiatric disorders and associated risk factors in women during their postpartum period: a major public health problem and global comparison. *Int J Womens Health* 2012; **4**: 191–200.
- World Health Organization. *World health report 2005: make every mother and child count*. Geneva: WHO, 2005.
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th edn. Washington, DC: American Psychiatric Association, 2000.
- Glasheen C, Richardson GA, Fabio A. A systematic review of the effects of postnatal maternal anxiety on children. *Arch Womens Ment Health* 2009; **13**(1): 61–74.
- [No authors listed]. Chronicity of maternal depressive symptoms, maternal sensitivity, and child functioning at 36 months. *Dev Psychol* 1999; **35**(5): 1297–1310.
- National Institute for Health and Care Excellence. *Antenatal and postnatal mental health: clinical management and service guidance*. CG192. London: NICE, 2014. <https://www.nice.org.uk/guidance/cg192> [accessed 26 Jul 2017].
- Turner KM, Sharp D, Folkes L, Chew-Graham C. Women's views and experiences of antidepressants as a treatment for postnatal depression: a qualitative study. *Fam Pract* 2008; **25**(6): 450–455.
- We Need To Talk Coalition. *We need to talk. Getting the right therapy at the right time*. MIND, 2010. <https://www.mind.org.uk/media/280583/We-Need-to-Talk-getting-the-right-therapy-at-the-right-time.pdf> [accessed 3 Aug 2017].
- Cooney GM, Dwan K, Greig CA, *et al*. Exercise for depression. *Cochrane Database Syst Rev* 2013; **9**: CD004366.
- National Institute for Health and Care Excellence. *Depression in adults: recognition and management*. CG90. London: NICE, 2009. <https://www.nice.org.uk/guidance/cg90> [accessed 2 Aug 2017].
- Coates R, Ayers S, de Visser R. Women's experiences of postnatal distress: a qualitative study. *BMC Pregnancy Childbirth* 2014; **14**: 359.
- Hight N, Stevenson AL, Purtell C, Co S. Qualitative insights into women's personal experiences of perinatal depression and anxiety. *Women Birth* 2014; **27**(3): 179–184.
- Daley A, Jolly K, MacArthur C. The effectiveness of exercise in the management of post-natal depression: systematic review and meta-analysis. *Fam Pract* 2009; **26**(2): 154–162.
- Albright C, Maddock JE, Nigg CR. Physical activity before pregnancy and following childbirth in a multiethnic sample of healthy women in Hawaii. *Women Health* 2005; **42**(3): 95–110.
- Fairley L, Santorelli G, Lawlor DA, *et al*. The relationship between early life modifiable risk factors for childhood obesity, ethnicity and body mass index at age 3 years: findings from the Born in Bradford birth cohort study. *BMC Obesity* 2015; **2**: 9.
- Starling AP, Brinton JT, Glueck DH, *et al*. Associations of maternal BMI and gestational weight gain with neonatal adiposity in the Healthy Start study. *Am J Clin Nutr* 2015; **101**(2): 302–309.
- Higgins JPT, Green S, eds. *Cochrane handbook for systematic reviews of interventions*. Version 5.1.0 [updated March 2011]. Cochrane Collaboration, 2011. <http://handbook.cochrane.org> [accessed 14 Aug 2017].
- Liberati A, Altman DG, Tetzlaff J, *et al*. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med* 2009; **6**(7): e1000100.
- Cox J, Holden J, Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry* 1987; **150**: 782–786.
- Ussher M, Lewis S, Aveyard P, *et al*. The London Exercise And Pregnant smokers (LEAP) trial: a randomised controlled trial of physical activity for smoking cessation in pregnancy with an economic evaluation. *Health Technol Assess* 2015; **19**(84): viiixiv, 1135.
- Armstrong K, Edwards H. The effects of exercise and social support on mothers reporting depressive symptoms: a pilot randomized controlled trial. *Int J Ment Health Nurs* 2003; **12**(2): 130–138.
- Da Costa D, Lowensteyn I, Abrahamowicz M, *et al*. A randomized clinical trial of exercise to alleviate postpartum depressed mood. *J Psychosom Obstet Gynaecol* 2009; **30**(3): 191–200.
- Huang TT, Yeh CY, Tsai YC. A diet and physical activity intervention for preventing weight retention among Taiwanese childbearing women: a randomised controlled trial. *Midwifery* 2011; **27**(2): 257–264.
- Robichaud KP. *The effects of an exercise intervention on the psychological well-being of postpartum women*. Graduate School. Middle Tennessee State University. Doctor of Philosophy, 2008. <http://jewlscholar.mtsu.edu/bitstream/handle/mtsu/4064/3322482.pdf?sequence=1&isAllowed=y> [accessed 3 Aug 2017].
- Norman E, Sherburn M, Osborne RH, Galea MP. An exercise and education program improves well-being of new mothers: a randomized controlled trial. *Phys Ther* 2010; **90**(3): 348–355.
- Surkan PJ, Gottlieb BR, McCormick MC, *et al*. Impact of a health promotion intervention on maternal depressive symptoms at 15 months postpartum. *Matern Child Health J* 2012; **16**(1): 139–148.
- Daley A, Winter H, Grimmer C, *et al*. Feasibility of an exercise intervention for women with postnatal depression: a pilot randomised controlled trial. *Br J Gen Pract* 2008; DOI: <https://doi.org/10.3399/bjgp08X277195>.
- Lewis BA, Gjerdingen DK, Avery MD, *et al*. A randomized trial examining a physical activity intervention for the prevention of postpartum depression: the healthy mom trial. *Ment Health Phys Act* 2014; **7**(1): 42–49.
- Thirupathi A, Prasana B, Mastanah E, *et al*. A structured physical activity and health care education beats postpartum depression for primipara mothers: a pilot randomized controlled trial. *Int J Physiother* 2014; **11**(3): 144–151.
- Keller C, Ainsworth B, Records K, *et al*. A comparison of a social support physical activity intervention in weight management among post-partum Latinas. *BMC Public Health* 2014; **14**: 971.
- Haruna M, Watanabe E, Matsuzaki M, *et al*. The effects of an exercise program on health-related quality of life in postpartum mothers: a randomised controlled trial. *Health* 2013; **5**(3): 432–439.
- Daley AJ, Blamey RV, Jolly K, *et al*. A pragmatic randomized controlled trial to evaluate the effectiveness of a facilitated exercise intervention as a treatment for postnatal depression: the PAMPeRS trial. *Psychol Med* 2015; **45**(11): 2413–2425.
- Boath E, Henshaw C, Forsyth J, eds. Exercise as an adjunct therapy for postnatal depression: a pilot study. [Abstract]. The International Marcé Society for Perinatal Mental Health Biennial Scientific Conference. Swansea University, Swansea: Springer Vienna, 2014. *Arch Womens Ment Health* 2015; **18**(2): 269–408.
- Gelenberg AJ, Freeman MP, Markowitz JC, *et al*. *Practice guideline for the treatment of patients with major depressive disorder*. 3rd edn. American Psychiatric Association, 2010. http://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/mdd.pdf [accessed 14 Aug 2017].
- Gibson J, McKenzie-McHarg K, Shakespeare J, *et al*. A systematic review of studies validating the Edinburgh Postnatal Depression Scale in antepartum and postpartum women. *Acta Psychiatr Scand* 2009; **119**(5): 350–364.
- Bower P, Kontopantelis E, Sutton A, *et al*. Influence of initial severity of depression on effectiveness of low intensity interventions: meta-analysis of individual patient data. *BMJ* 2013; **346**: f540.
- Mortensen K, Kam R. Exercise and breastfeeding. *Breastfeed Rev* 2012; **20**(3): 39–42.
- Arroll B, Elley CR, Fishman T, *et al*. Antidepressants versus placebo for depression in primary care. *Cochrane Database Syst Rev* 2009; **3**: CD007954.
- Groth SW, David T. New mothers' views of weight and exercise. *MNC Am J Matern Child Nurs* 2008; **33**(6): 364–370.