

Treatment burden for patients with multimorbidity:

cross-sectional study with exploration of a single-item measure

Abstract

Background

Treatment burden is the effort required of patients to look after their health, and the impact this has on their wellbeing. Quantitative data on treatment burden for patients with multimorbidity are sparse, and no single-item treatment burden measure exists.

Aim

To determine the extent of, and associations with, high treatment burden among older adults with multimorbidity, and to explore the performance of a novel single-item treatment burden measure.

Design and setting

Cross-sectional postal survey via general practices in Dorset, UK.

Method

Patients ≥ 55 years, living at home, with three or more long-term conditions (LTCs) were identified by practices. Treatment burden was measured using the Multimorbidity Treatment Burden Questionnaire. Data collected were sociodemographics, LTCs, medications, and characteristics including health literacy and financial resource. Associations with high treatment burden were investigated via logistic regression. Performance of a novel single-item measure of treatment burden was also evaluated.

Results

A total of 835 responses were received (response rate 42%) across eight practices. Patients' mean age was 75 years, 55% were female ($n = 453$), and 99% were white ($n = 822$). Notably, 39% of patients self-reported fewer than three LTCs ($n = 325$). Almost one-fifth (18%) of responders reported high treatment burden ($n = 150$); making lifestyle changes and arranging appointments were particular sources of difficulty. After adjustment, limited health literacy and financial difficulty displayed strong associations with high treatment burden; more LTCs and more prescribed regular medications were also independently associated. The single-item measure discriminated moderately between high and non-high burden with a sensitivity of 89%, but a specificity of 58%.

Conclusion

High treatment burden was relatively common, underlining the importance of minimising avoidable burden. More vulnerable patients, with less capacity to manage, are at greater risk of being overburdened. Further development of a single-item treatment burden measure is required.

Keywords

cross-sectional studies; general practice; multimorbidity; treatment burden.

INTRODUCTION

Treatment burden is the effort required of patients to look after their health and the impact this has on their functioning and wellbeing.¹⁻³ Recognition of treatment burden in people living with long-term conditions (LTCs) is increasingly important, given the ageing populations of many countries, and increasing prevalence of LTCs and multimorbidity.^{4,5} The workload of health care can include ordering and taking medications, organising and attending healthcare appointments, monitoring health conditions, and modifying lifestyle behaviours.⁶ For people living with multimorbidity, clinicians may be following multiple single-disease guidelines, an important driver of polypharmacy and potentially of treatment burden.^{7,8} In the UK, the National Institute for Health and Care Excellence 2016 guidance for the assessment and management of multimorbidity⁹ advises taking an approach to care that includes 'improving quality of life by reducing treatment burden'.

Negative impacts of treatment burden are various and include those of both a psychological and practical nature, such as interference with daily activities, negative emotions, and strained relationships.^{10,11}

Patients may seek to minimise disruptions not only through adaptation, but also by non-adherence to treatment.^{10,12} When thinking about treatment burden, it is important to consider patient 'capacity': the abilities, resources, and readiness to address the combined demands of treatment workload and daily life.¹³ Components of capacity may include socioeconomic resources such as financial wellbeing and support networks, health literacy, relevant knowledge and experience, and physical and mental functioning.^{13,14} When the workload exceeds available capacity, patients may be described not only as experiencing high treatment burden, but also as being overburdened — with the potential risk of involuntary non-adherence, disruption to care, and adverse health outcomes.^{13,15} A degree of treatment burden is perhaps inevitable in managing LTCs, but seeking to minimise unnecessary burden is important, as encapsulated by 'minimally disruptive medicine'.¹⁶

Validated instruments to measure patient-perceived treatment burden are available.¹⁷⁻²¹ However, quantitative data on the extent of, and factors associated with, high treatment burden for patients with multimorbidity are limited. Measuring treatment burden is not

JE Morris, MSc, MFPH, specialty registrar in public health; **PJ Roderick**, MD, professor of public health; **S Harris**, MSc, associate professor of medical statistics; **SDS Fraser**, DM, MRCP, FFPH, associate professor of public health, School of Primary Care, Population Sciences and Medical Education, Faculty of Medicine, University of Southampton, Southampton General Hospital, Southampton. **G Yao**, PhD, professor in health economics, Department of Health Sciences, University of Leicester, Leicester. **S Crowe**, MSc, FFPH, director of public health for Dorset Council and for Bournemouth, Christchurch and Poole Council; **D Phillips**, MB, FFPH, SFFMLM, former director of public health, Public Health Dorset, Dorchester. **P Duncan**, MPH, MRCP, GP and senior clinical research fellow, Centre for Academic Primary Care, University of Bristol, Bristol.

Address for correspondence

James E Morris, School of Primary Care, Population Sciences and Medical Education, Faculty of Medicine, University of Southampton, Southampton General Hospital, Southampton SO16 6YD, UK.

Email: j.e.morris@soton.ac.uk

Submitted: 24 September 2020; **Editor's response:**

21 October 2020; **final acceptance:**

26 November 2020.

©The Authors

This is the full-length article (published online 20 Apr 2021) of an abridged version published in print. Cite this version as: **Br J Gen Pract 2021;**

DOI: <https://doi.org/10.3399/BJGP.2020.0883>

How this fits in

Patients with multimorbidity may experience treatment burden from healthcare demands, but the extent of, and factors associated with, high treatment burden are not well understood. In this survey of older adults with multimorbidity, almost one-fifth reported 'high' treatment burden, which was associated not only with a greater number of long-term conditions and more prescribed regular medications, but also with characteristics indicating reduced 'capacity' to manage, namely limited health literacy and financial difficulty. GPs have a central role in incorporating these considerations into patient care, to try to ensure that patients are not overburdened. Additionally, existing measures of treatment burden are too time consuming for clinical use. A novel single-item measure of treatment burden was explored here, but performed moderately; further development of such a measure is therefore required.

currently part of routine care and existing measures are too long for time-constrained clinical encounters, particularly in UK general practice. The aims of this study were to determine the extent of treatment burden, and explore the characteristics associated with high treatment burden, among adults ≥ 55 years with three or more specified LTCs documented in their GP records; and to explore the performance of a novel single-item treatment burden measure in the same sample.

METHOD

Survey design and sample

A cross-sectional postal survey of older adults with multimorbidity was conducted in Dorset, England, between February and July 2019. Patients aged ≥ 55 years, living at home, and with three or more LTCs from a specified list, were identified for invitation from GP registers. Inclusion and exclusion criteria are shown in Box 1. Quality and Outcomes Framework clinical code clusters²² defined 14 LTCs, with Read codes²³ (structured clinical terms) defining a further five. The 19 LTCs selected were common, readily identified from GP records, and represented a range of body systems. Conditions considered distinct regarding the likely avoidability, or impact on perception, of treatment burden (such as cancer or severe mental health diagnoses) were not included. The cut-off of ≥ 55 years defining 'older' adults was lower than a more orthodox threshold of ≥ 65 years because more deprived groups may experience multimorbidity at younger ages.²⁴ Patients' treatment burden was measured using the 10-item Multimorbidity Treatment Burden Questionnaire (MTBQ):²¹ a concise, simply worded instrument suitable for self-completion, with good coverage of burden domains and validated in a multimorbid population similar to that in the current study. These characteristics conferred greatest suitability relative to other candidate measures,^{17–20} which variously exhibited limitations such as wording complexity, narrowness of focus, or considerable length. Each MTBQ item is scored as follows: 0 (not difficult/does not apply), 1 (a little difficult), 2 (quite difficult), 3 (very difficult), and 4 (extremely difficult); for those completing five or more items, average item score is multiplied by 25 to yield a global score of 0–100. Treatment burden is categorised as none (global score 0), low (>0 and <10), medium (≥ 10 and <22), or high (≥ 22).

Self-reported data on a range of variables were collected through the survey, in areas

Box 1. Inclusion and exclusion criteria for patients in this study

Inclusion criteria

- Aged ≥ 55 years
- Registered at a participating Dorset GP practice
- Diagnosed with ≥ 3 of the following long-term conditions, as recorded in the GP record (considered as separate conditions):^a
 1. Atrial fibrillation
 2. Coronary heart disease
 3. Heart failure
 4. Hypertension
 5. Peripheral arterial disease
 6. Stroke or transient ischaemic attack
 7. Diabetes
 8. Asthma
 9. Chronic obstructive pulmonary disease
 10. Depression
 11. Chronic kidney disease
 12. Epilepsy
 13. Osteoporosis
 14. Rheumatoid arthritis
 15. Parkinson's disease
 16. Multiple sclerosis
 17. Inflammatory bowel disease
 18. Coeliac disease
 19. Osteoarthritis

Additionally:

- Only one person from a given household was eligible to participate
- Participating GP practices used the 'SystmOne' electronic medical record system^b

^aLong-term conditions numbered 1–14 are Quality and Outcomes Framework (QOF) conditions; those numbered 15–19 are non-QOF conditions. ^bSystmOne is the electronic GP record system predominantly used in practices throughout Dorset.

Exclusion criteria

- Resident in a care home
- In receipt of palliative care
- Diagnosed with a severe mental health condition (psychosis, schizophrenia, bipolar disorder)
- Diagnosed with dementia
- Known to have active cancer (recorded within the last 3 years)
- Lacked mental capacity to participate
- Expressed wish not to participate in research
- Other reason, rendering a patient unsuitable to participate, at the discretion of a healthcare professional at the GP practice




Figure 1. Survey flowchart: recruitment, invitation, and response.^aSeven practices invited 250 people; one practice invited 233 people.

including sociodemographics, prescribed regular medications, specific LTCs (corresponding to survey inclusion criteria), travel for health care, recent healthcare resource use, and health status/quality of life. Home ownership (dichotomised as homeowner/non-homeowner) served to measure socioeconomic status. Characteristics indicative of capacity were captured, including financial resource via the perceived level of difficulty in meeting the financial costs of health care (on a 5-point Likert scale from 'no' to 'extreme' difficulty), and health literacy via the Single Item Literacy Screener — which asks about the perceived frequency of needing help to read health-related written material (on a 5-point Likert scale, with descriptors 'sometimes', 'often', and 'always' indicating limited health literacy).²⁵

A novel single-item measure of treatment burden was also explored: 'On a scale of 0–10, where 0 is no effort and 10 is the highest effort you can imagine, how would you rate the amount of effort you have to put in to manage your health conditions?', with responders circling along a number-line. This measure did not undergo formal development; however, the wording and format embodied principles of existing measures of self-rated health (the EQ visual analogue scale)²⁶ and pain (the numerical rating scale for pain).²⁷ Additionally, MTBQ unidimensionality lent legitimacy to trialling a measure comprised of a single item.

Based on a 26.6% prevalence for high treatment burden from MTBQ validation data,²¹ identifying those with high burden would require at least 300 survey responses

for a maximum 95% confidence interval (CI) width of $\pm 5\%$, considered sufficiently precise. nQuery (version 7.0) was used for the sample size calculation.

Survey documents were refined using a 'think-aloud' procedure (a cognitive-interviewing approach for pre-testing self-completion questionnaires),²⁸ conducted with colleagues and a lay representative (see Supplementary Box S1 for details of survey questions and response options). This article reports on a subset of data and analyses.

Recruitment, invitation, and response

Recruitment of GP practices sought eight geographically dispersed, socioeconomically diverse sites across Dorset. Each participating practice identified survey invitees by electronically searching their patient list with a supplied algorithm, and manually screening a random sample of resulting records — identifying at most 250 invitees meeting the inclusion/exclusion criteria. Reasons for manual exclusions were requested.

Practices posted survey packs to the invitees that comprised a personalised invitation letter, information sheet, consent form, survey booklet, and return envelopes. Freepost returns, GP practice endorsement, an uncomplicated means of participation, and a survey designed to minimise completion burden were used to maximise response rate, along with an online response platform offered via a weblink. Participants provided written (or online equivalent) consent; and could optionally consent to GP-record data sharing with the study team, and to receive a potential follow-up survey (both outwith the scope of this article).

Statistical analyses

Survey data underwent manual database input, with a 5% double-entered sample facilitating error-rate estimation. SPSS Statistics (version 24) was used for analysis. Descriptive statistics were used to explore characteristics of the responder sample, non-response bias based on age and sex, and distribution of treatment burden. A binary outcome of high treatment burden versus no/low/medium burden combined was used (this dichotomy lending a focus to the most impacted group). Other variables acted as exposures. Univariable and multivariable logistic regression were used to identify associations with high treatment burden. Variables were considered for inclusion in a final multivariable model if deemed clinically important a priori or

Table 1. Characteristics of survey responders (all data self-reported)

Variable ^a		Number	% (of N for variable)
Age in years, mean (SD)		75 (8.6)	
Age category in years	55–59	40	4.8
<i>N</i> = 827	60–64	58	7.0
Missing <i>n</i> = 8 (1.0%)	65–69	125	15.1
	70–74	175	21.2
	75–79	173	20.9
	80–84	136	16.4
	85–89	83	10.0
	90–94	34	4.1
	≥95	3	0.4
Sex	Male	376	45.4
<i>N</i> = 829	Female	453	54.6
Missing <i>n</i> = 6 (0.7%)			
Ethnicity	White	822	99.2
<i>N</i> = 829	Other than white ^b	7	0.8
Missing <i>n</i> = 6 (0.7%)			
Marital status	Married or in a partnership	520	63.0
<i>N</i> = 826	Widowed	180	21.8
Missing <i>n</i> = 9 (1.1%)	Divorced or dissolved partnership	94	11.4
	Single ^c	32	3.9
Living situation	Cohabiting ^d	565	68.3
<i>N</i> = 827	Lives alone	262	31.7
Missing <i>n</i> = 8 (1.0%)			
Home ownership	Home owner ^e	654	79.3
<i>N</i> = 825	Non-home owner	171	20.7
Missing <i>n</i> = 10 (1.2%)			
Employment status	Retired	716	86.6
<i>N</i> = 827	Employed (full/part time)	67	8.1
Missing <i>n</i> = 8 (1.0%)	Unemployed	22	2.7
	Other	22	2.7
Smoking status	Current smoker	44	5.3
<i>N</i> = 826	Ex-smoker	440	53.3
Missing <i>n</i> = 9 (1.1%)	Never smoked	342	41.4
Number of long-term conditions^f	0	7	0.8
<i>N</i> = 824	1	95	11.5
Missing <i>n</i> = 11 (1.3%)	2	223	27.1
	3	255	30.9
	4	138	16.7
	5	69	8.4
	≥6	37	4.5
Medications prescribed	0–3	130	15.7
(number on repeat)	4–6	315	38.1
<i>N</i> = 826	7–9	204	24.7
Missing <i>n</i> = 9 (1.1%)	10–14	131	15.9
	≥15	46	5.6
Treatment burden category	High	150	18.0
(Measure: MTBQ)	Medium	224	26.9
<i>N</i> = 833	Low	268	32.2
Missing <i>n</i> = 2 (0.2%)	None	191	22.9

shown to be statistically significant (at $P < 0.05$) at the univariable stage. Variables for healthcare resource use (number of GP and outpatient appointments in the previous 6 months) were excluded given

the risk of poor recall and greater amounts of missing data; travel time to hospital was also excluded since it was not relevant for all responders. The final, parsimonious model was mutually adjusted for age, sex, marital status, home ownership, number of LTCs, number of prescribed regular medications, health literacy, and financial difficulty.

For the single-item measure, the sensitivity, specificity, positive predictive value, and negative predictive value at each integer number-line cut-off were calculated. 'Correct' identification of high treatment burden was determined relative to the MTBQ as a reference standard. A receiver operating characteristic curve was plotted, and area under the curve computed to evaluate the ability of the single-item measure to discriminate between high and non-high treatment burden.

RESULTS

Eight practices were recruited, which exhibited heterogeneity in geographic location (including rurality/urbanity), deprivation level, and list size (see Supplementary Table S1 for details, which also presents practice demographics relative to national data). Manual exclusion of potentially eligible patients most commonly occurred at practice discretion (data from seven sites). In total, practices posted out 1983 surveys; 835 usable responses were received (response rate 42.1%), the vast majority by post ($n = 808$, 96.8%) (Figure 1).

With the exception of the questions relating to healthcare resource use, there were minimal missing data. A data entry error rate of <0.1% was estimated, which was deemed acceptably low. Age and sex distributions of the responder sample closely matched those of the invited sample (see Supplementary Table S2 for details). Responder characteristics are shown in Table 1. The mean age of responders was 75 years, 54.6% ($n = 453$) were females, and 99.2% ($n = 822$) were of white ethnicity. Responders were mostly retired ($n = 716$, 86.6%); many were married/in a civil partnership ($n = 520$, 63.0%); and most owned their own home ($n = 654$, 79.3%).

A substantial minority ($n = 325$, 39.4%) self-reported fewer than three survey-specified LTCs, despite the presence of three or more such conditions in the GP record being an inclusion criterion (Table 1). The most frequently self-reported LTCs were hypertension ($n = 531$, 64.4%), osteoarthritis ($n = 331$, 40.2%), and type 2 diabetes ($n = 248$, 30.1%) (see Supplementary Table S3 for details). Polypharmacy was common, with

Table 1 continued. Characteristics of survey responders (all data self-reported)

Variable ^a		Number	% (of N for variable)
Health literacy (frequency of needing reading help) (Measure: SILS) N = 830 Missing n = 5 (0.6%)	Never	554	66.7
	Rarely	137	16.5
	Sometimes	73	8.8
	Often	34	4.1
	Always	32	3.9
Financial difficulty with health care N = 828 Missing n = 7 (0.8%)	Not difficult or n/a	609	73.6
	A little	146	17.6
	Quite	44	5.3
	Very	24	2.9
	Extreme	5	0.6
Travel time to hospital N = 760 Missing or n/a, n = 75 (9.0%)	≤1 hour	707	93.0
	>1 hour	53	7.0
Travel time to GP N = 826 Missing n = 9 (1.1%)	≤10 minutes	505	61.1
	>10 minutes	321	38.9
Number of outpatient appointments in last 6 months N = 761 Missing n = 74 (8.9%)	0–2	605	79.5
	≥3	156	20.5
Number of GP appointments in last 6 months N = 724 Missing n = 111 (13.3%)	0–2	466	64.4
	≥3	258	35.6

^aMissing percentages are based on overall survey responder denominator of 835. ^b'Other than white' ethnicity comprises categories of 'Mixed/multiple ethnic groups', 'Asian/Asian British', 'Black/African/Caribbean/black British', or 'Other ethnic group', combined. ^cSingle means never married or in a civil partnership. ^dCohabiting means living with any of spouse/partner, child(ren), or other person. ^eHome owner means owning or jointly owning one's current home, including having a mortgage. ^fFor number of long-term conditions (LTCs): if responders had not ticked any LTC survey tick-box and not provided LTC freetext in the survey, LTC data are considered missing; if responders had not ticked any LTC tick-box but had provided some freetext (that did not itself indicate a survey-specified LTC), their LTC count is considered 0. MTBQ = Multimorbidity Treatment Burden Questionnaire. n/a = not applicable. p'ship = civil partnership. SILS = Single Item Literacy Screener.

46.1% of responders (n = 381) prescribed seven or more medications (Table 1).

Some financial difficulty with health care (level of difficulty described as 'a little', 'quite', 'very', or 'extreme') was reported by 26.4% of responders (n = 219), and limited health literacy was reported by 16.7% (n = 139). The number of GP appointments was recalled as being three or more over the previous 6 months by 35.6% of responders (n = 258), and the number of hospital outpatient appointments as three or more over the same time period by 20.5% of responders (n = 156). For both these types of healthcare resource, responders being unable to remember/provide the number of appointments attended (despite indicating at least one attendance) was the principal reason for missing data.

Distribution of global score for the 833 MTBQ responders is shown in Figure 2. 'High' treatment burden was reported by 18.0% (n = 150), 'medium' by 26.9% (n = 224), and 'low' by 32.2% (n = 268); no burden was reported by 22.9% (n = 191) (Table 1). Treatment burden domains in which responders most commonly reported 'some difficulty' (MTBQ item score >0) were: 'making recommended lifestyle changes (such as diet/exercise)' with 47.0% (n = 390) reporting difficulty; and 'arranging appointments with health professionals' (39.4%, n = 327) (see Supplementary Table 4 for details).

Associations with high treatment burden

On univariable analysis, high treatment burden was associated with younger age; female sex; non-home-ownership; greater number of LTCs; more prescribed regular medications; limited health literacy; financial difficulty with health care; longer travel time to hospital; and more outpatient appointments, and GP appointments, in the previous 6 months.

The final multivariable model was mutually adjusted for age, sex, marital status, home ownership, number of LTCs, number of prescribed regular medications, health literacy, and financial difficulty. Strong independent associations (at $P < 0.001$) were observed for both limited health literacy (odds ratio [OR] 3.64, 95% CI = 2.24 to 5.92) and financial difficulty (OR 3.94, 95% CI = 2.56 to 6.07), both being dichotomised variables. Independent associations were also seen for greater number of LTCs (overall $P = 0.008$ with ORs reaching significance at five or more LTCs, for example, OR 2.98 with 95% CI = 1.13 to 7.84 for 5 LTCs versus ≤1), and more prescribed regular medications ($P = 0.04$, OR 1.62, 95% CI = 1.03 to 2.55, for ≥7 medications versus <7) (Table 2).

Single-item treatment burden measure

Setting a threshold of ≥5 for high treatment burden on the single-item measure (optimising its performance) yielded a sensitivity of 89%, specificity of 58%, positive predictive value of 31%, and negative predictive value of 96%. See Supplementary Table 5 for details of the underlying data for the 826 responders. Figure 3 displays the receiver operating characteristic curve; the area under the curve was 0.77 (95% CI = 0.73 to 0.81).

DISCUSSION

Summary

This cross-sectional study of older adults with multimorbidity identified that high




Figure 2. Distribution of treatment burden (MTBQ global score) among survey responders.^a
^aGlobal score 0 indicates no treatment burden; global score cut-offs of ≥ 10 and ≥ 22 (out of 100), indicated by the vertical dashed lines, facilitate categorisation of responders into those with 'low', 'medium', or 'high' treatment burden. MTBQ = Multimorbidity Treatment Burden Questionnaire.

self-reported treatment burden was reported by almost one-fifth of participants. Making recommended lifestyle changes and arranging appointments with health professionals frequently contributed to the burden. High treatment burden was strongly associated with limited health literacy and reported financial difficulty with health care; a greater number of LTCs and more prescribed regular medications were also independently associated. These findings imply that more vulnerable patients, with less capacity to manage, are at particular risk of being overburdened by healthcare demands. A novel single-item measure of treatment burden was explored, but performed moderately: further development of such a measure is required.

Strengths and limitations

Strengths of this study include the number of responders considerably exceeding the minimum target sample size, generally little missing data, and virtually no non-response bias in age or sex. Regarding socioeconomic status, the proportion of homeowners, at 79%, is comparable to the 77% observed in those aged ≥ 55 years nationally (English Housing Survey 2018 to 2019 data),²⁹ and the predominantly postal response supports generalisability beyond just those responders who are computer literate. The survey captured numerous characteristics including those indicative

of patient capacity. Investigation of a single-item treatment burden measure was novel to the authors' knowledge.

However, there are important limitations. Given the cross-sectional design, the study's findings represent association, not causation. As is often seen with postal surveys, the absolute response rate was low (42%); while low response rate is not synonymous with non-response bias,³⁰ characteristics of non-responders could not be assessed beyond age and sex. The proportion of patients with high treatment burden may be an underestimate if those experiencing the highest levels of burden were, as a consequence, less likely to respond. If non-responders also had lower levels of health literacy (which is plausible), the positive association identified between limited health literacy and high treatment burden might be underestimated. Socioeconomic status was captured only via the proxy measure of home ownership: a limitation, although home ownership has been shown to have some value as a marker of socioeconomic status, particularly among older people in Britain.³¹ Unaccounted confounding may also be occurring, for example, through the potential influence of carer burden or functional incapacity among responders; importantly, individual LTC severity was also not measured. The Single Item Literacy Screener captured patient-perceived reading ability: a central component of, but not wholly characterising, the multifaceted concept of health literacy.³² Despite home ownership levels of responders being comparable to those in England, caution is required before generalising findings from this Dorset study to other areas, given the below-national average deprivation profile of participating practices and predominantly white ethnicity of the sample.

Almost 40% of responders self-reported fewer than three LTCs, at variance with their GP record. This could reflect limited recall or lack of awareness if diagnoses were not discussed by clinicians; this is well recognised in some conditions such as chronic kidney disease.³³ Imprecision associated with electronic patient record system coding may also be a contributory factor.³⁴ The association of treatment burden with LTC count should therefore be interpreted with caution, but is highly plausible.

The single-item treatment burden measure is acknowledged as entirely exploratory and was not subject to formal development. The item's discriminatory ability, based on the area under the curve, was fair; specificity and positive predictive

Table 2. Univariable and multivariable associations with high treatment burden (versus non-high treatment burden)

Characteristic ^a		Univariable			Multivariable ^b		
		OR	95% CI	P-value	OR	95% CI	P-value
Age category in years (vs. 55–64)	65–74	0.60	0.36 to 1.01	0.010	0.82	0.44 to 1.56	0.268
	75–84	0.39	0.23 to 0.68		0.55	0.28 to 1.07	
	≥85	0.56	0.30 to 1.06		0.64	0.27 to 1.48	
Sex (vs. male)	Female	1.45	1.01 to 2.08	0.046	1.40	0.91 to 2.18	0.130
Marital status (vs. married) ^c	Single	1.52	0.64 to 3.63	0.348	1.32	0.49 to 3.57	0.579
	Divorced ^d	1.86	1.10 to 3.13	0.020	1.66	0.88 to 3.11	0.117
	Widowed	1.45	0.94 to 2.23	0.090	1.37	0.76 to 2.47	0.298
Living situation (vs. cohabiting)	Lives alone	1.15	0.79 to 1.68	0.461	—		
Home ownership (vs. home owner)	Non-home owner	2.32	1.56 to 3.43	<0.001	1.34	0.82 to 2.19	0.239
Employment status (vs. employed)	Unemployed	1.98	0.70 to 5.61	0.198	—		
	Retired	0.71	0.39 to 1.31	0.272			
	Other	0.77	0.23 to 2.63	0.677			
Smoking (vs. never smoked)	Ex-smoker	0.88	0.61 to 1.26	0.478	—		
	Current smoker	1.25	0.59 to 2.67	0.558			
Number of long-term conditions (vs. 0 or 1 condition)	2	1.25	0.56 to 2.79	<0.001	1.06	0.44 to 2.54	0.008
	3	2.41	1.13 to 5.11		1.91	0.83 to 4.39	
	4	2.42	1.08 to 5.42		1.21	0.50 to 3.00	
	5	6.25	2.70 to 14.47		2.98	1.13 to 7.84	
	≥6	8.78	3.43 to 22.52		4.38	1.42 to 13.53	
Medications prescribed (number on repeat) (vs. <7 medications)	≥7	2.66	1.84 to 3.86	<0.001	1.62	1.03 to 2.55	0.038
Health literacy (vs. not limited)	Limited	4.92	3.29 to 7.36	<0.001	3.64	2.24 to 5.92	<0.001
Financial difficulty with health care (vs. not difficult or n/a)	Some difficulty	5.18	3.55 to 7.54	<0.001	3.94	2.56 to 6.07	<0.001
Travel time to hospital (vs. ≤1 hour)	>1 hour	2.24	1.22 to 4.11	0.009	—		
Travel time to GP (vs. ≤10 minutes)	>10 minutes	1.31	0.91 to 1.89	0.140	—		
Outpatient appointments in the last 6 months (vs. 0–2)	≥3	2.57	1.71 to 3.86	<0.001	—		
GP appointments in the last 6 months (vs. 0–2)	≥3	3.14	2.12 to 4.64	<0.001	—		

^aAll characteristics are self-reported. ^bAdjusting for age, sex, marital status, home ownership, number of long-term conditions, number of prescribed regular medications, health literacy, and financial difficulty. ^cMarried is used to refer to the category 'Married or in a civil partnership'. ^dDivorced is used to refer to the category 'Divorced or dissolved civil partnership'. CI = confidence interval. n/a = not applicable. OR = odds ratio (for high treatment burden).

value were low, but sensitivity and negative predictive value were high, for a number-line threshold ≥ 5 , suggesting some utility in 'ruling out' high treatment burden, rather than in screening for high burden *per se*. It is possible that the position of the single-item relative to other questions (the item was located part-way through the survey) influenced responses to the item. Further

development is clearly needed; current item formulation may provide a starting point for iterative work with patient groups.

Comparison with existing literature

Distribution of treatment burden was comparable to that observed during MTBQ validation in the 3D study.^{21,35} 3D exhibited a greater proportion with high burden




Figure 3. Receiver operating characteristic curve for the exploratory single-item treatment burden measure.

(27% versus 18%), potentially explained by a likely more multimorbid sample (from grouping of similar LTCs within selection criteria, and inclusion of severe mental health conditions), and inclusion of younger participants (≥ 18 years versus ≥ 55 years) who tended to report greater burden. An inverse age–burden relationship was also observed among European cohorts with chronic conditions^{36,37} (and here, although non-significant after adjustment); this could perhaps result from balancing employment demands or caring responsibilities with LTC management, or denote differing healthcare-related expectations with age. This study did not identify an independent association with female sex that has been noted previously.²¹ However, as here, during MTBQ development an association between the number of LTCs and high treatment burden was observed;²¹ such a relationship is consistent with presumed connections between multimorbidity and treatment burden.

Making recommended lifestyle changes and arranging appointments with health professionals were the domains most commonly generating ‘some difficulty’ – likewise observed during MTBQ

development.²¹ This consistent finding could direct burden-reduction initiatives.

Correlation between high treatment burden and markers of workload (more pharmacological treatments, more LTCs) was observed in a Swiss cohort with multimorbidity; however, this was for GP-assessed burden compared with patient-perceived burden.³⁷ Qualitative work indicates medications can contribute to burden (for example, by interfering with activities, or requiring coordination),³⁸ consistent with the association identified in this study with more prescribed regular medications (independent of number of LTCs).

Non-home-ownership was associated with high treatment burden on univariable analysis; in the multimorbid Swiss cohort, socioeconomic deprivation was independently associated with high burden.³⁷ A Danish study of 2111 people with cardiovascular disease and multimorbidity also found greater treatment burden among those with difficulty understanding health information³⁹ – consistent with the findings of the current study, and suggesting that enhancing health literacy (and thereby capacity) might mitigate burden.

Associations with limited health literacy and financial difficulty accord with Shippee *et al*'s¹³ ‘cumulative complexity model’: lower levels of capacity to manage workload relating to greater burden. A mixed-methods study in the US, however, found that capacity indicators were seldom documented in medical records,⁴⁰ implying under-recognition of capacity issues by clinicians: a potential barrier to prospective capacity-enhancing interventions.

Patients with multimorbidity report various strategies to lessen perceptions of burden, for example, maintaining a positive attitude or normalising self-care, and, notably, drawing on positive aspects of health care.⁴¹ Indeed, provider communication and interpersonal skills (‘relational quality’) have been correlated with lower treatment burden.⁴² Qualitative work with stroke professionals similarly identified communication and coordination, as part of individualised care, as vital to reducing treatment burden.⁴³ This links importantly to the current study because of the finding that limited health literacy was associated with high treatment burden.

Implications for research and practice

Almost one-fifth of older adults with multimorbidity in this study reported high treatment burden, underlining the importance of recognising, and seeking

Funding

Funding was provided by Public Health Dorset and the University of Southampton to support James E Morris in undertaking this work.

Ethical approval

Ethical approval was granted by the University of Southampton (ERGO reference 31695) and the London – City & East Research Ethics Committee together with the Health Research Authority and Health and Care Research Wales (REC reference 18/LO/2004, IRAS reference 242816).

Provenance

Freely submitted; externally peer reviewed.

Competing interests

Polly Duncan is an investigator for an industry–university collaboration study between the University of Bristol and Pfizer investigating the burden of lower respiratory tract infection and community-acquired pneumonia in primary care. The other authors have declared no competing interests.

Acknowledgements

The study team are grateful to all survey responders, participating practices, and colleagues at NHS Dorset Clinical Commissioning Group (CCG) who supported this work. A debt of thanks in particular is due to Carmen Taylor for her input. The study team would like to acknowledge the contribution of Dr Craig Wakeham, chief clinical information officer at NHS Dorset CCG and former Dorset GP, who made a significant contribution to facilitating and supporting this work and who tragically passed away in April 2020 after testing positive for coronavirus.

Open access

This article is Open Access: CC BY 4.0 licence (<http://creativecommons.org/licenses/by/4.0/>).

Discuss this article

Contribute and read comments about this article: bjgp.org/letters

to minimise, avoidable burden. While this study has focused on those with three or more documented LTCs, there remains the potential for patients with fewer conditions to also experience treatment burden. Further work is needed to investigate changes in treatment burden over time, and the impact of workload and capacity on perceived burden, adherence, and health outcomes. Research evaluating the impact of interventions that might reduce workload or enhance capacity would also be beneficial. As of 2020, such enquiry may be facilitated by innovations stimulated by the COVID-19 pandemic, including the introduction of potentially more sustainable models of healthcare delivery.⁴⁴ Social prescribing,⁴⁵ now increasingly available via primary care, might contribute to bolstering patient capacity, for example, by enhancing support networks or facilitating improvement in wellbeing. Principles of 'minimally disruptive medicine' will be key to reducing burden, including care coordination, development of clinical guidelines tailored to comorbidity, and prioritisation of care from the patient perspective.¹⁶ System-level

healthcare solutions are required, not only those applicable at individual patient level.

The ability to assess treatment burden swiftly and accurately is nevertheless key to optimising care, hence a single-item measure could ultimately find utility in clinical settings. Given the moderate performance of the single-item treatment burden measure explored in this study, this item should not yet be adopted into practice: further development is recommended.

Clinicians should be alert to potentially overburdened patients. The study findings suggest that, in addition to those with a greater number of LTCs and more prescribed regular medications, patients with limited health literacy and fewer financial resources are at increased risk of high treatment burden. Such factors indicate increased vulnerability and reduced capacity to manage the work of looking after one's health. GPs have a central role in incorporating these considerations into patient care for those with multimorbidity, to ensure that patients are not overburdened with healthcare demands.

REFERENCES

- Eton DT, Ramalho de Oliveira D, Egginton JS, *et al*. Building a measurement framework of burden of treatment in complex patients with chronic conditions: a qualitative study. *Patient Relat Outcome Meas* 2012; **3**: 39–49.
- Mair FS, May CR. Thinking about the burden of treatment. *BMJ* 2014; **349**: g6680.
- Sav A, King MA, Whitty JA, *et al*. Burden of treatment for chronic illness: a concept analysis and review of the literature. *Health Expect* 2015; **18**(3): 312–324.
- Kingston A, Robinson L, Booth H, *et al*. Projections of multi-morbidity in the older population in England to 2035: estimates from the Population Ageing and Care Simulation (PACSim) model. *Age Ageing* 2018; **47**(3): 374–380.
- United Nations, Department of Economic and Social Affairs, Population Dynamics. *World population prospects 2019*. <https://population.un.org/wpp/> [accessed 6 Apr 2021].
- Tran VT, Barnes C, Montori VM, *et al*. Taxonomy of the burden of treatment: a multi-country web-based qualitative study of patients with chronic conditions. *BMC Med* 2015; **13**: 115.
- Hughes LD, McMurdo MET, Guthrie B. Guidelines for people not for diseases: the challenges of applying UK clinical guidelines to people with multimorbidity. *Age Ageing* 2013; **42**(1): 62–69.
- Buffel du Vaure C, Ravaud P, Baron G, *et al*. Potential workload in applying clinical practice guidelines for patients with chronic conditions and multimorbidity: a systematic analysis. *BMJ Open* 2016; **6**(3): e010119.
- National Institute for Health and Care Excellence. *Multimorbidity: clinical assessment and management. NICE guideline [NG56]*. London: NICE, 2016. <https://www.nice.org.uk/guidance/ng56> [accessed 6 Apr 2021].
- Demain S, Gonçalves A-C, Areia C, *et al*. Living with, managing and minimising treatment burden in long term conditions: a systematic review of qualitative research. *PLoS One* 2015; **10**(5): e0125457.
- van Merode T, van de Ven K, van den Akker M. Patients with multimorbidity and their treatment burden in different daily life domains: a qualitative study in primary care in the Netherlands and Belgium. *J Comorb* 2018; **8**(1): 9–15.
- Gallacher K, May CR, Montori VM, Mair FS. Understanding patients' experiences of treatment burden in chronic heart failure using normalization process theory. *Ann Fam Med* 2011; **9**(3): 235–243.
- Shippee ND, Shah ND, May CR, *et al*. Cumulative complexity: a functional, patient-centered model of patient complexity can improve research and practice. *J Clin Epidemiol* 2012; **65**(10): 1041–1051.
- Gallacher KI, May CR, Langhorne P, Mair FS. A conceptual model of treatment burden and patient capacity in stroke. *BMC Fam Pract* 2018; **19**(1): 9.
- Boehmer KR, Shippee ND, Beebe TJ, Montori VM. Pursuing minimally disruptive medicine: disruption from illness and health care-related demands is correlated with patient capacity. *J Clin Epidemiol* 2016; **74**: 227–236.
- May C, Montori VM, Mair FS. We need minimally disruptive medicine. *BMJ* 2009; **339**: b2803.
- Eton DT, Yost KJ, Lai JS, *et al*. Development and validation of the Patient Experience with Treatment and Self-management (PETS): a patient-reported measure of treatment burden. *Qual Life Res* 2017; **26**(2): 489–503.
- Tran VT, Harrington M, Montori VM, *et al*. Adaptation and validation of the Treatment Burden Questionnaire (TBQ) in English using an internet platform. *BMC Med* 2014; **12**: 109.
- Boyd CM, Wolff JL, Giovannetti E, *et al*. Healthcare task difficulty among older adults with multimorbidity. *Med Care* 2014; **52** Suppl 3(03): S118–S125.
- Gibbons CJ, Kenning C, Coventry PA, *et al*. Development of a multimorbidity illness perceptions scale (MULTIPLEs). *PLoS One* 2013; **8**(12): e81852.
- Duncan P, Murphy M, Man M-S, *et al*. Development and validation of the Multimorbidity Treatment Burden Questionnaire (MTBQ). *BMJ Open* 2018; **8**(4): e019413.
- NHS Digital. *Quality and Outcomes Framework (QOF): business rules v41/v42*. Quality and Outcome Framework business rules. <https://digital.nhs.uk/data-and-information/data-collections-and-data-sets/data-collections/quality-and-outcomes-framework-qof> [accessed 7 Jan 2021].
- NHS Digital. *NHS UK Read codes*. <https://isd.digital.nhs.uk/trud3/user/guest/group/0/pack/9> [accessed 6 Apr 2021].
- Barnett K, Mercer SW, Norbury M, *et al*. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012; **380**(9836): 37–43.
- Morris NS, MacLean CD, Chew LD, Littenberg B. The Single Item Literacy Screener: evaluation of a brief instrument to identify limited reading ability. *BMC Fam Pract* 2006; **7**: 21.
- EuroQol Group. *EQ-5D-5L / About*. 2017. <https://euroqol.org/eq-5d-instruments/eq-5d-5l-about/> [accessed 6 Apr 2021].
- Farrar JT, Young JP Jr, LaMoreaux L, *et al*. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain* 2001; **94**(2): 149–158.
- Collins D. Pretesting survey instruments: an overview of cognitive methods. *Qual Life Res* 2003; **12**(3): 229–238.
- Ministry of Housing, Communities and Local Government. *English Housing Survey 2018 to 2019: headline report. Section 1: household tables* [data file]. 2020. <https://www.gov.uk/government/statistics/english-housing-survey-2018-to-2019-headline-report> [accessed 6 Apr 2021].
- Choung RS, Locke GR, Schleck CD, *et al*. A low response rate does not necessarily indicate non-response bias in gastroenterology survey research: a population-based study. *J Public Health* 2013; **21**: 87–95.
- Dalstra JAA, Kunst AE, Mackenbach JP. A comparative appraisal of the relationship of education, income and housing tenure with less than good health among the elderly in Europe. *Soc Sci Med* 2006; **62**(8): 2046–2060.
- Sørensen K, Van den Broucke S, Fullam J, *et al*. Health literacy and public health: a systematic review and integration of definitions and models. *BMC Public Health* 2012; **12**: 80.
- McIntyre NJ, Fluck R, McIntyre C, Taal M. Treatment needs and diagnosis awareness in primary care patients with chronic kidney disease. *Br J Gen Pract* 2012; DOI: <https://doi.org/10.3399/bjgp12X636047>.
- Majeed A, Car J, Sheikh A. Accuracy and completeness of electronic patient records in primary care. *Fam Pract* 2008; **25**(4): 213–214.
- Salisbury C, Man MS, Bower P, *et al*. Management of multimorbidity using a patient-centred care model: a pragmatic cluster-randomised trial of the 3D approach. *Lancet* 2018; **392**(10141): 41–50.
- Tran VT, Montori VM, Eton DT, *et al*. Development and description of measurement properties of an instrument to assess treatment burden among patients with multiple chronic conditions. *BMC Med* 2012; **10**: 68.
- Herzig L, Zeller A, Pasquier J, *et al*. Factors associated with patients' and GPs' assessment of the burden of treatment in multimorbid patients: a cross-sectional study in primary care. *BMC Fam Pract* 2019; **20**(1): 88.
- Rosbach M, Andersen JS. Patient-experienced burden of treatment in patients with multimorbidity: a systematic review of qualitative data. *PLoS One* 2017; **12**(6): e0179916.
- Friis K, Lasgaard M, Pedersen MH, *et al*. Health literacy, multimorbidity, and patient-perceived treatment burden in individuals with cardiovascular disease. A Danish population-based study. *Patient Educ Couns* 2019; **102**(10): 1932–1938.
- Boehmer KR, Kyriacou M, Behnken E, *et al*. Patient capacity for self-care in the medical record of patients with chronic conditions: a mixed-methods retrospective study. *BMC Fam Pract* 2018; **19**(1): 164.
- Ridgeway JL, Egginton JS, Tiedje K, *et al*. Factors that lessen the burden of treatment in complex patients with chronic conditions: a qualitative study. *Patient Prefer Adherence* 2014; **8**: 339–351.
- Eton DT, Ridgeway JL, Linzer M, *et al*. Healthcare provider relational quality is associated with better self-management and less treatment burden in people with multiple chronic conditions. *Patient Prefer Adherence* 2017; **11**: 1635–1646.
- Kyle J, Skleparis D, Mair FS, Gallacher KI. What helps and hinders the provision of healthcare that minimises treatment burden and maximises patient capacity? A qualitative study of stroke health professional perspectives. *BMJ Open* 2020; **10**(3): e034113.
- Collins B. *What is Covid-19 revealing about innovation in the NHS?* London: King's Fund, 2020. <https://www.kingsfund.org.uk/blog/2020/08/covid-19-innovation-nhs> [accessed 6 Apr 2021].
- Drinkwater C, Wildman J, Moffatt S. Social prescribing. *BMJ* 2019; **364**: l1285.