

Research

Angel YS Wong, Laurie Tomlinson, Jeremy P Brown, William Elson, Alex J Walker, Anna Schultze, Caroline E Morton, David Evans, Peter Inglesby, Brian MacKenna, Krishnan Bhaskaran, Christopher T Rentsch, Emma Powell, Elizabeth Williamson, Richard Croker, Seb Bacon, William Hulme, Chris Bates, Helen J Curtis, Amir Mehrkar, Jonathan Cockburn, Helen I McDonald, Rohini Mathur, Kevin Wing, Harriet Forbes, Rosalind M Eggo, Stephen JW Evans, Liam Smeeth, Ben Goldacre and Ian J Douglas (The OpenSAFELY Collaborative)

Association between oral anticoagulants and COVID-19-related outcomes:

a population-based cohort study

Abstract

Background

Early evidence has shown that anticoagulant reduces the risk of thrombotic events in those infected with COVID-19. However, evidence of the role of routinely prescribed oral anticoagulants (OACs) in COVID-19 outcomes is limited.

Aim

To investigate the association between OACs and COVID-19 outcomes in those with atrial fibrillation and a CHA₂DS₂-VASc score of 2.

Design and setting

On behalf of NHS England, a population-based cohort study was conducted.

Method

The study used primary care data and pseudonymously-linked SARS-CoV-2 antigen testing data, hospital admissions, and death records from England. Cox regression was used to estimate hazard ratios (HRs) for COVID-19 outcomes comparing people with current OAC use versus non-use, accounting for age, sex, comorbidities, other medications, deprivation, and general practice.

Results

Of 71 103 people with atrial fibrillation and a CHA₂DS₂-VASc score of 2, there were 52 832 current OAC users and 18 271 non-users. No difference in risk of being tested for SARS-CoV-2 was associated with current use (adjusted HR [aHR] 0.99, 95% confidence interval [CI] = 0.95 to 1.04) versus non-use. A lower risk of testing positive for SARS-CoV-2 (aHR 0.77, 95% CI = 0.63 to 0.95) and a marginally lower risk of COVID-19-related death (aHR, 0.74, 95% CI = 0.53 to 1.04) were associated with current use versus non-use.

Conclusion

Among those at low baseline stroke risk, people receiving OACs had a lower risk of testing positive for SARS-CoV-2 and severe COVID-19 outcomes than non-users; this might be explained by a causal effect of OACs in preventing severe COVID-19 outcomes or unmeasured confounding, including more cautious behaviours leading to reduced infection risk.

Keywords

COVID-19, warfarin, dabigatran, Factor Xa Inhibitors.

INTRODUCTION

Early studies have reported that heparin lowers the risk of pulmonary embolism and mortality during hospital admissions in patients with COVID-19.¹⁻⁴ However, the protective role of oral anticoagulants (OACs) in preventing severe COVID-19 outcomes is unclear. A randomised controlled trial has reported that therapeutic anticoagulation with rivaroxaban in patients admitted to hospital with COVID-19 did not improve all-cause mortality but increased the risk of major bleeding compared with prophylactic anticoagulation with enoxaparin or unfractionated heparin during hospital admissions.⁵ Notably, as this study excluded people with an indication for therapeutic anticoagulation, it remains uncertain whether there is an effect of routinely prescribed OACs for atrial fibrillation on COVID-19 outcomes. In

particular, anticoagulants are underutilised among people with atrial fibrillation in the UK.⁶ Understanding the effects of routinely prescribed OACs on COVID-19 may be of significant clinical importance that further informs OAC prescribing guidance in the context of the COVID-19 pandemic, given that people with atrial fibrillation are at higher risk of severe COVID-19 outcomes.⁷

Some observational studies have investigated the role of OACs in COVID-19 outcomes but the findings were conflicting.⁸⁻¹⁹ Importantly, most studied were of small sample size or in hospital settings only, leading to a lack of power and/or selection bias. In addition, people taking OACs are likely to have more comorbidities than those who are not, and so comparing COVID-19 outcomes for OAC use with non-use could be subject to confounding in these studies.

AYS Wong, PhD, assistant professor; **L Tomlinson**, PhD, associate professor; **JP Brown**, MSc, PhD student; **W Elson**, MBBS, general practitioner; **A Schultze**, PhD, research fellow; **K Bhaskaran**, PhD, professor of statistical epidemiology; **CT Rentsch**, PhD, assistant professor; **E Powell**, MSc, PhD student; **E Williamson**, PhD, biostatistician; **R Mathur**, PhD, assistant professor; **K Wing**, PhD, assistant professor; **IJ Douglas**, PhD, professor; **RM Eggo**, PhD, associate professor; **SJW Evans**, PhD, professor, Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London. **AJ Walker**, PhD, epidemiologist; **CE Morton**, MBChB, epidemiologist; **D Evans**, MPhil, software developer; **P Inglesby**, MPhil, consultant engineer; **B MacKenna**, MPharm, honorary clinical researcher; **R Croker**, MSc, honorary research fellow; **S Bacon**, BA, chief technical officer; **W Hulme**, PhD, statistician; **HJ Curtis**, DPhil, postdoctoral researcher; **A Mehrkar**, nMRCGP, honorary clinical researcher; **B Goldacre**, MRCPsych, director, The DataLab, Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford. **C Bates**, PhD, director of research and analytics; **J Cockburn**,

BSc, developer, TPP, Leeds. **HI McDonald**, PhD, clinical research fellow; **L Smeeth**, PhD, professor of clinical epidemiology, Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London and NIHR Health Protection Research Unit (HPRU) in Immunisation, London School of Hygiene and Tropical Medicine, London. **H Forbes**, PhD, epidemiologist, Bristol Medical School, University of Bristol, Bristol.

Address for correspondence

Angel YS Wong, Department of Non-Communicable Disease Epidemiology, Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK.

Email: Angel.Wong@lshtm.ac.uk

Submitted: 8 December 2021; **Editor's response:** 19 January 2022; **final acceptance:** 6 February 2022.

©The Authors

This is the full-length article (published online 20 Apr 2022) of an abridged version published in print. Cite this version as: **Br J Gen Pract 2022;** DOI: <https://doi.org/10.3399/BJGP.2021.0689>

How this fits in

Early studies reported that heparin lowers the risk of pulmonary embolism and mortality during hospitalisation among patients with COVID-19. However, the protective role of oral anticoagulants in severe COVID-19 outcomes is unclear. This study showed a lower risk of testing positive for SARS-CoV-2 and COVID-19-related deaths associated with current OAC use, versus non-use among people with atrial fibrillation and a low baseline risk of stroke.

In order to address these limitations, a population-based cohort study was conducted to investigate the association between routinely prescribed OACs and COVID-19 outcomes versus non-use, restricting the study population to people in England with atrial fibrillation who had a CHA₂DS₂-VASc score of 2. The CHA₂DS₂-VASc score is a validated measure to predict the risk of stroke among people with atrial fibrillation. The use of the CHA₂DS₂-VASc score is to determine the need for prescribed anticoagulants in people with non-valvular atrial fibrillation as a prophylactic therapy against stroke. According to the guidelines for the management of patients with atrial fibrillation,²⁰⁻²² people with a CHA₂DS₂-VASc score of ≥ 2 should be offered an anticoagulant. For those with a score of 2, there is possibly a degree of variation in OAC prescribing, offering a useful group in which OAC users are likely to be more comparable with non-users to minimise confounding. Also, a UK study has shown that patients with a CHA₂DS₂-VASc score of 2 were more likely to remain untreated with anticoagulants than patients with a score of ≥ 3 .⁶ A better understanding of the impact of OACs on COVID outcomes may alter the balance of benefits and risks of prescribing OACs for those around such a threshold.

METHOD

Study design

A population-based cohort study was conducted between 1 March 2020 and 28 September 2020.

Data source

Primary care records managed by the software provider TPP were linked to SARS-CoV-2 antigen testing data from the Second Generation Surveillance System, COVID-19-related hospital admissions from the Secondary Uses Service, and Office for National Statistics death data through OpenSAFELY, a data analytics platform created by the author team on behalf of

NHS England.²³ The dataset analysed within OpenSAFELY is based on 24 million people currently registered with primary care practices using TPP SystemOne software, representing 40% of the English population. It includes pseudonymised data such as coded diagnoses, prescribed medications, and physiological parameters. Details on information governance can be found in Supplementary Information S1.

Study populations

People with a diagnosis of atrial fibrillation on or before the study start date (1 March 2020) were identified. To reduce confounding, the cohort was limited to those with a CHA₂DS₂-VASc score of 2, as the indication for OAC therapy would typically be borderline among people with this score. Their CHA₂DS₂-VASc score was calculated based on their records relating to demographics and diagnoses, which contributed to the score before cohort entry.

People were excluded if they had missing data for sex, Index of Multiple Deprivation, <1 year of primary care records, or aged <18 years or >110 years and prescribed injectable anticoagulants 4 months before the study start date.

Exposure

Current OAC users were people who ever had an OAC prescription (that is warfarin, dabigatran, rivaroxaban, apixaban, edoxaban) in the 4 months before the study start, and the non-users were people with no record of an OAC prescription in the same period.

Outcomes and follow-up

The outcomes were:

- testing positive for SARS-CoV-2;
- a COVID-19-related hospital admission; and
- a COVID-19-related death (defined as the presence of ICD-10 codes U071 [confirmed COVID-19] and U072 [suspected COVID-19] on the death certificate).

Testing outcomes were obtained from the UK's Pillar 1 (NHS and Public Health England laboratories) and Pillar 2 (commercial partners) testing strategies and results from polymerase chain reaction (PCR) swab tests used to identify individuals who were symptomatic.²⁴ As pre-specified analyses, time to the first SARS-CoV-2 test was also included as a negative control outcome. It was anticipated that, within




Figure 1. Study diagram, illustrating the follow-up from 1 March 2020 to the latest at death, outcome occurrence, deregistration from TPP (TPP SystemOne software practice), or 28 September 2020 and identification of covariates before 1 March 2020. Date in brackets shows the range of the time window (days in unit). A&E = accident & emergency. EFU = end of follow-up. OAC = oral coagulation. TIA = transient ischaemic attack.

this population of people with atrial fibrillation, there were unlikely to be marked differences in the likelihood of being tested for SARS-CoV-2 infection in relation to drug treatment with OACs.

Follow-up began on the 1 March 2020 and ended at the latest at the outcome of interest in each analysis, deregistration from the TPP practice, death, or study end date (28 September 2020) (Figure 1).

Covariates

Clinical covariates were identified using the Read clinical classification system and included body mass index (BMI) to classify obesity and smoking status. Obesity was classified using BMI in kg/m²: no record of obesity or BMI <30.0; obese I (BMI 30.0–34.9 kg/m²); obese II (BMI 35.0–39.9 kg/m²) and obese III (≥ 40 kg/m²). Smoking status was grouped into current, former, and never smokers. Those with missing smoking status were categorised as never smokers. The following covariates were pre-specified using a directed acyclic graph (DAG) (Supplementary Figure S1), including age, sex, obesity, smoking status, hypertension, heart failure, myocardial infarction, peripheral arterial disease, stroke/transient ischaemic attack, venous thromboembolism, diabetes, flu

vaccination, current antiplatelet use, current oestrogen/oestrogen-like therapy use, and Index of Multiple Deprivation. Dementia was added to the regression during the peer-review process as it could be an important confounder. The authors of the present study identified these covariates, which are both associated with the exposure to OACs and severe COVID-19 outcomes.²³ Some covariates that are associated with the exposure and venous thromboembolism, possibly leading to severe COVID-19 outcomes, were also included.^{25,26} All code lists for identifying exposures, covariates, and outcomes are openly shared at <https://codelists.opensafely.org/>.

Statistical methods

Adjusted cumulative incidence/mortality curves are presented using the Royston-Parmar model. Hazard ratios (HRs) with 95% confidence intervals (CIs) were estimated using Cox regression with time since cohort entry as the underlying timescale. Competing risk was accounted for by modelling the cause-specific hazard (that is censoring non-COVID-19 other deaths for COVID-19 death analysis, and censoring any death for other outcomes analysis). Graphical methods and tests based on Schoenfeld residuals were used

Box 1. List of sensitivity analyses

| Sensitivity analysis | Justification |
|---|---|
| 1. In addition to the covariates identified by DAG, other covariates based on prior evidence of likely confounders such as chronic obstructive pulmonary disease, other respiratory diseases, cancer, immunosuppression, chronic kidney disease, general practice attendance rate in the year before cohort entry, and A&E attendance rate in the year before cohort entry were also included in the fully adjusted models (stratified by general practice) | To test the robustness of the covariate selection |
| 2. Additionally adjusted for ethnicity in DAG and fully adjusted models. In the fully adjusted models, additional covariates included chronic obstructive pulmonary disease, other respiratory diseases (not including asthma), cancer, immunosuppression, chronic kidney disease, general practice attendance rate in the year before cohort entry, and A&E attendance rate in the year before cohort entry | In the main analysis, ethnicity is not adjusted for because of a sizable proportion of individuals with missing ethnicity (~23%). Complete case analysis was undertaken to address missing data |
| 3. Repeated main analysis excluding people prescribed antiplatelets 4 months before study start date | To explore the impact of use of antiplatelets, which can reduce the risk of blood clots |
| 4. Limited the study cohort to people aged ≥ 55 years | To explore the impact of potential confounders at a young age, for example, pregnancy |
| 5. Stratified the cohort by care home residence for the outcome of testing positive for SARS-CoV-2 (<i>post hoc</i> analysis) | To explore the impact of health behaviour, as people living in care homes were less likely to have differences in behaviour |

A&E = accident & emergency. DAG = directed acyclic graph.

to explore violations of the proportional hazards assumption.

Unadjusted models, models adjusted for age (using restricted cubic splines) and sex, and a DAG-adjusted model were constructed.

Quantitative bias analysis

The authors considered that OAC users may lower their COVID-19 risk through more risk-averse health behaviour (for example wearing face masks, avoiding close proximity to others) than non-users. Given that health behaviour is not captured in databases, quantitative bias analyses (both the E-value and the Cornfield condition)²⁷ were conducted to assess the sensitivity of the results to this potential unmeasured confounder.

Sensitivity analyses

Box 1 shows the list of sensitivity analyses.

Data management was performed using Python 3.8 and SQL, with analysis carried out using Stata 16.1. All study analyses were pre-planned unless otherwise stated. All code for data management and analyses are archived at <https://github.com/opensafely/oral-anticoagulant-covid->

outcome, in addition to the pre-specified protocol (https://github.com/opensafely/anticoagulants-research/blob/master/protocol/Protocol_%20Anticoag%20OpenSAFELY_v3.docx). Deviations from pre-specified protocol, with reasons are provided in Supplementary Information S2 and Supplementary Table S1.

RESULTS

Of 71 103 people with atrial fibrillation and a CHA₂DS₂-VASc score of 2, there were 52 832 current OAC users and 18 271 non-users (Figure 2, Supplementary Table S2). The median age was 71 years (interquartile range [IQR] 66–75 years) among current users and 69 years (IQR 63–74 years) among non-users.

Current OAC users were more likely to be male, obese, former smokers, and have a medical history of heart failure, chronic obstructive pulmonary disease, and chronic kidney disease, but were less likely to have myocardial infarction, peripheral artery disease, venous thromboembolism, immunosuppression, and diabetes than non-users. Current users were less likely to have a prescription for oestrogen/oestrogen-like drugs, antiplatelets, non-steroidal anti-inflammatory drugs, and aspirin, but to have had more primary care consultations and a flu vaccination than non-users.

Supplementary Figure S2 presents time to COVID-19-related outcomes in adjusted cumulative incidence plots. A lower risk of testing positive for SARS-CoV-2 (480 events in 2 107 517 person-weeks; unadjusted HR 0.75 [95% CI = 0.62 to 0.91]; DAG-adjusted HR 0.77 [95% CI = 0.63 to 0.95]) was associated with current use, compared with non-use (Figure 3, Supplementary Table S3). Similarly, a lower risk of COVID-19-related hospital admission was observed with wide CIs (226 events in 2 110 854 person-weeks; unadjusted HR 0.89 [95% CI = 0.67 to 1.19]; DAG-adjusted HR 0.85 [95% CI: 0.62 to 1.15]), and COVID-19-related deaths (185 events in 2 113 796 person-weeks; unadjusted HR 0.81 [95% CI = 0.59 to 1.11]; DAG-adjusted HR 0.74 [95% CI = 0.53 to 1.04]), comparing current use with non-use.

For the negative control outcome, there was no difference in risk of being tested for SARS-CoV-2 comparing current use with non-use (11 190 events in 1 994 072 person-weeks; unadjusted HR 0.95 [95% CI = 0.91 to 0.99]; DAG-adjusted HR 0.99 [95% CI = 0.95 to 1.04]) (see Figure 3 & Supplementary Table S3).




Figure 2. Flow chart of inclusion of participants. IMD = Index of Multiple Deprivation. TPP = TPP SystemOne software practice.

Figure 3. Hazard ratios of the association between current use of oral anticoagulants and COVID-19-related outcomes versus non-use in people with atrial fibrillation who had a CHA₂DS₂-VASc score of 2. CI = confidence interval. DAG = directed acyclic graph.

Sensitivity analyses

In the fully adjusted model, a lower risk of COVID-19-related death comparing current OAC use with non-use (HR 0.66, 95% CI = 0.45 to 0.97) was observed. Results of all other sensitivity analyses were similar to those in the main analyses (Supplementary Tables S4–S6). Although there was no strong evidence of a different

association between current use of OACs and testing positive for SARS-CoV-2 according to care home residence (Supplementary Table S7), this comparison was underpowered.

Quantitative bias analysis


To potentially fully explain the observed associations, either non-anticoagulant use would need to be associated with at least a 1.29 increased risk of unmeasured risk-prone behaviour; or risk-prone behaviour would need to be associated with a 1.29 times increased risk of each COVID-19 outcome. Alternatively, both non-use and each COVID-19 outcome would need to be associated with at least a 1.05 times increased risk or unmeasured risk-prone behaviour (Supplementary Table S8).

DISCUSSION

Summary

A lower risk of testing positive for SARS-CoV-2 was associated with current OAC use versus non-use among people with atrial fibrillation and a low baseline risk of stroke. With small absolute numbers of COVID-19-related hospital admissions and COVID-19-related deaths, a marginally lower risk of these outcomes was associated with current OAC use compared with non-use. No difference in the risk of being tested for SARS-CoV-2 between current users and non-users was found, indicating that the lower risk of testing positive was unlikely to be because of the chance of being tested.

Consideration needs to be given to whether these associations are causal, or because of other differences between groups. There is no clear evidence that the current users were generally frailer in terms of their comorbidities than non-users. The inverse associations in OAC users were specific to COVID-19 outcomes, with no protective association seen against having a COVID-19 test, which would support a possible causal association. OAC users had a reduced risk of receiving a positive test and severe COVID-19-related outcomes, suggesting a lower risk of acquiring test-detected infection in this group. An experimental study suggested that direct factor Xa inhibitors may prevent SARS-CoV entry to human cells by preventing the spike protein cleavage into the S1 and S2 subunits²⁸ but the clinical evidence is limited. Additionally, the authors of the present study considered that anticoagulation might inhibit the PCR for SARS-CoV-2^{29,30} but the evidence of a specific inhibitory effect of OACs on PCRs for SARS-CoV-2 is lacking.



Funding:

The OpenSAFELY data science platform is funded by the Wellcome Trust. OpenSAFELY work was jointly funded by UKRI [COV0076;MR/V015737/1], NIHR and Asthma UK-BLF and the Longitudinal Health and Wellbeing strand of the National Core Studies programme. TPP provided technical expertise and infrastructure within their data centre pro bono in the context of a national emergency. Ben Goldacre's work on better use of data in healthcare more broadly is currently funded in part by: National Institute for Health Research (NIHR) Oxford Biomedical Research Centre, NIHR Applied Research Collaboration Oxford and Thames Valley, the Mohn-Westlake Foundation, NHS England, and the Health Foundation; all DataLab staff are supported by Ben Goldacre's grants on this work. Liam Smeeth reports grants from Wellcome, Medical Research Council (MRC), NIHR, UK Research and Innovation (UKRI), British Council, GlaxoSmithKline (GSK), British Heart Foundation (BHF), and Diabetes UK outside this work. Angel YS Wong holds a fellowship from BHF. Jeremy P Brown is funded by a studentship from GSK. Anna Schultze is employed by the London School of Hygiene and Tropical Medicine (LSHTM) on a fellowship sponsored by GSK. Krishnan Bhaskaran holds a Sir Henry Dale fellowship jointly funded by Wellcome and the Royal Society (107731/Z/15/Z). Helen I McDonald is funded by the NIHR Health Protection Research Unit in Immunisation, a partnership between Public Health England and LSHTM. Rohini Mathur holds a Sir Henry Wellcome fellowship. Elizabeth Williamson holds grants from MRC. Ian J Douglas holds grants from NIHR and GSK. Harriet Forbes holds a UKRI fellowship. The views expressed are those of the authors and not necessarily those of the NIHR, NHS England, Public Health England or the Department of Health and Social Care. Funders had no role in the study design, collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the article for publication.

Ethical Approval

This study was approved by the Health Research Authority (REC reference 20/LO/0651) and by the LSHTM Ethics Board (reference 21863).

Notably, these results do not necessarily mean OACs reduce the risk of infection; in many instances the outcome of a positive COVID-19 test reflects both infection and symptom severity leading to test seeking. It is possible that anticoagulant activity lessens symptoms/severity, leading to a lower rate of infection detection.

The authors of the present study have also considered the non-causal explanation that risk behaviours may differ between OAC users and non-users and this may have an impact on the probability of being infected with SARS-CoV-2 and severe COVID-19 outcomes. Although the authors cannot fully capture the behavioural differences between groups in the database, it was observed that OAC users were less likely to be current smokers, had less hazardous alcohol use, and were more likely to have had flu vaccination than non-users but the differences were small. A quantitative bias analyses was performed and found that an unmeasured confounder would need to be of moderate strength to potentially explain the observed associations. Although this is always possible, notably the very wide range of well measured confounders the authors did have information on had little impact on these findings, suggesting confounding may not be a substantial problem. Nonetheless, further studies that can account for behavioural differences between groups are required to confirm the findings as the authors of the present study cannot rule this out as a possible contributor to these findings.

Strengths and limitations

The greatest strength of this study was the power enabling the examination of the association between OACs and various COVID-19 outcomes as the dataset included medical records from 24 million individuals. A quantitative bias analyses was also conducted to explore the impact of unmeasured confounding on the observed results, complementing the interpretation. The breadth of data available in primary care makes it possible to account for a wide range of potential confounders.

Possible limitations are recognised. First, as in any observational study, residual confounding could not be eliminated. Further, there may be misclassification bias in measuring covariates, leading to incomplete adjustment for confounding. Although the authors of the present study attempted to reduce confounding by limiting the cohort to people who had a threshold CHA₂DS₂-VASc score for being prescribed anticoagulants, results may not

be generalisable to all patients with atrial fibrillation. In particular, females, people with stroke, transient ischaemic attack or venous thromboembolism may be under-represented in this study given that these alone would have led to a CHA₂DS₂-VASc score of 2, so any additional risk factors would mean therefore they would be excluded from the exposed group.

Second, it is not known whether patients took the medications as prescribed. However, non-adherence of OAC treatment would only bias the estimate towards null.

Third, it was not possible to capture anticoagulant use (for example low molecular weight heparin or unfractionated heparin) during admissions to hospital. However, this would tend to make the comparison groups more similar to each other during admissions to hospital, which would most likely lead to an underestimation of any effect of routine OAC use before hospital admission.

Fourth, there may be misclassification in ascertaining atrial fibrillation using diagnostic codes alone when deriving the study population. Some recorded atrial fibrillation might resolve at the study start and thus not require anticoagulants. However, this is considered to be less common³¹ and is unlikely to substantially bias these results. Fifth, some cases of people having COVID-19 in the early pandemic may have been missed because of limited testing capacity. Importantly, the effect estimate in this analysis would not be biased assuming non-differential misclassification bias of outcome.

Comparison with existing literature

Although the effects of COVID-19 may predispose patients to thromboembolic disease through severe illness, hypoxia, or severe inflammatory response,³² anticoagulation may have a role in preventing thrombotic events in patients with COVID-19. Recent studies investigating the potential effects of early initiation of anticoagulation resulted in conflicting findings.^{3,8-15,33-37}

Seven studies^{8,9,13,14,16-18} focusing on prehospital use of anticoagulants and one study³⁴ focusing on therapeutic use of anticoagulants found no difference in risk of mortality, mechanical ventilation, or acute respiratory distress syndrome. Notably, some were of small sample size,^{8,16-18} with unclear exposure definition,⁹ classifying patients who initiated therapeutic anticoagulation on day 3 or later after intensive care unit admission in the control group³⁴ or used composite outcomes with varying clinical importance,^{13,14} limiting the

Data

All data were linked, stored, and analysed securely within the OpenSAFELY platform. Detailed pseudonymised patient data are potentially re-identifiable and therefore not shared. The authors rapidly delivered the OpenSAFELY data analysis platform without previous funding to deliver timely analyses of urgent research questions in the context of the global COVID-19 health emergency: now that the platform is established, the authors are developing a formal process for external users to request access in collaboration with NHS England. Details of this process will be published in the near future on the OpenSAFELY website.

Provenance

Freely submitted; externally peer reviewed.

Competing interests

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare the following: Ben Goldacre has received research funding from Health Data Research UK (HDRUK), the Laura and John Arnold Foundation, the Wellcome Trust, the NIHR Oxford Biomedical Research Centre, the NHS National Institute for Health Research School of Primary Care Research, the Mohn-Westlake Foundation, the Good Thinking Foundation, the Health Foundation, and the World Health Organization; he also receives personal income from speaking and writing for lay audiences on the misuse of science. Ian J Douglas has received unrestricted research grants and holds shares in GlaxoSmithKline. All other authors have declared no competing interests.

Contributors

Ben Goldacre and Ian J Douglas are joint principal investigators.

Acknowledgements

The authors are very grateful for all the support received from the TPP Technical Operations team throughout this work; for generous assistance from the information governance and database teams at NHS England/NHSX.

Open access

This article is Open Access: CC BY 4.0 licence (<http://creativecommons.org/licenses/by/4.0/>).

Discuss this article

Contribute and read comments about this article: bjgp.org/letters

interpretation of the results. Two cohort studies showed a higher risk of admission to an intensive care unit, intubation, or death associated with anticoagulants in patients with COVID-19 disease versus non-use without restricting to a study population with a specific OAC indication, but the findings were possibly the result of confounding by indication.^{12,36} A cohort study with propensity score matching reported a lower risk of all-cause mortality associated with OACs in people with COVID-19 disease compared with non-use, supporting the results in the present study.¹⁹

Implications for research and practice

Notably, this study was undertaken during the early pandemic before vaccines and COVID-19 treatments were available to prevent or treat severe COVID-19 disease. Future work is needed to confirm these

findings of an inverse association between OACs and severe COVID-19-related outcomes in people with atrial fibrillation at the threshold for OAC treatment, and to establish causality; either randomised trials or observational studies with detailed data on risk factors for COVID-19 infection. If confirmed to be a causal effect, this could be of significant clinical importance, particularly as the older age and comorbidities in this group are independent risk factors for severe COVID-19 outcomes. Choice of whether to prescribe routine anticoagulant therapy represents a complex balance of expected risks, benefits, and patient preference; the authors do not recommend changes to ongoing anticoagulant therapy based on these results.

REFERENCES

- Fauvel C, Weizman O, Trimaille A, *et al*. Pulmonary embolism in COVID-19 patients: a French multicentre cohort study. *Eur Heart J* 2020; **41(32)**: 3058–3068.
- Paranjpe I, Fuster V, Lala A, *et al*. Association of treatment dose anticoagulation with in-hospital survival among hospitalized patients with COVID-19. *J Am Coll Cardiol* 2020; **76(1)**: 122–124.
- Nadkarni GN, Lala A, Bagiella E, *et al*. Anticoagulation, mortality, bleeding and pathology among patients hospitalized with COVID-19: a single health system study. *J Am Coll Cardiol* 2020; **76(16)**: 1815–1826.
- Tang N, Bai H, Chen X, *et al*. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost* 2020; **18(5)**: 1094–1099.
- Lopes RD, de Barros E Silva PGM, Furtado RHM, *et al*. Therapeutic versus prophylactic anticoagulation for patients admitted to hospital with COVID-19 and elevated D-dimer concentration (ACTION): an open-label, multicentre, randomised, controlled trial. *Lancet* 2021; **397(10291)**: 2253–2263.
- Lacoin L, Lumley M, Ridha E, *et al*. Evolving landscape of stroke prevention in atrial fibrillation within the UK between 2012 and 2016: a cross-sectional analysis study using CPRD. *BMJ Open* 2017; **7(9)**: e015363.
- Inciardi RM, Adamo M, Lupi L, Metra M. Atrial fibrillation in the COVID-19 era: simple bystander or marker of increased risk? *Eur Heart J* 2020; **41(32)**: 3094.
- Russo V, Di Maio M, Attena E, *et al*. Clinical impact of pre-admission antithrombotic therapy in hospitalized patients with COVID-19: a multicenter observational study. *Pharmacol Res* 2020; **159**: 104965.
- Tremblay D, van Gerwen M, Alsen M, *et al*. Impact of anticoagulation prior to COVID-19 infection: a propensity score-matched cohort study. *Blood* 2020; **136(1)**: 144–147.
- Chocron R, Galand V, Cellier J, *et al*. Anticoagulation before hospitalization is a potential protective factor for COVID-19: insight from a French multicenter cohort study. *J Am Heart Assoc* 2021; **10(8)**: e018624.
- Harrison RF, Forte K, Buscher MG Jr, *et al*. The association of preinfection daily oral anticoagulation use and all-cause in hospital mortality from novel coronavirus 2019 at 21 days: a retrospective cohort study. *Crit Care Explor* 2021; **3(1)**: e0324.
- Rivera-Caravaca JM, Núñez-Gil IJ, Vivas D, *et al*. Clinical profile and prognosis in patients on oral anticoagulation before admission for COVID-19. *Eur J Clin Invest* 2021; **51(1)**: e13436.
- Ho G, Dusendang JR, Schmittiel J, *et al*. Association of chronic anticoagulant and antiplatelet use on disease severity in SARS-COV-2 infected patients. *J Thromb Thrombolysis* 2021; **52(2)**: 476–481.
- Flam B, Wintzell V, Ludvigsson JF, *et al*. Direct oral anticoagulant use and risk of severe COVID-19. *J Intern Med* 2021; **289(3)**: 411–419.
- Fröhlich GM, Jeschke E, Eichler U, *et al*. Impact of oral anticoagulation on clinical outcomes of COVID-19: a nationwide cohort study of hospitalized patients in Germany. *Clin Res Cardiol* 2021; **110(7)**: 1041–1050.
- Rossi R, Coppi F, Talarico M, Boriani G. Protective role of chronic treatment with direct oral anticoagulants in elderly patients affected by interstitial pneumonia in COVID-19 era. *Eur J Intern Med* 2020; **77**: 158–160.
- Spiegelberg JP, van Gelder MMHJ, Maas ML, *et al*. Prior use of therapeutic anticoagulation does not protect against COVID-19 related clinical outcomes in hospitalized patients: a propensity score-matched cohort study. *Br J Clin Pharmacol* 2021; **87(12)**: 4839–4847.
- Schiavone M, Gasperetti A, Mancone M, *et al*. Oral anticoagulation and clinical outcomes in COVID-19: An Italian multicenter experience. *Int J Cardiol* 2021; **323**: 276–280.
- Denas G, Gennaro N, Ferroni E, *et al*. Reduction in all-cause mortality in COVID-19 patients on chronic oral anticoagulation: A population-based propensity score matched study. *Int J Cardiol* 2021; **329**: 266–269.
- January CT, Wann LS, Calkins H, *et al*. 2019 AHA/ACC/HRS Focused Update of the 2014 AHA/ACC/HRS Guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society in Collaboration with the Society of Thoracic Surgeons. *Circulation* 2019; **140(2)**: e125–e151.
- National Institute for Health and Care Excellence. *Recommendations. Atrial fibrillation: diagnosis and management*. [NG196]. 2021. <https://www.nice.org.uk/guidance/ng196/resources/atrial-fibrillation-diagnosis-and-management-pdf-66142085507269> (accessed 3 March 2022).
- National Institute for Health and Care Excellence. *In those patients with atrial fibrillation with a record of a CHA2DS2-VASc score of 2 or more, the percentage of patients who are currently treated with anticoagulation drug therapy. NICE Quality and Outcomes Framework indicator. Standards and Indicators*. 2014. <https://www.nice.org.uk/standards-and-indicators/qofindicators/in-those-patients-with-atrial-fibrillation-whose-latest-record-of-a-cha2ds2-vasc-score-is-2-or-above-the-percentage-of-patients-who-are-currently-treated-with-anti-coagulation-drug-therapy> (accessed 3 March 2022).
- Williamson EJ, Walker AJ, Bhaskaran K, *et al*. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020; **584**: 430–436.
- UK Government. *Coronavirus (COVID-19): scaling up our testing programmes*. 2021. <https://www.gov.uk/government/publications/coronavirus-covid-19-scaling-up-testing-programmes/coronavirus-covid-19-scaling-up-our-testing-programmes> (accessed 3 March 2022).
- Vinogradova Y, Coupland C, Hippisley-Cox J. Use of hormone replacement therapy and risk of venous thromboembolism: nested case-control studies using the QRResearch and CPRD databases. *BMJ* 2019; **364**: k4810.
- Steinhubl SR, Eikelboom JW, Hylek EM, *et al*. Antiplatelet therapy in prevention of cardio- and venous thromboembolic events. *J Thromb Thrombolysis* 2014; **37(3)**: 362–371.
- Ding P, VanderWeele TJ. Sensitivity analysis without assumptions. *Epidemiology* 2016; **27(3)**: 368–377.
- Du L, Kao RY, Zhou Y, *et al*. Cleavage of spike protein of SARS coronavirus by protease factor Xa is associated with viral infectivity. *Biochem Biophys Res Commun* 2007; **359(1)**: 174–179.
- Miyachi H, Masukawa A, Ohshima T, *et al*. Monitoring of inhibitors of enzymatic amplification in polymerase chain reaction and evaluation of efficacy of RNA extraction for the detection of hepatitis C virus using the internal control. *Clin Chem Lab Med* 1998; **36(8)**: 571–575.
- Kirstein LM, Mellors JW, Rinaldo CR Jr, *et al*. Effects of anticoagulant, processing delay, and assay method (branched DNA versus reverse transcriptase PCR) on measurement of human immunodeficiency virus type 1 RNA levels in plasma. *J Clin Microbiol* 1999; **37(8)**: 2428–2433.
- Ruigómez A, Johansson S, Wallander M-A, García Rodríguez LA. Predictors and prognosis of paroxysmal atrial fibrillation in general practice in the UK. *BMC Cardiovasc Disord* 2005; **5**: 20.
- Bikdeli B, Madhavan MV, Jimenez D, *et al*. COVID-19 and thrombotic or thromboembolic disease: implications for prevention, antithrombotic therapy, and follow-up: JACC state-of-the-art review. *J Am Coll Cardiol* 2020; **75(23)**: 2950–2973.
- Hanif A, Khan S, Mantri N, *et al*. Thrombotic complications and anticoagulation in COVID-19 pneumonia: a New York City hospital experience. *Ann Hematol* 2020; **99(10)**: 2323–2328.
- Al-Samkari H, Gupta S, Leaf RK, *et al*. Thrombosis, bleeding, and the observational effect of early therapeutic anticoagulation on survival in critically ill patients with COVID-19. *Ann Intern Med* 2021; **174(5)**: 622–632.
- Hsu A, Liu Y, Zayac AS, Olszewski AJ, *et al*. Intensity of anticoagulation and survival in patients hospitalized with COVID-19 pneumonia. *Thromb Res* 2020; **196**: 375–378.
- Zhou J, Lee S, Guo CL, *et al*. Anticoagulant or antiplatelet use and severe COVID-19 disease: A propensity score-matched territory-wide study. *Pharmacol Res* 2021; **165**: 105473.
- Rentsch CT, Beckman JA, Tomlinson L, *et al*. Early initiation of prophylactic anticoagulation for prevention of coronavirus disease 2019 mortality in patients admitted to hospital in the United States: cohort study. *BMJ* 2021; **372**: n311.