

Accepted Manuscript

British Journal of General Practice

Recommendations for the recognition, diagnosis, and management of long covid: A Delphi study

Nurek, Martine; Rayner, Clare; Freyer, Anette; Taylor, Sharon; Järte, Linn; MacDermott, Nathalie; Delaney, Brendan

DOI: <https://doi.org/10.3399/BJGP.2021.0265>

To access the most recent version of this article, please click the DOI URL in the line above.

Received 20 April 2021

Revised 28 May 2021

Accepted 07 June 2021

© 2021 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by British Journal of General Practice. For editorial process and policies, see: <https://bjgp.org/authors/bjgp-editorial-process-and-policies>

When citing this article please include the DOI provided above.

Author Accepted Manuscript

This is an 'author accepted manuscript': a manuscript that has been accepted for publication in British Journal of General Practice, but which has not yet undergone subediting, typesetting, or correction. Errors discovered and corrected during this process may materially alter the content of this manuscript, and the latest published version (the Version of Record) should be used in preference to any preceding versions

TITLE

Recommendations for the recognition, diagnosis, and management of long covid: A Delphi study

AUTHORS

Martine Nurek¹ (BSc, MSc, PhD), Clare Rayner² (MFOM, MRCP, MB. ChB.), Anette Freyer³ (MBBCh, PhD, FRCP), Sharon Taylor⁴ (MBBS, BSc, MRCP, FRCPsych, CASLAT, Dip Family Therapy, ILT), Linn Järte⁵ (MBBS, iBSc (Hons), DTM&H, DMCC), Nathalie MacDermott⁶ (BSc, MBBCh, MRCPCH), and Brendan C Delaney^{7,*} (BM BCh, MD, FRCP, FRCGP, FFCI), on behalf of the Delphi panellists (listed).

1. Postdoctoral Research Associate (Imperial College London, Department of Surgery and Cancer, 5th Floor Medical School Building, St Mary's Campus, Norfolk Place, London W2 1PG, UK);
 2. Occupational Physician (Independent);
 3. Consultant in Acute and Obstetric Medicine (Nottingham University Hospitals NHS Trust, Department of Acute Medicine, Queen's Medical Centre, Nottingham NG7 2UH, UK);
 4. Consultant Child and Adolescent Psychiatrist (Central and North West London NHS Foundation Trust/Imperial College School of Medicine, 7a Woodfield Road, London W9 3XZ, UK) and Honorary Senior Clinical Lecturer (Imperial College London);
 5. CT1 Anaesthetist (Swansea Bay University Health Board, Anaesthetics Department, Morriston Hospital, Heol Maes Eglwys, Morriston, Cwmrhydyceirw, Swansea SA6 6NL);
 6. NIHR Academic Clinical Lecturer (King's College London, Floor 5 Becket House, St Thomas' Hospital, Westminster Bridge Road, London SE1 7EH, UK);
 7. Chair in Medical Informatics and Decision Making (Imperial College London, Department of Surgery and Cancer, Medical School Room 506, St Mary's Campus, Norfolk Place, London W2 1PG, UK) and Principal in General Practice (Albion Street Group Practice, London SE16, UK).
- * Correspondence concerning this article should be addressed to Prof Brendan Delaney, Imperial College London, Department of Surgery and Cancer, Medical School Room 506, St Mary's Campus, Norfolk Place, London W2 1PG, UK (email: brendan.delaney@imperial.ac.uk; tel: +44 (0)20 7594 3427)

ABSTRACT

Background: In the absence of research into therapies and care pathways for long covid, guidance based on “emerging experience” is needed.

Aim: To provide a rapid expert guide for GPs and long covid clinical services.

Design: Delphi study.

Setting: Primary and secondary care doctors.

Method: The authors generated recommendations pertaining to the investigation and management of long covid. These were distributed online to a panel of UK doctors (any specialty) with an interest in, lived experience of, and/or experience treating long covid. Over two rounds of Delphi testing, panellists indicated their agreement with each recommendation (5-point Likert scale) and gave comments. Recommendations eliciting a response of “strongly agree”, “agree”, or “neither agree nor disagree” from $\geq 90\%$ of respondents were taken as showing consensus.

Results: Thirty-three clinicians representing 14 specialties reached consensus on 35 recommendations. Chiefly: GPs should consider long covid in the presence of a wide range of presenting features (not limited to fatigue and breathlessness) and exclude differential diagnoses where appropriate. Detailed history and examination with baseline investigations should be conducted in primary care. Indications for further investigation and specific therapies (for myocarditis, PoTS, mast cell disorder) include hypoxia/desaturation, chest pain, palpitations, and histamine-related symptoms. Rehabilitation should be individualised, with careful activity pacing (to avoid relapse) and multidisciplinary support.

Conclusions: Long covid clinics should operate as part of an integrated care system, with GPs playing a key role in the multidisciplinary team. Holistic care pathways, investigation of specific complications, management of potential symptom clusters, and tailored rehabilitation are needed.

Keywords: Post Covid-19 Condition, long covid, long-hauler, Post-Acute Sequelae of Covid-19 (PASC), general practice, guidelines

HOW THIS FITS IN

- There is an urgent need to devise clinical pathways and guidance for long covid (thought to affect 10% of those diagnosed with covid-19).
- In the absence of conclusive research to inform clinical practice, “expert physician-patients” (i.e., doctors with long covid and those involved in nascent clinics) are a source of professional expertise.
- Using robust consensus methodology, we derived 35 clear and practical recommendations to assist in the organisation of clinics, and the diagnosis and management of patients with long covid.
- Medically-led multidisciplinary clinics are required as serious cardiovascular, neurocognitive, respiratory and immune sequelae such as can present with only non-specific symptoms.

Accepted Manuscript – BJGP – BJGP:2021.0265

INTRODUCTION

“Post Covid-19 Condition” is an umbrella term for a complex, multi-system illness that follows on from an acute covid-19 infection, irrespective of severity, either immediately or sometime after apparent recovery. The terms “long covid” (UK) and “long-haulers” (US) were adopted by patient groups as people came together to compare their experiences.¹ Much of the publicity and campaigning has been around the term “long covid”, however WHO and SNOMED International have adopted the term “Post Covid-19 Condition” for classification. We will use the term “long covid” in this article but accept that medical terminology may continue with “Post Covid-19 Condition” or “Post-Acute Sequelae of SARS-CoV-2” in the USA.

It is estimated that over one million people in the UK (population: 66 million) are currently living with long covid (prolonged symptoms at 4+ weeks).² No such data are available for other countries, but numbers are likely to be proportionate, placing huge and immediate pressure on health services worldwide. Associated conditions have yet to be fully delineated, but examples are included in Box 1.³ The illness commonly has an unpredictable, relapsing-remitting pattern with significant associated conditions often appearing weeks to months into the disease course. Therefore, a high index of suspicion and a low threshold for referral to secondary care specialists or doctor-led long covid clinics with diagnostic capabilities, as per local availability, is advised.

In December 2020, the UK National Institute for Health and Care Excellence (NICE) produced a “rapid guideline”,⁴ alongside the launch of both community-based and specialist clinics for long covid. Discussion amongst an online community of doctors with long covid (members of the “UK doctors #longcovid” group on Facebook) quickly identified that there was a practical gap between best consensus practice in clinical care and the cautious, evidenced-based approach adopted by NICE (itself limited by the paucity of evidence concerning investigation and treatment of this new condition).⁵ In such particular circumstances, we must turn to clinical experience during 2020 as a guide for managing long covid. Clinicians with lived experience of the condition have

been instrumental in helping to define the problem, and “expert clinician-patient” self-help groups remain a vital resource in providing advice to the wider clinical community. To capture, organise, and disseminate this knowledge, we used robust consensus-based methods to derive recommendations for best practice in the recognition, investigation, and management of long covid. These recommendations are intended to guide generalist doctors providing medical supervision of a community-based long covid clinic with access to specialist referrals if required.

METHOD

Panel selection

A Delphi panel should consist of experts in the area of interest. In long covid, there is developing expertise through experience. Research studies have only recently begun to receive funding, but doctors with long covid have over the past year carried out a dynamic discussion on social media, highlighting new case reports, important studies, and potential clinical advances. Many of these doctors are involved in research and/or publishing articles. At this point in our knowledge, a pool of doctors with lived experience of long covid, combined with UK-based clinicians (of varying specialities) involved in service provision for long covid, form a suitable expert group.⁶

A call for panellists was placed on the “UK doctors #longcovid” support group, hosted by the social media platform Facebook. This is a closed group, exclusive to UK doctors with an interest in long covid, many of whom are seeing patients with long covid and/or have lived experience of the condition themselves (currently, approximately 1,100 members). Doctors interested in joining the panel were asked to provide their e-mail address by direct message. To ensure representation of all relevant specialties, specialist experts known to the authors were approached directly (via e-mail) and invited to join the panel.

The Delphi process

The present Delphi study comprised three stages, described below.

Initial identification of items

Following the publication of the NICE rapid guideline⁴ and a “living review” by the UK National Institute of Health Research (NIHR),⁷ the named authors generated a list of potential recommendations (“items”) to cover common clinical problems that were not fully addressed in the review. These were refined in a series of Zoom meetings to create 33 statements. Panellists had all seen the NIHR review and NICE guidance, had been following the research literature, and – more importantly in this case – following evolving clinical experience.

Round 1

Panellists were e-mailed a link to an online questionnaire (hosted by Qualtrics) with a request to respond within four days. Upon clicking the link, they gave their contact information (name, e-mail address), qualifications (specialty, year of qualification) and consent to participate in the Delphi study (we made clear that the results would be published in the form of a scientific article authored by the panel as a whole, and that each panellist would have the opportunity to review the article prior to submission). Thereafter, they read a brief introduction to the study processes and a definition of long covid (with examples).³ They then indicated the degree to which they agreed with each of the 33 initial items (presented sequentially) on a 5-point Likert scale (1=“strongly disagree”, 2=“disagree”, 3=“neither agree nor disagree”, 4=“agree”, 5=“strongly agree”). They were also encouraged to comment on each item in a free-text box, particularly if they disagreed with the item. Item scoring and comments were downloaded from Qualtrics (MN), anonymised and summarised (MN), and circulated to the named authors (CR, AF, ST, LJ, NM, BD) to facilitate review and revision of the items.

Items eliciting a response of “strongly agree”, “agree”, or “neither agree nor disagree” from $\geq 90\%$ of panellists were taken as showing consensus. Items with consensus were subject to minor amendment for sense only. If more substantial amendments were needed, the item was reworked on the basis of panel comments and retested in Round 2, alongside items that did not obtain consensus initially.

Round 2

Respondents that took part in Round 1 were e-mailed a link to a second Qualtrics questionnaire (Round 2), with a request to respond within five days. As in Round 1, they indicated their level of agreement with sequentially-presented items (either amended from Round 1 or newly-added to Round 2) using the 5-point Likert scale and gave comments. Items that achieved consensus were included in the final list of recommendations (with minor amendments as before to reflect participants’ feedback). Items that failed to achieve consensus were not included in the final list.

RESULTS

Panel characteristics

Thirty-seven doctors responded to the Facebook call and were e-mailed a link to Round 1; of these, 29 (78%) completed it. A further eight clinicians involved in service provision for long covid (known to the research team) were e-mailed a link to Round 1; of these, four (50%) completed it and four (50%) were too busy with acute covid work. All of the doctors that completed Round 1 also completed Round 2 (i.e., no doctors lost to follow up).

The panel therefore comprised 33 UK-based clinicians, representing a wide range of specialties (Fig. 1). The median number of years since qualification was 21 (range 0-41). Twenty-nine (88%) were recruited via social media and four (12%) via direct e-mail.

Twenty-nine (88%) had lived experience of long covid and five (15%) were clinicians developing services for long covid.

Round 1

Data collection took place over a nine-day period (24 Jan-1 Feb 2021). All 33 items obtained consensus. Of these, 18 required no/minor changes and were incorporated into the final list of recommendations (Supplementary Table 1: Round 1, blue text). The remaining 15 required more substantial work: 13 were amended to reflect panellists' feedback (Supplementary Table 1: Round 1, green text) and two were excluded (Supplementary Table 1: Round 1, red text). One of the excluded items was deemed self-evident (item 38), the other was strongly opposed by a relevant (respiratory) specialist (item 39).

Round 2

Round 2 data were collected over a six-day period (11 Feb-16 Feb 2021). Of the 19 items presented, 13 were amended from Round 1 and six were newly-added (usually to disentangle diagnosis and treatment when a Round 1 statement had addressed both). Eighteen items achieved consensus; of these, 17 were added to the final list of recommendations (Supplementary Table 1: Round 2, blue text) and one was excluded (Supplementary Table 1: Round 2, item 36): it was superfluous to another item that had obtained better agreement (agreement for item 36=97% vs. agreement for item 20=100%). One item did not obtain consensus (item 37, agreement=64%) and we did not feel that there would be a route to achieving this; therefore, this item was also excluded. Supplementary Figure 1 displays the results of the Delphi process graphically.

Final list of recommendations

The final list comprises 35 recommendations: six pertaining to clinic organisation (Box 2), 13 to diagnosis of the underlying disorder (Box 3), and 16 to management (Boxes 4 and 5). For a full and printable list of the recommendations, see Supplementary Box 1.

Recommendations pertaining to clinic organisation (Box 2)

Covid-19 is a new condition with increasing evidence of serious long-term sequelae, including cardiac, respiratory and renal disease, new-onset diabetes and excess deaths reported.⁸ It cannot therefore be assumed that patients are suffering from a self-limiting post-viral fatigue and that rehabilitation is sufficient. However, input from expert physiotherapists and occupational therapists who are familiar with the condition is an important aspect of caring for long covid patients.^{9,10} Patients require a holistic clinical approach that prioritises investigation of potential physical pathology.^{3,11} The lead clinician should be a doctor, “well versed in multi-system disorders”, working across disciplines, and able to refer to specialists. A respondent who runs a long covid clinic noted the importance of “easy access to multi-speciality input without multiple onward referrals (e.g., via multi-speciality post covid MDT)” and another that “isolated consultant clinics (without full MDT) will not work.” Long covid is not a primary mental health problem, but mental health specialists such as neuropsychiatrists can offer a supporting role to the multidisciplinary team. Psychological aspects of disease should be managed as part of the recovery process, but not seen as the primary treatment focus.^{3,4} Panellists were clear that no discrimination should exist in the treatment of patients with pre-existing mental health difficulties with regard to equal access to care for their long covid, and appropriate investigations for organ damage.¹¹ Regarding children, a Consultant Paediatrician should lead the service.³ NICE guidelines recommend considering referral from four weeks for specialist advice for children with ongoing symptomatic covid-19 or long covid.⁴

Recommendations pertaining to diagnosis of underlying disorder (Box 3)

At present there is a considerable risk to patient safety if appropriate investigation of common symptoms of long covid (such as chest pain, breathlessness, palpitations, abdominal pain, fatigue) which have wide differential diagnoses is not undertaken. Serious conditions, related to SARS-CoV-2 infection or not, must be adequately excluded¹² and investigations should be appropriately guided by the history. Long covid-specific examination (e.g., The NASA Lean Test for Postural Tachycardia Syndrome (PoTS)) or tests such as ECG are best conducted in person, and chest x-ray (CXR) may be appropriate. CXR may exclude relevant pathology such as Tuberculosis but is less relevant in investigating cardiac, pulmonary vascular, or autonomic causes for breathlessness where Computed Tomography or Ventilation/Perfusion (V/Q) scans are more likely to be indicated.¹³ In keeping with NICE guidance on asthma management,¹⁴ the panel agreed that spirometry with beta-agonist reversibility could be used to diagnose airway hyper-reactivity. Studies on venous thromboembolic disease are limited to patients with acute covid-19,¹⁵ severe disease,^{16,17} or based on expert opinion, but are an important diagnosis to consider.¹⁸ Oxygen desaturation on exertion occurs in both acute and long covid and should form part of baseline assessment. The only thresholds for defining levels of concern for hypoxaemia and desaturation with exercise pertain to acute covid-19^{19,20} and no agreed thresholds are available in long covid. Doctors working in existing clinics indicated that assessments like 1-minute sit-to-stand tests²¹ and 6-minute walk tests²² do or should form part of the assessment in community or specialist clinics. The exertional test chosen should take account of any pre-covid-19 limitations and should include heart rate as this may help to assess autonomic function. Referral for more detailed assessment is required in the following scenarios: desaturation with or without overt/reported dyspnoea; nocturnal desaturation; extreme fatigue; behavioural change in those who struggle with verbal communication; patient reports significant post exercise malaise after such testing (lasting beyond the next day); severe tachycardia; postural blood pressure drop.

There is increasing evidence of cardiovascular complications with covid-19.²³⁻³⁰ Patients with long covid (of all ages) have been diagnosed with arrhythmias, autonomic dysfunction, myocarditis, pericarditis, and microvascular ischaemia.³¹ The latter three

may only be seen on Cardiac Magnetic Resonance (cMRI) scans (gadolinium-enhanced; stress). Echocardiography has a low diagnostic yield for myocarditis in long covid,^{31,32} but diagnosis is important as people experience significant improvement in daily function through specific treatment. Pulmonary embolism appears to be rare more than six weeks after the acute illness and there are feasibility concerns about a potential surge of investigations for long covid. Usual risk scoring calculators were not valid in this context³³ and research is needed. Autonomic dysfunction, especially manifesting as PoTS, occurs commonly post covid-19.³⁴ There is a need to consider a differential for tachycardia and palpitations which, in long covid, includes pulmonary embolus, cardiac and respiratory causes. It was noted that autonomic dysfunction should also be suspected in patients with light-headedness, chest pain, and nausea, and the association of autonomic dysfunction with mast cell disorders considered.

As there are no accepted UK criteria for the diagnosis of “Mast Cell Activation Syndrome” and it remains an area of controversy, we used the term “mast cell disorder” to describe patients who present with a range of features listed in recommendation 17.³⁵ The list is not exhaustive and other serious disorders need to be excluded. This area is an important target of mechanistic and potential therapeutic studies in long covid. Neurocognitive testing is a particularly scarce resource⁴ and neurology review and brain MRI may be more helpful early in the illness. The benefits of testing would support the need for rehabilitation from occupational therapy or a neuropsychologist.^{4,36} NICE guidelines on long covid advise considering neuro-psychometric testing after six months if no improvement/worsening of cognitive function, as many will resolve.³⁷ At present, evidence related to joint swelling and arthralgia consists of only case reports, but clinicians should include long covid in a differential diagnosis of arthritis once other known autoimmune causes have been excluded.³⁸

Recommendations pertaining to management (Boxes 4 and 5)

The experience of many patients is of post-exertional symptom relapse. Physical or cognitive workload beyond the patient’s “energy envelope” may cause an exacerbation

of symptoms including fatigue, fever, myalgia, and breathlessness.³⁹ Exacerbations may manifest immediately or after a delay of 24-48 hours and may last days or months. As the threshold for this effect varies not only by patient but over time, pacing needs to be flexible and careful. Doctors play a key role in supporting patients through the complexity of specialist investigations and differential diagnoses, and considering symptomatic treatments. In addition, occupational health service referrals and medical reports supporting the return-to-work process are needed. Employers should discuss with their employee suitable adjustments to aid a return to work, and both parties should be provided with written advice such as the leaflet “covid-19 return to work guide for recovering workers” by the Society of Occupational Medicine.⁴⁰ The relapsing-remitting nature of the illness needs to be emphasised as employer pressure may result in patients returning to work too soon. The onus is on the doctor with current clinical responsibility for the patient to complete the fit note; this includes secondary care doctors.⁴¹ The content of the fit note should be agreed between the patient and doctor, including a “medically-recognised diagnosis”. For NHS staff to receive “covid pay” during absence, the fit note must mention covid.⁴² The ability to return to work after illness is a marker of recovery and clinicians must, therefore, record work status in the clinical notes in situations of chronic ill-health.^{43,44} From a public health perspective, counting days lost to sickness and lost income on account of long covid is essential.

Long covid, like all long-term conditions, impacts on many aspects of life and is best managed holistically with physical, psychological, and social factors addressed.⁴⁵ Prolonged illness following SARS-CoV-2 infection is characterised by the development of new symptoms at different timepoints.⁴⁶ Clinicians need to provide patient “safety-netting” advice and guidance on expected patterns of illness. Although NICE guidance states that new symptoms after three months are unlikely to be due to covid,⁴ this is not borne out by research⁴⁷ or patient and specialist experience. The importance of research into long covid was emphasised by many, which should include quantification of the burden of disease. To this end, one panellist advised that case reporting should be mandatory. The length of the World Health Organisation (WHO) long covid case report form was noted, however, and flagged as a potential barrier to case reporting in

practice. Patients should be made aware of research studies: participation could add meaning to what is often a very negative experience.

Ongoing exertional chest pain may warrant referral to a Rapid Access Chest Pain Clinic and/or cardiac MRI.⁴⁸ In patients diagnosed with myocarditis, exercise to 60% maximum heart rate can be advised but patients need to work out their own limits, which may be lower than this. In some cases of myocarditis or pericarditis, there is a difficulty in managing tachycardia and pharmacological approaches are needed, such as beta blockers or Ivabradine. As Colchicine and anti-anginals may also be helpful, advice from a long covid assessment clinic needs to be sought.⁸ PoTS (and other dysautonomic symptoms such as breathlessness, orthostatic intolerance, dizziness, and tremor) is an unfamiliar diagnosis for many clinicians, but seems to affect a significant subgroup of long covid patients. Whilst many would advocate specific investigation, NHS autonomic services are patchy, and if they are not to be overwhelmed, there will be significant educational needs for referring clinicians.⁴⁹ PoTS treatment can start with fluids, compression, and lifestyle adaptations (for which specific patient support materials are available),^{50,51} but may need to escalate to medication if symptoms are not improved.^{52,53} Midodrine may be helpful, though this is only available following secondary care initiating the prescription in many parts of the UK. There is an urgent need for research, education, and clear guidance to help GPs in managing this condition.

Similarly to treating urticaria, mast cell features require two-to-four-fold larger doses of antihistamines to suppress them.⁵⁴ Dermatologists and GPs with an interest in mast cell disorders have experience in counselling patients about such off-label use and an individual therapeutic trial is simple to arrange. Some patients exhibit sensitivity to histamine-rich foods and prominent GI symptoms (bloating, cramping pain, diarrhoea, acid reflux). These and other known triggers of mast cell activation should be avoided; the aim is to switch off the immune overreaction.⁵⁵ Unfortunately, H2-receptor antagonists are not readily available in the UK, although at time of writing Famotidine is again available. Further research including clinical trials are needed in this area, but the

recommendations represent a simple solution to dealing with very troublesome symptoms in some long covid patients.⁸ We used the term “breathing pattern disorder” to describe the subjective experience of patients which is not “breathlessness” in the strict sense of the word.⁵⁶ Its aetiology is unknown but may represent a disorder of central breathing control. Whilst specialist physiotherapy should be available to patients being seen in clinics, many patients seek help from alternative therapy, such as pranayama breathing. Meditation/mindfulness is promoted in the NHS as an effective therapy for anxiety and sensation of breathlessness.⁵⁷

It has been long established that chronic physical diseases have an increased risk of secondary mental health problems. A meta-analysis showed that 36.6% of people with a chronic physical disease had a co-existent mental health disorder.⁵⁸ Having a mental health disorder should not preclude investigation of any organic disease and unexplained symptoms or signs, and neuropsychiatric features should always prompt exclusion of organic pathology in the first place. Addressing epistemic injustice issues in the investigation and management of long covid should be a priority for local services.⁵⁹

Patients with long covid commonly refer to taking “the stack” or “the supplement stack”, which includes high dose vitamin C and D, niacin (nicotinic acid), quercetin, zinc, selenium, and sometimes also magnesium.⁶⁰⁻⁶² Further research is needed to confirm or refute supplement impact in long covid.⁶³ Examples of noteworthy interactions with supplements include: niacin causing an increased risk of bleeding events when combined with SSRIs or NSAIDs, increased risk of rhabdomyolysis together with statins,⁶⁴ and quercetin causing inhibition and induction of various human cytochrome P450 enzymes.⁶⁵

A recommendation concerning the name (“long covid”) did not obtain consensus and was ultimately excluded (Supplementary Table 1: item 37). In Round 1, we suggested the term “long covid” in preference to “Post Covid-19 Syndrome” and this achieved consensus (94% agreement); in Round 2, we decided to use the recently-approved WHO term “Post Covid-19 Condition” but this did not obtain consensus (64%

agreement). The naming of the condition is a subject of considerable controversy⁶⁶: “long covid” and “long-hauler” have been adopted by patients in the UK and USA (respectively) as neutral terms that make no assumptions about aetiology, presence/absence of ongoing infection, or prognosis.⁴ NICE suggested adoption of two terms: “Ongoing Symptomatic Covid-19” (4-12 weeks) and “Post Covid-19 Syndrome” (12+ weeks).⁴ In the USA the term “Post-Acute Sequelae of Covid-19” (PASC) has been adopted.⁶⁷ In contrast, the WHO have adopted “Post Covid-19 Condition” for ICD 11, reflected also in SNOMED coding.⁶⁸ This enables the creation of subclassifications or SNOMED coordinated terms for any future subcategories of long covid. Given the desire for international adoption and its use in coding, we accept the use of “Post Covid-19 Condition” as another medical term to describe long covid.

DISCUSSION

Summary

With a novel viral pathogen giving rise to a newly-recognised condition and a worldwide pandemic, there is at present very little empirical evidence on which to base recommendations for clinical care. Mechanisms which aim to identify, appraise, and use evidence in guidelines cannot function effectively when there is little evidence, often only hypothecated comparisons with other conditions. In this situation emerging clinical experience is the best practical guide to both practice and research. Cognisant of this, we used a robust Delphi method to derive 35 clear and practical recommendations to assist in the recognition, diagnosis, and management of patients with long covid.

Strengths and limitations

Recognising the need for strong consensus in the absence of evidence, we chose a high threshold for agreement (90%). The mid-point of the Likert scale (“neither agree nor disagree”) was interpreted as “agreement”, but classifying this as “disagreement” would affect only three recommendations (7, 17 and 33), whose consensus would fall

below 90% but nonetheless clear a threshold of 70%.⁶⁹ A rich set of views and relevant literature surrounding the recommendations has been summarised, even when a clear consensus was present.

Other than process, the key question for the validity of a Delphi study is the composition of the expert panel. We therefore recruited a panel of active “clinician-patients”, and augmented it with clinicians involved in newly-established long covid clinics. We ensured that a wide range of specialisms was represented in the panel, and that all panellists completed both rounds of the Delphi. This ensures that the recommendations were derived by clinicians highly active in this developing field rather than simply appraising an as-yet sparse literature. The recommendations were written in the spirit of promoting patient safety by ensuring differential diagnosis and potential for serious underlying conditions are considered. This is the precautionary principle. We accept that the full range of experience of long covid in the community is not yet known and that many patients will recover spontaneously without a need for treatment or referral. However, it is also quite possible that a lack of provision has failed to meet need, and many will have undiagnosed pathology.

Comparison with existing literature

The statements go further than NICE in many areas, particularly in the need to investigate potential cardiac conditions, dysautonomia, and immune dysfunction. The recently published NIHR “Living with Covid-19 – second review” echoes many of the themes explored here, including potential subgroups, need for investigation, and the relapsing-remitting nature of the condition, but offers a broad narrative rather than specific, practical statements for clinical use.⁷⁰ Practical management of such issues as PoTS and mast cell dysfunction (which are part of the therapeutic approach to chronic fatigue in the USA) can be very helpful for many long covid patients and need to be debated and tried in therapeutic settings, eventually as part of controlled studies.

Implications for research and practice

GPs need to be able to give practical advice to patients with long covid especially considering the simple nature of many treatments (e.g., compression stockings, fluids, and electrolytes for PoTS; over the counter antihistamines for mast cell suppression; pranayama breathing, meditation, and respiratory physiotherapy for breathing pattern disorder). We hope that these statements will act as a stimulus for research, medical education, and discussions around services for long covid. Research needs to prioritise rapid learning from long covid clinics, with mixed methods improvement science, learning from best practice by sharing data and by targeted mechanistic studies leading eventually to an evidence-based guideline on patient investigation, segmentation, and specific therapies. We acknowledge that there are significant challenges in resourcing services for long covid and that strict adherence to these recommendations will not always be possible but we need to meet the needs of this large group of high-need patients.

NHS England recently announced further funding for long covid clinics (there are no dedicated services in the other three UK nations). In due course, services for investigation, advice, and rehabilitation should be available to all practices. At present, however, the sheer practicalities of managing hundreds of thousands of patients after two waves of covid-19 mean that GPs need to develop confidence in good practice with long covid patients, basic investigations, deciding on need for referral or investigation, differential diagnosis, safety netting, empathy and support. These statements are a carefully considered and reasonable approach to helping patients until further evidence is available, generated by a robust consensus method from a unique group of “lived experience” professionals and front-line clinicians in the field.

ADDITIONAL INFORMATION

Additional authors forming the Delphi panel:

- Dr Nisreen Alwan (MBE, PhD, MBChB, MRCP, FFPH, MPH, MSc, FHEA; Associate Professor and Honorary Consultant in Public Health, University of Southampton and University Hospital Southampton NHS Foundation Trust);
- Dr Emily Attree (MBBS MA Camb MRCS MRCGP (2012); General Practitioner, Ridgmount Practice and The Lawson Practice);
- Dr Jennifer Blair (MB BCh, FFARCS; Epsom and St Helier University Hospitals NHS Trust);
- Dr Mary-Ann Bowen (MRCGP, BMedSci, BMBS, DCH, DRCOG, DFSRH; General Practitioner, Church Street Surgery, Wyre Forest Health Partnership);
- Dr Nicola J Brobbel (M.B.CH.B; Royal Surrey County Hospital / University of Surrey R and D);
- Dr Ciara Burgess (MB, BCh, BAO, MSc in Forensic Mental Health, Diploma in Management for Medical Doctors, MRCPsych; Consultant Psychiatrist, Norfolk and Suffolk NHS Foundation Trust);
- Dr Michael Cannell (MBBS DRCOG MRCGP MFHom (SpR) D.K (Bologna) BMAS (CoBC) PGDip WMA (Western Medical Acupuncture) PGDipNutrMed (Surrey); Bassett Clinic (St Albans), Hollybush Surgery (Hertfordshire), and Biolab Medical Unit (London));
- Dr Christopher Dixon (B Med Sci (Hons) BM BS MRCGP; General Practitioner);
- Dr Nell Freeman-Romilly (MA(Oxon), MSc(LSHTM), MBBS(Imperial); Oxford University Hospitals);
- Dr Sonali Gaur (MBBS, FRCOG; Frimley Health NHS Trust);
- Dr Thea Haldane (Bsc(Hons), MBBS, MMedEd, PGCert (clin nut), FRCP; Worcestershire Acute NHS Trust);
- Dr Melissa Heightman (FRCP; University College London Hospitals NHS Foundation Trust (UCLH));

- Dr Theresa Howe (MBBCh, BMedSci, MRCP (UK), FRCR; Consultant Clinical Oncologist, Velindre University NHS Trust, Cardiff);
- Dr Parul Kalia (MBBS; London North West University Healthcare NHS Trust);
- Dr Ramzi Khamis (MB ChB (Bristol) MRCP PhD FESC; Charing Cross Hospital and Hammersmith Hospital);
- Dr Muhammed Asad Khan (MBBS, FRCP, FRACP, PGCertClinEd (Auckland); Consultant Respiratory Physician, Manchester University NHS Foundation Trust);
- Dr Emma Ladds (MBBCh, MA(Oxon) MRCS MPH; GP Academic Clinical Fellow, Eynsham Medical Group and University of Oxford);
- Dr Amali Lokugamage (MBChB, BSc, MSc, MD, FRCOG, SFHEA; Whittington Health NHS Trust and University College London);
- Dr Harsha Master (BSc Infection & Immunity, MBBS, MRCGP; GP Lead Long Covid Rehabilitation Service, Hertfordshire Community Trust);
- Dr Rebecca Macfarlane (MA, MBBS, FRCEM, PhD; Epsom and St Helier University Hospitals NHS Trust);
- Dr Anna Paes (MRCGP, MBChB with Honours; General Practitioner, Richford Gate Medical Practice, Hammersmith and Fulham);
- Dr Sonia Parmar (MBBS, BSc, MRCP; Fairfield General Hospital);
- Dr Elizabeth Potter (MA, MBBCh, MRCGP; General Practitioner);
- Dr Manoj Sivan (MD, FRCP Ed; Associate Clinical Professor and Honorary Consultant in Rehabilitation Medicine, University of Leeds and Leeds Community Healthcare NHS Trust);
- Dr Sarah May Taylor (MBChB, MRCGP; General Practitioner, Ormskirk Medical Practice);
- Dr Margarita Thomson (BA, MA, BSc, MBBS, MRCGP (2010), DFSRH; General Practitioner, South West London Clinical Commissioning Group (Merton Borough) Cancer Clinical Lead and Macmillan GP);

- Dr Avril Washington (MBBS MRCP FRCPCH; Consultant Paediatrician, St George's University Hospitals NHSFT);
- Dr Katherine Wildon (MBChB, MRCP, DFSRH, DRCOG; General Practitioner).

To maintain the independence of the panel from authoring/publishing the recommendations, BD (corresponding author) and MN (first author) did not participate in the Delphi panel.

Acknowledgments:

The authors gratefully acknowledge infrastructure support from the NIHR Imperial Patient Safety Translational Research Centre and the NIHR Imperial Biomedical Research Centre.

Funding:

The first author is funded by the National Institute for Health Research (NIHR) Imperial Patient Safety Translational Research Centre (award number: PSTRC-2016-004); however, the present research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. No sponsor(s) or funder(s) had any role in the study design; in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. All authors had full access to all of the data in the study and accept responsibility to submit for publication. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Ethical approval:

This work is intended as a rapid guide to clinical practice in an emerging area and to inform priorities for research in long covid; as such, the Health Research Authority define the work as service development rather than research, not requiring ethical approval. Despite this, we followed best practice and Delphi participants gave consent to take part. All the participants in the study are study authors and approved the final manuscript.

Declaration of interests:

There are no conflicts of interest.

Author contributions:

BD devised the study and is guarantor, BD CR AF ST LJ NM MN wrote and revised the Delphi Statements and worked on the clinical content, MN managed the Qualtrics and Delphi processes and drafted the manuscript. Additional listed authors formed the rest of the Delphi panel and provided comments and references for the discussion on statements. All panellists were involved in re-drafting the manuscript and approving the final draft. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Accepted Manuscript – BJGP – BJGP:2021:0265

REFERENCES

1. Callard F, Perego E. How and why patients made Long Covid. *Soc Sci Med* 2021; **268**: 113426.
2. Office for National Statistics (ONS). Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK: 1 April 2021. 2021; published online Apr 1. <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/prevalenceofongoingsymptomsfollowingcoronaviruscovid19infectionintheuk/1april2021> (accessed Apr 4, 2021).
3. Gorna R, MacDermott N, Rayner C, et al. Long COVID guidelines need to reflect lived experience. *Lancet* 2021; **397**(10273): 455-57.
4. National Institute for Health and Care Excellence (NICE). COVID-19 rapid guideline: managing the long-term effects of COVID-19. 2020; published online Dec 18. <https://www.nice.org.uk/guidance/ng188> (accessed Mar 9, 2021).
5. Long Covid Support. Long Covid patient experience: NHS England Taskforce on Long Covid. 2021; published online Feb 4. <https://www.pslhub.org/learn/coronavirus-covid19/patient-recovery/long-covid-patient-experience-nhs-england-taskforce-on-long-covid-4-february-2021-r4209/> (accessed Mar 18, 2021).
6. Taylor AK, Kingstone T, Briggs TA, et al. 'Reluctant pioneer': a qualitative study of doctors' experiences as patients with long COVID. *Health Expect* Published Online First: Mar 22, 2021. <https://onlinelibrary.wiley.com/doi/10.1111/hex.13223> (accessed Mar 22, 2021). doi: <https://doi.org/10.1111/hex.13223>
7. National Institute for Health Research (NIHR). Living with Covid19: a dynamic review of the evidence around ongoing Covid19 symptoms (often called Long Covid). 2020; published online Oct 15. <https://evidence.nihr.ac.uk/themedreview/living-with-covid19/> (accessed Mar 22, 2021).
8. Dennis A, Wamil M, Kapur S, et al. Multi-organ impairment in low-risk individuals with long COVID. *medRxiv* 2020; published online Oct 16. <https://www.medrxiv.org/content/10.1101/2020.10.14.20212555v1> (accessed Mar 10, 2021).
9. Iacobucci G. Long covid: damage to multiple organs presents in young, low risk patients. *BMJ* 2020; **371**: m4470.
10. Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020; **584**(7821): 430-36. doi: 10.1038/s41586-020-2521-4
11. Sivan M, Taylor S. NICE guideline on long covid: research must be done urgently to fill the many gaps in this new "living guideline". *BMJ* 2020; **371**: m4938. doi: <http://dx.doi.org/10.1136/bmj.m4938>
12. Ayoubkhani D, Khunti K, Nafilyan V, et al. Epidemiology of post-COVID syndrome following hospitalisation with coronavirus: a retrospective cohort study. *medRxiv* 2021; published online Jan 15. <https://www.medrxiv.org/content/10.1101/2021.01.15.21249885v1> (accessed Mar 22, 2021).

13. Speets AM, van der Graaf Y, Hoes AW, et al. Chest radiography in general practice: indications, diagnostic yield and consequences for patient management. *Br J Gen Pract* 2006; **56**(529): 574-78.
14. National Institute for Health Care Excellence (NICE). Asthma: diagnosis, monitoring and chronic asthma management. 2017; published online Nov 29. <https://www.nice.org.uk/guidance/ng80> (accessed Mar 9, 2021).
15. Rosovsky RP, Grodzin C, Channick R, et al. Diagnosis and treatment of pulmonary embolism during the COVID-19 pandemic: a position paper from the national PERT consortium. *Chest* 2020; **158**(6): 2590-601.
16. Ackermann M, Verleden SE, Kuehnel M, et al. Pulmonary vascular endothelialitis, thrombosis, and angiogenesis in Covid-19. *N Engl J Med* 2020; **383**(2): 120-28.
17. Thillai M, Patvardhan C, Swietlik EM, et al. Functional respiratory imaging identifies redistribution of pulmonary blood flow in patients with COVID-19. *Thorax* 2021; **76**: 182-84.
18. Dhawan RT, Gopalan D, Howard L, et al. Beyond the clot: perfusion imaging of the pulmonary vasculature after COVID-19. *Lancet Respir Med* 2021; **9**(1): 107-16. doi: 10.1016/S2213-2600(20)30407-0
19. National Institute for Health and Care Excellence (NICE). COVID-19 rapid guideline: managing suspected or confirmed pneumonia in adults in the community. 2020; published online Apr 3. www.nice.org.uk/guidance/ng165 (accessed Mar 9, 2021).
20. Greenhalgh T, Javid B, Knight M, et al. What is the efficacy and safety of rapid exercise tests for exertional desaturation in covid-19? *Oxford COVID-19 Evidence Service* 2020; published online Apr 26. https://www.researchgate.net/profile/Trisha-Greenhalgh/publication/340934179_What_is_the_efficiency_and_safety_of_rapid_exercise_tests_for_exertional_desaturation_in_covid-19/links/5ea5ba53a6fdccd7945722aa/What-is-the-eficacy-and-safety-of-rapid-exercise-tests-for-exertional-desaturation-in-covid-19.pdf (accessed Mar 22, 2021).
21. Bui K-L, Nyberg A, Maltais F, et al. Functional tests in chronic obstructive pulmonary disease, part 1: clinical relevance and links to the international classification of functioning, disability, and health. *Ann Am Thorac Soc* 2017; **14**(5): 778-84.
22. du Bois RM, Weycker D, Albera C, et al. Six-minute-walk test in idiopathic pulmonary fibrosis: test validation and minimal clinically important difference. *Am J Respir Crit Care Med* 2011; **183**(9): 1231-7. doi: 10.1164/rccm.201007-1179OC
23. Zheng Y-Y, Ma Y-T, Zhang J-Y, et al. COVID-19 and the cardiovascular system. *Nat Rev Cardiol* 2020; **17**(5): 259-60.
24. Nishiga M, Wang DW, Han Y, et al. COVID-19 and cardiovascular disease: from basic mechanisms to clinical perspectives. *Nat Rev Cardiol* 2020; **17**(9): 543-58.
25. Huang L, Zhao P, Tang D, et al. Cardiac involvement in patients recovered from COVID-2019 identified using magnetic resonance imaging. *JACC Cardiovasc Imaging* 2020; **13**(11): 2330-39.
26. Luetkens JA, Isaak A, Zimmer S, et al. Diffuse myocardial inflammation in COVID-19 associated myocarditis detected by multiparametric cardiac magnetic resonance imaging. *Circ Cardiovasc Imaging* 2020; **13**(5): e010897.
27. Ho JS, Sia C-H, Chan MY, et al. Coronavirus-induced myocarditis: a meta-summary of cases. *Heart Lung* 2020; **49**(6): 681-85.

28. Dweck MR, Bularga A, Hahn RT, et al. Global evaluation of echocardiography in patients with COVID-19. *Eur Heart J Cardiovasc Imaging* 2020; **21**(9): 949-58.
29. Shchendrygina A, Nagel E, Puntmann VO, et al. COVID-19 myocarditis and prospective heart failure burden. *Expert Rev Cardiovasc Ther* 2021; **19**(1): 5-14.
30. Varga Z, Flammer AJ, Steiger P, et al. Endothelial cell infection and endotheliitis in COVID-19. *Lancet* 2020; **395**(10234): 1417-18.
31. Starekova J, Bluemke DA, Bradham WS, et al. Evaluation for myocarditis in competitive student athletes recovering from coronavirus disease 2019 with cardiac magnetic resonance imaging. *JAMA Cardiol* Published Online First: Jan 14, 2021. <https://jamanetwork.com/journals/jamacardiology/article-abstract/2775372> (accessed Mar 22, 2021). doi: 10.1001/jamacardio.2020.7444
32. Puntmann VO, Carerj ML, Wieters I, et al. Outcomes of cardiovascular magnetic resonance imaging in patients recently recovered from coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020; **5**(11): 1265-73. doi: 10.1001/jamacardio.2020.3557
33. Arunthari V, Burger CD. Utility of d-dimer in the diagnosis of patients with chronic thromboembolic pulmonary hypertension. *Open Respir Med J* 2009; **3**: 85.
34. Dani M, Dirksen A, Taraborrelli P, et al. Autonomic dysfunction in 'long COVID': rationale, physiology and management strategies. *Clin Med* 2021; **21**(1): e63.
35. Theoharides TC. Potential association of mast cells with coronavirus disease 2019. *Ann Allergy* 2021; **126**(3): 217-18. doi: 10.1016/j.anai.2020.11.003
36. Almeria M, Cejudo JC, Sotoca J, et al. Cognitive profile following COVID-19 infection: clinical predictors leading to neuropsychological impairment. *Brain Behav Immun Health* 2020; **9**: 100163. doi: 10.1016/j.bbih.2020.100163
37. Beaud V, Crottaz-Herbette S, Dunet V, et al. Pattern of cognitive deficits in severe COVID-19. *J Neurol Neurosurg Psychiatry* Published Online First: Nov 20, 2020. <https://jnnp.bmj.com/content/early/2020/11/19/jnnp-2020-325173.info> (accessed Mar 22, 2021). doi: http://dx.doi.org/10.1136/jnnp-2020-325173
38. Parisi S, Borrelli R, Bianchi S, et al. Viral arthritis and COVID-19. *Lancet Rheumatol* 2020; **2**(11): e655-e57. doi: 10.1016/S2665-9913(20)30348-9
39. Shepherd CB. NICE guideline on Long Covid fails to acknowledge important clinical and pathological overlaps with ME/CFS. *BMJ* 2020; **371**: m4938. doi: <https://doi.org/10.1136/bmj.m4938>
40. The Society of Occupational Medicine. COVID-19 return to work guide for recovering workers. 2021. https://www.som.org.uk/COVID-19_return_to_work_guide_for_recovering_workers.pdf (accessed Mar 10, 2021).
41. National Institute for Health Care Excellence (NICE). Workplace health: long-term sickness absence and capability to work. 2019; published online Nov 20. <https://www.nice.org.uk/guidance/ng146/chapter/Recommendations#assessing-and-certifying-fitness-for-work> (accessed Feb 19, 2021).
42. NHS Employers. COVID-19 sickness absence recording. 2020; published online Oct 2. <https://www.nhsemployers.org/news/2020/10/covid-19-sickness-absence-recording?fbclid=IwAR3eikoh6tSHdQbPjHFfPPcq8zP9z5I9M8VbpSiljyUQDKxZnDjTcJXBWc> (accessed Feb 19, 2021).
43. Public Health England. Work conversations in healthcare: how, where, when and by whom? 2019; published online Aug 19.

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/832876/Work_Conversations_in_Healthcare_How_where_when_and_by_whom.pdf (accessed Feb 19, 2021).

44. Waddell G, Burton AK. *Is Work Good For Your Health and Well-being?* London, UK: The Stationery Office, 2006.
45. Wade DT, Halligan PW. The biopsychosocial model of illness: a model whose time has come. *Clin Rehabil* 2017; **31**(8): 995-1004. doi: 10.1177/0269215517709890
46. Ontario Agency for Health Protection and Promotion (Public Health Ontario). *Long-Term Sequelae and COVID-19 – What We Know So Far*. Toronto, ON: Queen's Printer for Ontario, 2020. <https://www.publichealthontario.ca/-/media/documents/ncov/covid-wwksf/2020/07/what-we-know-covid-19-long-term-sequelae.pdf?la=en> (accessed Feb 28, 2021).
47. Altmann DM, Boyton RJ. Confronting the pathophysiology of long covid. 2020; published online Dec 9. <https://blogs.bmj.com/bmj/2020/12/09/confronting-the-pathophysiology-of-long-covid/> (accessed Feb 28, 2021).
48. Phelan D, Kim JH, Elliott MD, et al. Screening of potential cardiac involvement in competitive athletes recovering from COVID-19: an expert consensus statement. *JACC Cardiovasc Imaging* 2020; **13**: 2635-52. doi: 10.1016/j.jcmg.2020.10.005
49. Raj SR, Arnold AC, Barboi A, et al. Long-COVID postural tachycardia syndrome: an American Autonomic Society statement. *Clin Auton Res* 2021; published online Mar 19. <https://link.springer.com/article/10.1007/s10286-021-00798-2> (accessed Mar 22, 2021). doi: 10.1007/s10286-021-00798-2
50. Dysautonomia Support Network. 2019. <https://www.dysautonomiasupport.org> (accessed Mar 18, 2021).
51. Sheldon RS, Grubb BP, Olshansky B, et al. 2015 Heart Rhythm Society expert consensus statement on the diagnosis and treatment of postural tachycardia syndrome, inappropriate sinus tachycardia, and vasovagal syncope. *Heart Rhythm* 2015; **12**(6): e41-e63.
52. Gee ME, Watkins AK, Brown JN, et al. Ivabradine for the treatment of postural orthostatic tachycardia syndrome: a systematic review. *Am J Cardiovasc Drugs* 2018; **18**(3): 195-204.
53. Schreglmann SR, Büchele F, Sommerauer M, et al. Pyridostigmine bromide versus fludrocortisone in the treatment of orthostatic hypotension in Parkinson's disease – a randomized controlled trial. *Eur J Neurol* 2017; **24**(4): 545-51. doi: <https://doi.org/10.1111/ene.13260>
54. Afrin LB, Weinstock LB, Molderings GJ. Covid-19 hyperinflammation and post-Covid-19 illness may be rooted in mast cell activation syndrome. *Int J Infect Dis* 2020; **100**: 327-32.
55. The UK Mastocytosis Support Group. What triggers mast cells? 2021. <https://ukmasto.org/living-with-mcd/what-triggers-mast-cells/> (accessed Mar 10, 2021).
56. Patient-Led Research Collaborative. What does COVID-19 recovery actually look like? An analysis of the prolonged COVID-19 symptoms survey by patient-led research team. 2020; published online May 11. <https://patientresearchcovid19.com/research/report-1/> (accessed Mar 18, 2021).

57. Liang N-C, Visger TV, Devereaux A. Mindfulness for those with COPD, asthma, lung cancer, and lung transplantation. *Am J Respir Crit Care Med* 2020; **202**(4): 11.
58. Daré LO, Bruand P-E, Gérard D, et al. Co-morbidities of mental disorders and chronic physical diseases in developing and emerging countries: a meta-analysis. *BMC Public Health* 2019; **19**(1): 304. doi: 10.1186/s12889-019-6623-6
59. Goldbeck-Wood SJ. Minding the epistemic gap in covid-19 and beyond. *BMJ* 2020; **369**: m2379. doi: <https://doi.org/10.1136/bmj.m2379>
60. Alexandrov N. The team of doctors and biohackers who seem to be successfully treating “Long Covid”. 2020; published online Nov 12. <https://nkalex.medium.com/the-team-of-front-line-doctors-and-biohackers-who-seem-to-have-solved-long-covid-5f9852f1101d> (accessed Mar 10, 2021).
61. “RUN-DMC”. Here's how you treat long covid: lessons from MCAS. *YouTube* 2020; published online Nov 24. <https://www.youtube.com/watch?v=sICD0Kn6pR4> (accessed Mar 10, 2021).
62. “RUN-DMC”. 10 top tips for recovering from coronavirus: the lessons from 9 months of Long Covid studies. *YouTube* 2021; published online Jan 5. <https://www.youtube.com/watch?v=0EaHFuGJI78> (accessed Mar 10, 2021).
63. Mousavi S, Bereswill S, Heimesaat MM. Immunomodulatory and antimicrobial effects of vitamin C. *Eur J Microbiol Immunol (Bp)* 2019; **9**(3): 73-79. doi: 10.1556/1886.2019.00016
64. National Institute for Health and Care Excellence (NICE). British National Formulary (BNF). 2021; updated online Mar 4. <https://bnf.nice.org.uk> (accessed Mar 10, 2021).
65. Hakkola J, Hukkanen J, Turpeinen M, et al. Inhibition and induction of CYP enzymes in humans: an update. *Arch Toxicol* 2020; **94**(11): 3671-722. doi: 10.1007/s00204-020-02936-7
66. Perego E, Callard F, Stras L, et al. Why we need to keep using the patient made term “Long Covid”. *BMJ Opinion* 2020; published online Oct 1. <https://blogs.bmj.com/bmj/2020/10/01/why-we-need-to-keep-using-the-patient-made-term-long-covid/> (accessed Mar 22, 2021).
67. Rodriguez A. Dr. Anthony Fauci aims to answer 'a lot of important questions' about 'COVID long-haulers' in new nationwide initiative. *USA Today News* 2021; published online Feb 24. <https://eu.usatoday.com/story/news/health/2021/02/24/covid-19-long-haulers-fauci-announces-launch-nationwide-initiative/4572768001/> (accessed Mar 22, 2021).
68. SNOMED International. SNOMED CT COVID-19 Related Content. 2021; updated online Feb 2. <https://confluence.ihtsdotools.org/display/snomed/SNOMED+CT+COVID-19+Related+Content> (accessed Mar 9, 2021).
69. Diamond IR, Grant RC, Feldman BM, et al. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014; **67**(4): 401-09. doi: <https://doi.org/10.1016/j.jclinepi.2013.12.002>
70. National Institute for Health Research (NIHR). Living with Covid19 – second review: a dynamic review of the evidence around ongoing Covid19 (often called Long Covid). 2021; published online Mar 16. <https://evidence.nihr.ac.uk/themedreview/living-with-covid19-second-review/> (accessed Mar 19, 2021).

71. World Health Organization (WHO). WHO COVID-19: case definitions. 2020; published online Aug 7. https://apps.who.int/iris/bitstream/handle/10665/333912/WHO-2019-nCoV-Surveillance_Case_Definition-2020.1-eng.pdf?sequence=1&isAllowed=y&fbclid=IwAR06Y91HMyerQwOTfGVjmnFNYIv82c_fils6iw8SZx2YnmnncP7XjGrDGrE (accessed Mar 9, 2021).
72. World Health Organization (WHO). Global COVID-19 clinical platform Case Report Form (CRF) for Post COVID condition (Post COVID-19 CRF). 2021; published online Feb 9. [https://www.who.int/publications/i/item/global-covid-19-clinical-platform-case-report-form-\(crf\)-for-post-covid-conditions-\(post-covid-19-crf-\)](https://www.who.int/publications/i/item/global-covid-19-clinical-platform-case-report-form-(crf)-for-post-covid-conditions-(post-covid-19-crf-)) (accessed Mar 10, 2021).
73. Bhatia RT, Marwaha S, Malhotra A, et al. Exercise in the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) era: a question and answer session with the experts endorsed by the section of Sports Cardiology & Exercise of the European Association of Preventive Cardiology (EAPC). *Eur J Prev Cardiol* 2020; **27**(12): 1242-51.
74. Molderings GJ, Haenisch B, Brettner S, et al. Pharmacological treatment options for mast cell activation disease. *Naunyn Schmiedebergs Arch Pharmacol* 2016; **389**(7): 671-94.
75. Comas-Basté O, Sánchez-Pérez S, Veciana-Nogués MT, et al. Histamine intolerance: the current state of the art. *Biomolecules* 2020; **10**(8): 1181. doi: 10.3390/biom10081181

Accepted Manuscript – BJGP – E-GP – 2021-0265

Figure title:

Fig. 1. Number of panellists per specialty.

Box 1: Known examples of conditions associated with long covid.

- Myocarditis or pericarditis
- Microvascular angina
- Cardiac arrhythmias, including inappropriate sinus tachycardia, atrial flutter, atrial fibrillation, and high burden of ventricular ectopics
- Dysautonomia, including postural (orthostatic) tachycardia syndrome (PoTS)
- Mast cell activation, including urticaria, angioedema, and histamine intolerance
- Interstitial lung disease
- Thromboembolic disease (e.g. pulmonary emboli, microthrombi, or cerebral venous thrombosis)
- Myelopathy, neuropathy, and neurocognitive disorders
- Renal impairment
- New-onset diabetes and thyroiditis
- Hepatitis and abnormal liver enzymes
- Persistent GI disturbance including heartburn, diarrhoea, and loss of appetite
- New-onset allergies and anaphylaxis
- Dysphonia

Box 2: Recommendations pertaining to clinic organisation.

- 1) Consider long covid in patients with a clinical diagnosis of covid-19 as per WHO criteria⁷¹ or test-positive history with new or fluctuating symptoms including but not limited to breathlessness, chest pain, palpitations, inappropriate tachycardia, wheeze, stridor, urticaria, abdominal pain, diarrhoea, arthralgia, neuralgia, dysphonia, fatigue including neurocognitive fatigue, cognitive impairment, prolonged pyrexia, and neuropathy occurring beyond four weeks of initial covid-19** (strongly agree=19, 58%; agree=11, 33%; neither agree nor disagree=0, 0%; disagree=2, 6%; strongly disagree=1, 3%).
- 2) Multispecialty long covid clinics should be led by a doctor with cross-specialty knowledge and experience of managing this condition** (strongly agree=29, 88%; agree=2, 6%; neither agree nor disagree=1, 3%; disagree=0, 0%; strongly disagree=1, 3%).
- 3) Consider individualised investigations, management, and rehabilitation planning via a multispecialty long covid assessment service as local services allow. Prioritise physician-led medical assessments and diagnostics initially, and consider allied health professionals including physiotherapy and OT input as adjuncts** (strongly agree=23, 70%; agree=8, 24%; neither agree nor disagree=1, 3%; disagree=1, 3%; strongly disagree=0, 0%).
- 4) It is inappropriate for long covid clinics to be led by mental health specialists e.g., IAPT, clinical or health psychologist. They may be useful in supporting the multispecialty team but do not have the expertise to investigate and manage potential organ damage** (strongly agree=27, 82%; agree=5, 15%; neither agree nor disagree=1, 3%; disagree=0, 0%; strongly disagree=0, 0%).
- 5) All under-18-year-olds need access to similar services run by paediatric specialists with knowledge of how presentations and treatments differ to adults and with close liaison with school** (strongly agree=26, 79%; agree=7, 21%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).
- 6) Patients with comorbid mental health difficulties should have equal access to medical care as a patient without mental health difficulties and should not be triaged away from services** (strongly agree=28, 85%; agree=5, 15%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).

Box 3: Recommendations pertaining to diagnosis of underlying disorder.

General approach:

- 7) **In someone with long covid, symptoms of possible non-covid-19 related issues should be investigated and referred as per local guidelines. Long covid alone is not a sufficient diagnosis unless other causes have been excluded** (strongly agree=21, 64%; agree=8, 24%; neither agree nor disagree=2, 6%; disagree=1, 3%; strongly disagree=1, 3%).
- 8) **Carry out a face-to-face assessment including a thorough history and examination, consider other non-covid-19 related diagnoses, and measure FBC, renal function, CRP, LFT, thyroid function, HbA1c, Vitamin D, Magnesium*, B12, folate, ferritin, and bone studies** (strongly agree=24, 73%; agree=9, 27%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).

Respiratory:

- 9) **In those with respiratory symptoms, consider CXR at an early stage. Be aware that a normal appearance does not exclude respiratory pathology** (strongly agree=27, 82%; agree=4, 12%; neither agree nor disagree=1, 3%; disagree=1, 3%; strongly disagree=0, 0%).
- 10) **Be aware that simple spirometry may be normal but patients may have diffusion defects indicative of scarring, chronic PEs, or microthrombi. Consider referral to respiratory for full lung function testing** (strongly agree=23, 70%; agree=10, 30%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).
- 11) **Measure oxygen saturation at rest and after an age-appropriate brief exercise test in people with breathlessness and refer for investigation if hypoxaemic or if any desaturation on exercise** (strongly agree=17, 52%; agree=14, 42%; neither agree nor disagree=2, 6%; disagree=0, 0%; strongly disagree=0, 0%).

Cardiac:

- 12) **Consider the possibility of a cardiac cause of breathlessness** (strongly agree=27, 82%; agree=5, 15%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=1, 3%).
- 13) **Be aware that a normal D-dimer may not exclude thromboembolism, especially in a chronic setting, and referral for investigation is therefore indicated if there is a clinical suspicion of pulmonary emboli. Additionally, be mindful that thromboembolism may occur at any stage during the disease course** (strongly agree=26, 79%; agree=6, 18%; neither agree nor disagree=1, 3%; disagree=0, 0%; strongly disagree=0, 0%).

- 14) In patients with inappropriate tachycardia and/or chest pain, carry out ECG, troponin, Holter monitoring, and echocardiography. Be aware that myocarditis and pericarditis cannot be excluded on echocardiography alone** (strongly agree=22, 67%; agree=8, 24%; neither agree nor disagree=2, 6%; disagree=1, 3%; strongly disagree=0, 0%).
- 15) In patients with chest pain consider a referral to cardiology as cardiac MRI may be indicated in a normal echo to rule out myopericarditis and microvascular angina** (strongly agree=25, 76%; agree=6, 18%; neither agree nor disagree=1, 3%; disagree=1, 3%; strongly disagree=0, 0%).
- 16) In patients with palpitations and/or tachycardia, consider autonomic dysfunction** (strongly agree=25, 76%; agree=7, 21%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=1, 3%).

Others:

- 17) In patients with urticaria, conjunctivitis, wheeze, inappropriate tachycardia, palpitations, shortness of breath, heartburn, abdominal cramps or bloating, diarrhoea, sleep disturbance, or neurocognitive fatigue,⁵⁴ consider mast cell disorder** (strongly agree=15, 46%; agree=14, 42%; neither agree nor disagree=4, 12%; disagree=0, 0%; strongly disagree=0, 0%).
- 18) In patients with cognitive difficulties sufficient to interfere with work or social functioning, consider neurocognitive assessment** (strongly agree=23, 70%; agree=9, 27%; neither agree nor disagree=0, 0%; disagree=1, 3%; strongly disagree=0, 0%).
- 19) In patients with joint swelling and arthralgia, consider a diagnosis of reactive arthritis or new connective tissue disease and investigate and refer as appropriate** (strongly agree=20, 61%; agree=12, 36%; neither agree nor disagree=1, 3%; disagree=0, 0%; strongly disagree=0, 0%).

* Magnesium level may not be available in general practice.

Box 4: Recommendations pertaining to management: general approach.

- 20) For patients with fatigue and worsening symptoms hours to days following an activity, emphasise the importance of an initial phase of convalescence followed by careful pacing and rest** (strongly agree=27, 82%; agree=6, 18%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).
- 21) Support patients in shifting their mental timeline of recovery to reflect the likely prolonged course, with a possibly long phased return to work** (strongly agree=24, 73%; agree=9, 27%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).
- 22) Further support patients with signposting to patient resources. Applicable resources may include: management of post-exertional symptom exacerbation, activity pacing, acupuncture, diagnosis-specific management as relevant** (strongly agree=14, 42%; agree=16, 49%; neither agree nor disagree=1, 3%; disagree=2, 6%; strongly disagree=0, 0%).
- 23) Provide patients with signposting to social prescribing, sickness certification, and financial advice. Discuss with the patient whether sickness certification will state long covid as diagnosis** (strongly agree=26, 79%; agree=6, 18%; neither agree nor disagree=1, 3%; disagree=0, 0%; strongly disagree=0, 0%).
- 24) Clinicians should ensure that the occupational status of patients with long covid is recorded (in/out of work, part/full time, student)** (strongly agree=25, 76%; agree=8, 24%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).
- 25) Follow patients up regularly to monitor progress from a full biopsychosocial and occupational perspective** (strongly agree=19, 58%; agree=13, 39%; neither agree nor disagree=1, 3%; disagree=0, 0%; strongly disagree=0, 0%).
- 26) Encourage reporting of new symptoms (expected) and expectation of waxing-waning course** (strongly agree=25, 76%; agree=8, 24%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).
- 27) Consider contributing patient data to research on long covid, using the WHO Case Report Form or similar⁷²** (strongly agree=22, 67%; agree=9, 27%; neither agree nor disagree=2, 6%; disagree=0, 0%; strongly disagree=0, 0%).

Box 5: Recommendations pertaining to management: specific conditions.

- 28) Patients with cardiac symptoms should be advised to limit their HR to 60% of maximum (usually around 100-110bpm) and investigated with at least ECG and echocardiogram before taking up exercise. Supervised exercise testing should be considered for this patient group as they may have perimyocarditis and exercise carries risk of arrhythmia and worsening cardiac function⁷³ (strongly agree=16, 49%; agree=14, 42%; neither agree nor disagree=2, 6%; disagree=1, 3%; strongly disagree=0, 0%).**
- 29) For autonomic dysfunction including PoTS (Postural Orthostatic Tachycardia Syndrome), consider first increased fluids, salts, compression hosiery, and specific rehabilitation³⁴ (strongly agree=18, 55%; agree=13, 39%; neither agree nor disagree=2, 6%; disagree=0, 0%; strongly disagree=0, 0%).**
- 30) If PoTS and no or inadequate response to non-pharmacological therapy consider beta blocker, Ivabradine, or Fludrocortisone (with BP and response monitoring) (strongly agree=18, 55%; agree=13, 39%; neither agree nor disagree=1, 3%; disagree=1, 3%; strongly disagree=0, 0%).**
- 31) In patients with possible mast cell disorder, consider a one-month trial of initial medical treatment and dietary advice. Higher than standard dose of antihistamines are commonly used for this indication. If partial effect consider adding second level treatment such as Montelukast, as well as referral to allergy or immunology specialists^{74,75} (strongly agree=17, 52%; agree=14, 42%; neither agree nor disagree=2, 6%; disagree=0, 0%; strongly disagree=0, 0%).**
- 32) Be aware that adverse drug reactions are more common in patients with mast cell disorder, for example to beta lactam antibiotics, NSAIDs, codeine, morphine, or buprenorphine (strongly agree=17, 52%; agree=13, 39%; neither agree nor disagree=3, 9%; disagree=0, 0%; strongly disagree=0, 0%).**
- 33) For breathing pattern disorder, consider specialist physiotherapy and/or using alternative therapies such as pranayama breathing and meditation (strongly agree=12, 36%; agree=14, 42%; neither agree nor disagree=4, 12%; disagree=3, 9%; strongly disagree=0, 0%).**
- 34) In patients expressing distress, significant low mood, anxiety, or symptoms of PTSD, consider mental health assessment (strongly agree=20, 61%; agree=13, 39%; neither agree nor disagree=0, 0%; disagree=0, 0%; strongly disagree=0, 0%).**
- 35) Over the counter supplementation is common, including vitamin C, D, niacin (nicotinic acid), and quercetin. Be aware of significant drug interactions, such as with niacin or quercetin (strongly agree=21, 64%; agree=10, 30%; neither agree nor disagree=1, 3%; disagree=0, 0%; strongly disagree=1, 3%).**




Fig. 1. Number of panellists per specialty.

244x135mm (600 x 600 DPI)

Accepted Manuscript – BJGP – BJGP.2021.0265