British Journal of General Practice

Remote primary care during the COVID-19 pandemic for people experiencing homelessness: a qualitative study.

Howells, Kelly; Amp, Mat; Burrows, Martin; Brown, Jo; Brennan, Rachel; Dickinson, Joanne; Jackson, Shaun; Yeung, Wan-Ley; Ashcroft, Darren; Campbell, Stephen; Blakeman, Thomas; Sanders, Caroline

DOI: <https://doi.org/10.3399/BJGP.2021.0596>

To access the most recent version of this article, please click the DOI URL in the line above.

Received 18 October 2021

Revised 17 January 2022

Accepted 21 February 2022

© 2022 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by British Journal of General Practice. For editorial process and policies, see: <https://bjgp.org/authors/bjgp-editorial-process-and-policies>

When citing this article please include the DOI provided above.

Author Accepted Manuscript

This is an 'author accepted manuscript': a manuscript that has been accepted for publication in British Journal of General Practice, but which has not yet undergone subediting, typesetting, or correction. Errors discovered and corrected during this process may materially alter the content of this manuscript, and the latest published version (the Version of Record) should be used in preference to any preceding versions

Title page

Remote primary care during the COVID-19 pandemic for people experiencing homelessness: a qualitative study.

Dr Kelly Howells, Research Associate, PhD ^{1,2}

Kelly.howells@manchester.ac.uk

5th Floor, Williamson Building
The University of Manchester
Oxford Road, Manchester
M13 9PL.

Mr Mat Amp, Project Research Worker³

Canterbury Court (CC3.44)
1-3 Brixton Road
London
SW9 6DE

Mr Martin Burrows, Research Manager MSc³

(No business address available as works from home-formally employed by Groundswell).

Dr Jo Brown, Research Manager, PhD³

Canterbury Court (CC3.44)
1-3 Brixton Road
London
SW9 6DE

Miss Rachel Brennan HealthNow Lead, MA³

Canterbury Court (CC3.44)
1-3 Brixton Road
London
SW9 6DE

Ms Joanne Dickinson, Senior Nurse Practitioner, RGN⁴

Waters Meeting Health Centre, Waters Meeting Road, Bolton, BL1 8TT

Dr Shaun Jackson, GP, MD⁵

Urban Village Medical Practice
Ancoats Primary Care Centre
Old Mill Street
Ancoats
Manchester
M4 6EE

Dr Wan-Ley Yeung, GP, MD⁶

The Willow Tree Healthy Living Centre
94 Littleton Road
Salford
M7 3SE

Professor Darren Ashcroft, Professor of Pharmacoepidemiology, PhD¹
1st Floor, Stopford Building
School of Pharmacy
University of Manchester
Oxford Road,
M13 9PL.

Professor Stephen Campbell, Chair in Primary Care, PhD^{1,2}
7th Floor, Williamson Building
The University of Manchester
Oxford Road, Manchester
M13 9PL

Dr Thomas Blakeman, Senior Clinical Lecturer, MD, PhD²
5th Floor, Williamson Building
The University of Manchester
Oxford Road, Manchester
M13 9PL

Professor Caroline Sanders, Professor of Medical Sociology, PhD^{1,2}
5th Floor, Williamson Building
The University of Manchester
Oxford Road, Manchester
M13 9PL

1. **NIHR GM PSTRC, The University of Manchester**
2. **Division of Population Health, Health Services Research and Primary Care, Centre for Primary Care and Health Services Research, The University of Manchester.**
3. **Groundswell, London.**
4. **Bolton NHS Foundation Trust**
5. **Urban Medical Village, Manchester.**
6. **Salford Together, Salford CCG.**

Remote primary care during the COVID-19 pandemic for people experiencing homelessness: a qualitative study.

Background

The COVID-19 pandemic has caused unprecedented disruption and change for the organisation of primary care, **including for** people experiencing homelessness who may not have access to a phone. Little is known about whether the recent changes required to deliver services to people experiencing homelessness will help to address or compound inequality in accessing care.

Aim

To explore the experience and impact of organisational and technology changes in response to COVID-19 on access to healthcare for people experiencing homelessness.

Design and setting

An action led and participatory research methodology was employed in 3 case study sites made up of primary care services delivering care for people experiencing homelessness.

Method

Individual semi-structured interviews were conducted with 21 people experiencing homelessness and 22 clinicians and support workers. Interviews were analysed using a framework approach.

Results

The move to remote telephone consultations highlighted the difficulties experienced by participants in accessing healthcare. These barriers including problems at the practice level associated with remote triage as participants did not always have access to a phone or the means to pay for a phone call. This fostered increased reliance on support workers and clinicians working in the community to provide or facilitate a primary care appointment.

Conclusion

The findings have emphasised the importance of addressing practical and technology barriers as well as supporting communication and choice for mode of consultation. We argue that consultations should not be remote 'by default' and instead take into consideration both the clinical and social factors underpinning health.

How this fits in

Pre-COVID-19, there was mounting policy pressure to implement digital innovations to increase staff capacity and access to primary care services. However, there were concerns about a so-called 'digital divide' with certain population groups being left behind and excluded from potential benefits of technologies that could further exacerbate existing health inequalities. This qualitative study explores how the rapid change to remote care during the COVID-19 pandemic impacted the access and safety of care for people experiencing homelessness. We argue that while the remote model may facilitate more 'timely' access for some individuals, remote care should not be the default approach. The evidence highlights that the remote model may compromise a relationship-based approach, which is particularly important for building trust and continuity and to mitigate against existing health inequalities.

Introduction

Homelessness is a persistent and growing problem in **England and elsewhere (1). Shelter calculates approximately 253,000 people are experiencing homelessness in England**, with many thousands more at risk (2, 3). People experiencing homelessness are heterogeneous falling into multiple categories of homelessness including: roofless (rough sleepers or in emergency accommodation); houseless (those in homeless shelters); insecure (under threat of eviction); and inadequate (people living in temporary or unfit structures) (4). The numbers of people falling in these categories are a moving picture, and the numbers in hotels and other temporary accommodation increased during Covid-19 lockdowns periods. Recent evidence shows widening health inequalities for people experiencing homelessness (5) with common experiences of both acute and ongoing complex mental, social and physical care needs and reduced life expectancy (6). Inequitable access to healthcare is identified as a key social determinant of health (7, 8). Known as the 'inverse care law', those most in need of healthcare are often the least likely to receive it (9-11).

The NHS Long Term Plan (2019) seeks to close the inequality gap and recommends prioritisation of underserved groups in primary care (12). Dedicated homeless and inclusion health services are available within mostly urban areas in the UK and they provide 'drop in' and out of hour's clinics and outreach healthcare (13-15). However COVID-19 has disrupted

this model of care with telephone or video consultations recommended to reduce infections (16). Pre-COVID-19, there were already calls to implement remote care via digital innovations to maximise staff capacity and access to primary care services(12). However there were concerns about a 'digital divide' (17, 18) and compounding inequalities for groups that are digitally excluded (18, 19). Digital transformation to remote services including triage, as well as consultations, also raises questions about how to support people to access the right type of consultation (face-to-face, phone, or video) appropriate for their needs. This has implications regarding clinician-patient relationships (14, 20-22) such as relational continuity of care, which is viewed by The Royal College of General Practitioners (RCGP) as crucial in the diagnosis and treatment of conditions, especially amongst people with long-term or sensitive health conditions where stigma is often experienced (23-25).

Through data drawn from qualitative research methods, we explored how people experiencing homelessness accessed primary care services during the COVID-19 pandemic. We also explored how clinicians and support workers **(based in hostels or charities)** provided and facilitated this care to gauge how the change to remote consultations impacted access and service delivery within primary care.

Methods

An action led and participatory qualitative research methodology was employed in three case study sites **from September 2020 till the end of January 2021** (see table 1), made up of two commissioned primary care services and a community nurse inclusion health service. Three separate urban case study sites were chosen to compare organisational and local policy context. This included interviews with clinicians, stakeholders and people experiencing homelessness. Action research and participatory methods refer to styles of research emphasising collaboration and democratic working between multiple partners to bring about change (26, 27). It has been widely used for conducting community or service-based research, including amongst people experiencing homelessness and service providers (28-31).

Table 1 HERE

Flexible semi-structured interviews were conducted with health professionals and other

staff (such as care coordinators and managers) at each case study site who also facilitated snowball sampling of additional stakeholders that worked closely with the case study sites (such as hostel workers, volunteer organisations). Due to COVID-19 social distancing restrictions, all interviews were conducted remotely over the telephone and were audio recorded. Interviews with clinicians and other stakeholders were conducted by KH. To observe change in practice through the pandemic, one GP kept an audio diary to record their daily observations. We held team meetings with the primary care providers every three months to reflect on organisational changes and issues across the case study sites in 'real time', and to discuss emerging findings regarding barriers to accessing care. Notes were taken in these meetings but they were not audio recorded.

The interviews with people experiencing homelessness were led by a researcher (MH) employed by Groundswell, (a homeless charity) following a peer research approach (32-34). Researchers with lived experience of homelessness receive research skills training (via Groundswell) to enable them to more easily engage with the target research audience, who may be wary of participating in research. Their shared experience sought to promote a sense of trust and integrity, which encourages participation in research and may help people speak more openly about their experiences (34-36).

As a result of social distancing measures, we engaged with local hostels, day centres, charities and outreach healthcare workers in the case study sites to act as gatekeepers for recruitment. Sampling was pragmatic aiming to maximise diverse participation but the final sample was dependent on responses to information disseminated via gatekeepers. Participants were eligible to participate if they were currently homeless or had recent (during the pandemic) lived experience of homelessness. An information sheet and consent form were given to eligible participants and they contacted the researcher directly if interested. In some circumstances, the gatekeeper facilitated contact where participants did not have access to a phone. Participants experiencing homelessness were asked to consent to a follow up interview 3-6 months after the initial interview to reflect **on their experiences of changes to accessing care during the pandemic. All interviews were audio-recorded using an encrypted digital recorder and were immediately downloaded to an encrypted university. Audio files were then transferred to an approved university transcription**

provider as an encrypted file. Once transcribed, all audio recordings were deleted and transcripts were pseudonymised before sharing with the wider team for data analysis.

Ethical approval was obtained from a NHS REC committee (London Stanmore REC 20/HRA3356). Interview schedules were developed from the research and policy literature, with some changes added following interviews and research meetings with primary care providers.

Analysis

The audio recorded interviews and audio diary were transcribed verbatim and analysed using a framework method, producing a matrix of summarised data and a structure to analyse and summarise data within key themes. It is particularly suited to enabling analysis effectively within teams of multiple researchers (37, 38). The approach allows for analysis according to predefined themes and themes that emerge more inductively from the data (37). Our initial analytical framework was completed in Microsoft Excel and structured by the main aspects of experience reflected in the literature on homelessness and access (39). The University research team worked closely with the research lead and researchers from Groundswell to do all stages of coding and thematic analysis. To increase rigour, the initial coding of the interview transcripts was conducted separately by KH (UoM) and MA (Groundswell) to detect similarities and variations of interpretation across the sample. A sample of transcripts were also read and coded by the CS (UoM) and MB (Groundswell) to develop the main and sub-themes. This formed the basis for further discussion and critical analysis for further iteration of the main themes within the wider team, including our primary care partners.

Results

Sample Characteristics

In total, 22 stakeholders, including GPs (n= 4), community and practice nurses (n=7) other staff based in general practice (n=2) and support workers based in hostels or working in outreach (n=9) were interviewed across 3 case study sites in North West England. Twenty-one people experiencing homelessness were also interviewed, including 6 follow up

interviews to gauge any changes in accessing care over the course of the pandemic (see table 2).

Table 2: Demographic characteristics of people experiencing homelessness sample (n=21)

TABLE 2 HERE

Findings

1. The impact of remote triage on accessing appointments

Experiences of remote care were mixed with some participants preferring the change as it offered them more convenient and **timely access**. This was particularly the case for participants that struggled to attend GP appointments physically due to anxiety or practical barriers and for patients that needed a medication review or a repeat prescription, as such requests were easily resolved over the phone.

However, several frustrations with new COVID19 lockdown enforced appointment systems were reported. Participants missed the ability to 'drop in' at their GP practice or daycentre for a same day appointment. All appointments needed to be made either online or by phone with patients advised to call early in the morning for a same day appointment. This often resulted in long waiting times to get through, **with some participants struggling to make appointments as they did not always have access to a phone or the funds to pay for the phone call**:

It is harder now because you're just on hold all the time. So when you've got credit, it sort of runs out. If you're on a contract it wouldn't bother you (ID 14, Male participant experiencing homelessness, aged 35-44, white British, more than 4 chronic conditions).

I found that really quite difficult with the phone calls and stuff like that. Like getting through the doctors. It was really hard work. Not all the time do I have (access to) a computer or like a phone with internet on it. The staff obviously here at the hostel are busy, so we can't really use the phones or use the computer. You can't even touch the computer because of the coronavirus pandemic that's going on and social distancing. I just find it, like the whole... it's really hard (ID 8, Female experiencing homelessness, aged 35-44, white British, 4 chronic conditions).

Once successfully through to the practice, patients then had to discuss their health concern with the GP receptionist to determine the urgency of their concern and the nature of appointment required (triage). Participants were **sometimes** reluctant to discuss concerns with receptionists **as they are not medically qualified, with some participants referring to them as strangers:**

Yeah, it's an issue for me because I've got such a complicated medical history with having cancer and the permanent side effects from the treatments that I've got now, I don't want to be explaining them to a stranger (ID 11, Male experiencing homelessness, aged 25-34, Eastern European, one long term condition).

Once triage was completed, participants were not always given a specific appointment time, which sometimes resulted in missed or delayed consultations. Participants with immediate health concerns expressed concern that a late call back would delay access to urgently needed medications or treatment. This resulted in some participants accessing care elsewhere, such as at a pharmacy or walk-in centre. **This frustration was echoed by the community nurses interviewed, as they often had to support their patients to reorganise the telephone consultations they had missed as appointment times was not always specified:**

Like today I went into one of the hostels and a lady in there who I went to see, to provide wound care, she said that the GPs had phoned... they said they would phone her at nine o'clock and they didn't ring her until four o'clock in the afternoon. So she missed that call because again, it wasn't that they stuck to the agreed time (Community nurse, ID 6).

2. Consequences of remote care for communication and therapeutic relationships

Despite the possibility of delivering remote consultations by telephone or video, all the participants interviewed had only participated in a telephone consultation. At the beginning of the pandemic, video-consultations were encouraged to bridge the gap between remote and face-to-face care but, the clinicians interviewed found this mode of consulting sometimes difficult and it was therefore rarely offered to patients experiencing homelessness:

Speaking to patients who get a really poor reception and phones are cutting out. Some use Wi-Fi in the hostel and it's not very good Wi-Fi and then there is a lag with the speech and the picture. It becomes pixilated and this is unhelpful. The technology has got to be good enough quality for us to do our work (GP 1, audio diary).

Two of the case study sites primarily preferred to use telephone consultations to provide care. At the practical level, both clinicians and patients noted that technical issues could impede communication during the consultation with poor phone signal a recurring problem. The clinicians interviewed were concerned that telephone consultations could potentially compromise patient safety as opportunities for patients to discuss other health issues would not necessarily arise as non-verbal aspects of the consultations were described as 'lost', making it more **difficult to provide a timely diagnosis and assess risk**. Body language was considered important in assessing the wellbeing of patients as often they struggled to discuss social and mental health issues with clinicians they did not always know or trust:

It's like clues that you pick up isn't it from people's nonverbal communication as well, just the way they present, their eye contact. Whether somebody's depressed, withdrawn. A lot of these, it's not just that...you can't gather that over a telephone (Community Nurse ID 6).

This view was shared by some participants who found it difficult talking to their GP over the phone with the sensory aspects of the consultation implicitly linked to safety. This was reflected particularly in concerns highlighted about remote clinical assessment as a basis for diagnosis and management:

Seeing a doctor face-to-face is a lot more personal, and then you can show the doctor where you're hurting. I'm finding it quite difficult to understand how doctors can make diagnosis over a phone when they're not seeing it (ID 11, Male experiencing homelessness, aged 25-34, Eastern European, one long term condition).

The dynamics of the doctor-patient relationship were described as ‘transactional’ by some clinicians as consultations were primarily symptom focussed. However, a pre-existing relationship between doctor and patient was considered to mitigate against the focus on diagnosis:

If you have the right rapport with a patient I think this kind of consultation can still work but I guess I question new patients because it's difficult to build that early rapport without being there in the same room as someone and there's much more of a transactional exchange....Pre-existing relationships can make the consultation easier and less 'transactional' (GP 1).

This dynamic was also reflected by some participants experiencing homelessness who struggled to verbalise their concerns, particularly their social situation, over the phone. One participant felt they received less ‘empathy’ over the phone, perhaps due to the more transactional nature of telephone consultations:

If it was face to face so much easier because you can explain your situation to gain either empathy or...I don't want sympathy but somewhere in the middle of these, do you know what I mean? But I'll tell you what you get, you do get something in the middle because you get apathy (ID 3, Male experiencing homelessness, aged 45-54, white British, 3 chronic conditions).

Although some participants felt nervous and distrustful of remote care, others found the anonymity empowering as it avoided perceived judgement and therefore reduced anxiety:

*Because of the anxiety as well, I don't really like associating with people, so it **makes it (having a consultation over the phone)** a bit easier for me (ID 10, Female experiencing homelessness, aged 45-54, white British, one chronic condition).*

3. Facilitating access to care: collaborative working across health and social care boundaries

The three case study sites often worked in collaboration with community and social care sectors pre-pandemic. The pandemic highlighted the importance of this collaborative approach with hostel and charity outreach workers being key facilitators of care. This included making appointments/medical queries on behalf of people experiencing homelessness, supplying pre-paid phones and digital devices to hostels (such as iPads), to enable easier and timely access, engaging with people to raise awareness of organisational changes, facilitating GP registration and discussing specific patients as part of a multidisciplinary team meeting (MDTs) within the primary care network (PCNs).

Not all the participants interviewed were registered with an enhanced general practice or had access to support via a hostel or outreach service. In one case study site (3), care continued to be provided face-to-face by the community nurse team during the first and subsequent lockdowns via both a walk-in clinic and outreach care. However, as the only face-to-face provider of care in the local area, their roles evolved to become not only providers of care but 'mediators'. This involved making appointments on behalf of patients and facilitating appointments by allowing their patients to use their smart phones. The nurses interviewed often went above and beyond their expected role and working hours to secure appointments for their patients. As one nurse described, they became the 'eyes and ears' of the GP.

I think the main thing is, is not being able to access GPs for patients, you know, most patients don't have telephones to make telephone calls. So they would come to our clinic, and if they needed a telephone consultation with the GPs, they would ask us if we would be able to do that for them. And then you try ringing the GP up, and unfortunately you have to ring at a certain time, which is eight o'clock in the morning. Now I don't work at eight o'clock, but I know in the past I have rung to make appointments for patients at eight o'clock in the morning, you know, which can be...you know, I'm not in work, I've had to go out of my house to do this telephone call (Community nurse, ID 10).

The community nurse team was particularly helpful for participants who were not currently registered with a GP and were either unaware of their right to register or they lacked the confidence to challenge general practices that refused registration requests. The

community nurses would advocate for their patients and ensure they were not illegitimately turned away from practices. Over the course of the pandemic, this prompted a new partnership between the community nurses and local GPs who worked together to manage the complex needs of people experiencing homelessness:

GPs have had to learn a new way of working. It's not easy for them because they've had to learn new telephone skills and trying to do consultations over the phone without actually visually looking at a patient. We have had issues where we've actually gone to see a patient because they couldn't get a GP appointment. We've basically been the GPs eyes, to sort of say, look X, Y and Z is happening here...Yeah, it's very difficult. I mean, we do feel that we've plugged the gap for a lot of services (Community nurse, ID 6).

However the community nurse team and support workers interviewed found it more difficult to facilitate for people registered with a mainstream primary care service as they were not always aware of or had received inclusion healthcare training:

You know, our patients trust us, they respect us, we don't get any grief from anybody. Our patients unfortunately get grief from other health professionals, which isn't fair. It's judgemental, but you know unless you work with people who are dependent on drug or alcohol or both, plus have mental health issues, unless you work in that field all the time, it is difficult not to judge people (Practice Nurse).

For this reason, there was concern from the support workers and clinicians as to 'who was being missed' as it was acknowledged that not all people experiencing homelessness would have access a support worker or outreach team to facilitate the healthcare system for them. For this reason, the support workers interviewed often encouraged their clients to register with an enhanced GP service if possible as these services were more aware of the barriers faced by people experiencing homelessness during the pandemic:

I am aware that there are people now who will need an individual to help them access health services, an individual like myself who can liaise with somebody. In a way that introduces an extra barrier if you don't have access to IT. Whereas if there was a drop-in service that they could access at times that they chose themselves they may get to talk to

somebody faster.....I have quite close connections with homeless primary care services but if there's another GP service there are a lot more barriers there, because they're not used to dealing with a third party, they've not had any contact with me before. That's why I would encourage somebody to register with a specialist homeless GP service unless they were very keen to remain at their current GP service (Outreach worker, ID 62).

However the GPs interviews highlighted that relying on a support workers could eventually be problematic for anyone experiencing homelessness as they may lose touch with their support worker or no longer have access once they transition to living in permanent accommodation. For this reason, the GPs and nurses interviewed felt it was important to support and enable people to access healthcare services for themselves by offering a 'flexible' model of care that recognised that services could not be exclusively remote or digital:

The system (remote only) how it currently stands will perpetuate inequality – there are positives but there are negatives but the people who will experience problems using it often don't have a voice so it is sometimes assumed it is working (GP 2).

Discussion

Summary

This study identified key access points and barriers to accessing, receiving and delivering primary care, at a time of rapid transformation, during the COVID-19 pandemic for people experiencing homelessness. While some of the barriers existed pre-COVID-19, the move to remote telephone consultations brought sharply into focus the difficulties experienced by participants, including problems at the practice level associated with remote triage, as participants did not always have access to a smart phone to enable a video consultation or the means to pay for a phone call. This fostered increased reliance on support workers and clinicians working in the community to provide or facilitate a primary care appointment. At the interactional level, people experiencing homelessness and health staff found that face-to-face care that enabled eye contact and non-verbal communication was important for patient safety (including for those with complex needs and other vulnerable groups). Such direct interaction can be important for **making a timely diagnosis** and continuity of care, but

also in enabling rapport and trust so that sensitive issues, such as mental and sexual health were more likely to be discussed.

We found that face-to-face interactions- remained a fundamental aspect of delivering healthcare during the pandemic delivered either by clinicians in general practice, the wider primary health care team or **facilitated by support workers and outreach teams**. The findings highlight that the responsiveness and success of implementing a remote model for people experiencing homelessness relies heavily on flexible and collaborative working across health **and community organisations**. This involves nurses and support workers, having face-to-face contact and supporting access for remote consultations where needed.

Strengths and limitations

We endeavoured to capture a wide range of diverse experiences by engaging with local stakeholders and outreach workers to act as gatekeepers to recruit people experiencing homelessness. While we acknowledge these findings will not be representative of all people experiencing homelessness, a key strength of our work is that we captured the patient and clinician experience. However, quantitative data may have provided more evidence on variation, such as mode of consultation and number of appointments.

Peer-led research is associated strongly with social change, inclusivity and breaking down the traditional hierarchies between researchers and participants(34) although some argue that peer-researchers can lack detachment from the research issue leading to bias in findings (36, 40). However, a key strength is that all research interviews were reviewed by the core research team and feedback provided to help guide ongoing interviews.

Comparison with existing literature

Much of the existing literature on the use of digital health innovations for people experiencing homelessness originates from the US (41). It has demonstrated that telephone consultations can improve access to care for patients and reduce workload for staff, and some research has pointed to enhanced benefits of video consultations, particularly when patients are anxious (42). Systematic reviews indicate they can provide safe and high quality care for 'clinically appropriate' patients but most evidence has been based on people with stable chronic conditions (17, 43). Greenhalgh et al (16) suggest that within the context of

COVID-19, video and telephone consultations are unlikely to be appropriate replacements for seriously ill patients or for those who are not experienced using digital technologies (44). Relationship-based care is a key strategic priority of the RCGP. A relationship-based care model is centred on a '*high quality relationship between doctor and patient*', in which continuity, compassion and trust are deemed essential in improving the quality and safety of care (22). However, the RCGP has raised concerns that the relationship-based care model has been compromised by the transactional nature of remote telephone care during the pandemic. (45). Our research supports this view as the interviews with participants experiencing homelessness and clinicians alike highlight how implementing the relationship-based model can be more challenging when consulting remotely, particularly if there is no prior relationship or the patient is vulnerable or dealing with sensitive and complex health issues. We argue that the relational aspects of care are an important in enabling access to care and reducing health inequalities. This relationship-based approach extends to non-clinical primary care staff that often engage with people experiencing homelessness at the first point of access. Our findings highlight the importance of having responsive and flexible organisational processes in place to enable a relationship-based approach at each stage of the patient journey.

Implications for practice

While a collaborative approach mitigated against some of the potential pitfalls of the remote model, there was recognition from the clinicians interviewed that not all people experiencing homelessness, particularly the 'hidden' homeless and people at severe risk of homelessness would necessarily have someone to facilitate an appointment for them. For this reason, case study sites 1 and 2 adapted their organisational processes to accommodate the complex needs of people experiencing homelessness. There was recognition that long-term organisational changes were required to enable access, such as the re-introduction of 'flexibility' that had not been possible at the beginning of the pandemic. In the case study sites, this included adapting appointment protocols so that people experiencing homelessness could bypass triage and be offered a face-to-face appointment if they appeared distressed or unable to articulate their health problem. Equally, we acknowledge that decisions regarding the mode of the consultation are complex and in some cases in order to offer timely access to care, a telephone consultation may be most suitable. We

recognise the limits of workforce capacity in providing face to face care for all patients and that certain trade-offs are inherent with this approach, particularly for the fostering of trusting relationships, reflected in the reticence of participants to discuss their health needs with a receptionist and their ambivalence regarding the safety of the care they were receiving. The study highlights that the development of support roles within PCNs, such as care coordinators, are important not only in facilitating timely access to care but they also enable the embedding of relationship-based care into and across routine general practice (46).

While there are undoubtedly benefits to the remote model, we argue that remote care should not be the default approach. The vulnerability of the patient in addition to their clinical needs should be taken into consideration. The evidence highlights that a relationship-based approach is important for building trust and continuity and to mitigate against existing inequalities. As suggested in recent related research, there should be more of a focus on collaborative working within the NHS (including secondary care and community pharmacy) and other services, such as charities, hostels, alcohol and drug services and job centres that have specific knowledge and experience of working with people experiencing homelessness and other vulnerable groups. Specific training on inclusion health for reception staff across primary care and not just for clinicians and others working in enhanced services would also be beneficial (14, 47).

Funding statement:

This work was funded by the National Institute for Health Research (NIHR) through the Greater Manchester Patient Safety Translational Research Centre (award number: PSTRC-2016-003).

Disclosure statement:

The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

References

1. Fitzpatrick S, Pawson H, Bramley G, et al. The homelessness monitor: England 2019. London: Crisis. 2019. <https://researchportal.hw.ac.uk/en/publications/the-homelessness-monitor-england-2019> (accessed 13th December 2021).
2. Shelter. Far from Alone: Homelessness in Britain in 2017, London. 2017. https://england.shelter.org.uk/professional_resources/policy_and_research/policy_library/report_far_from_home_alone (accessed October, 2021).

3. Shelter. Homeless in a pandemic: 253,000 people are trapped in temporary accommodation. 2020https://england.shelter.org.uk/media/press_release/homeless_in_a_pandemic_253000_people_are_trapped_in_temporary_accommodation.
4. Edgar B. The ETHOS definition and classification of homelessness and housing exclusion. *Eur J Homelessness*. 2012;6(2):219-25.
5. Marmot M, Allen J. COVID-19: exposing and amplifying inequalities. *J Epidemiol Community Health*. 2020;74(9):681-2.
6. Marmot M. Health equity in England: the Marmot review 10 years on. *Bmj*. 2020;368.
7. Marmot M. Social determinants of health inequalities. *The lancet*. 2005;365(9464):1099-104.
8. Marmot M. Fair society, healthy lives. *Fair society, healthy lives*. 2013:1-74.
9. Hart JT. The inverse care law. *The Lancet*. 1971;297(7696):405-12.
10. Mercer SW, Guthrie B, Furler J, et al. Multimorbidity and the inverse care law in primary care. *BMJ* 2012; 344 :e4152.
11. Moss C, Sutton M, Cheraghi-Sohi S, et al. Comparative 4-year risk and type of hospital admission among homeless and housed emergency department attendees: longitudinal study of hospital records in England 2013–2018. *BMJ Open*. 2021;11(7):e049811.
12. England N. The NHS long term plan. London: NHS England; 2019. 2019. <https://www.england.nhs.uk/long-term-plan/> (accessed October 2021).
13. Brennan R. Delivering Healthcare to Homeless People: lessons from Scandinavia and the USA. Winston Churchill Trust. 2018.
14. Hirst V, Cuthill F. An exploration of GP care in outreach settings for people experiencing homelessness: a qualitative study. *Br J Gen Pract*. 2021; 71 (709): e596-e603. DOI: <https://doi.org/10.3399/BJGP.2020.0749>
15. Clark E, Player E, Gillam T, et al Evaluating a specialist primary care service for patients experiencing homelessness: a qualitative study. *BJGP Open*. 2020;4(3): bjgpopen20X101049.
16. Greenhalgh T, Wherton J, Shaw S, et al Video consultations for covid-
19. *BMJ* 2020; 368 :m998 doi:10.1136/bmj.m998
17. Greenhalgh T, Papoutsi C. Spreading and scaling up innovation and improvement. *BMJ*. 2019; 365 :l2068 doi:10.1136/bmj.l2068.
18. Sanders C, Rogers A, Bowen R, et al. Exploring barriers to participation and adoption of telehealth and telecare within the Whole System Demonstrator trial: a qualitative study. *BMC Health Serv Res*. 2012;12(1):220.
19. Ong BN, Hodgson D, Small N, et al Implementing a digital patient feedback system: an analysis using normalisation process theory. *BMC Health Serv Res*. 2020;20:1-16.
20. Greenhalgh T. Future-proofing relationship-based care: a priority for general practice. *Br J Gen Pract*. 2014;64(628):580-.
21. Jones R. General practice in the years ahead: relationships will matter more than ever. *Br J Gen Pract*; 2021; 71(702):4-5. doi: 10.3399/bjgp21X714341.
22. Practitioners TRCoG. General practice COVID-19 recovery: the power of relationships. <https://www.rcgp.org.uk/policy/general-practice-covid-19-power-of-relationships.aspx> 2021.
23. Rhodes P, Campbell S, Sanders C. Trust, temporality and systems: how do patients understand patient safety in primary care? A qualitative study. *Health Expect*. 2016;19(2):253-63.
24. Rhodes P, McDonald R, Campbell S, et al. Sensemaking and the co-production of safety: a qualitative study of primary medical care patients. *Sociol Health Illn*. 2016;38(2):270-85.
25. Rhodes P, Sanders C, Campbell S. Relationship continuity: when and why do primary care patients think it is safer? *Br J Gen Pract*. 2014;64(629):e758-e64.
26. Greenhalgh T, Collard A, Begum N. Sharing stories: complex intervention for diabetes education in minority ethnic groups who do not speak English. *BMJ*. 2005;330(7492):628.

27. Sanders C, Seymour J, Clarke A, et al. Development of a peer education programme for advance end-of-life care planning. *Int J Palliat Nurs*. 2006;12(5):216-23.
28. Groundswell. The Escape Plan: A participatory Research Study, London. 2008. https://groundswell.org.uk/wp-content/uploads/2018/10/Groundswell-The_Escape_Plan_Report-2008.pdf (access Sept, 2021)
29. Groundswell. Out of pain: breaking the cycle of physical pain and homelessness: London. 2018 <https://groundswell.org.uk/wp-content/uploads/2018/08/Groundswell-Out-of-Pain-Report-Final.pdf> (accessed Sept 2021).
30. De Oliveira B. On the news today: challenging homelessness through participatory action research. *Housing, Care and Support*. 2018. <https://www.emerald.com/insight/content/doi/10.1108/HCS-01-2018-0002/full/html> (access Sept 2021)
31. Franco A, Ngaruiya C. 275 Targeting Homeless Population Needs Using Community-Based Participatory Research Methods in an Urban Emergency Department. *Ann Emerg Med*. 2016;4(68):S106-S7.
32. Burns S, Schubotz D. Demonstrating the merits of the peer research process: A Northern Ireland case study. *Field Methods*. 2009;21(3):309-26.
33. Lushey CJ, Munro ER. Participatory peer research methodology: An effective method for obtaining young people's perspectives on transitions from care to adulthood? *Qual Soc Work*. 2015;14(4):522-37.
34. Terry L, Cardwell V. Refreshing perspectives: Exploring the application of peer research with populations facing severe and multiple disadvantage. London: Revolving Doors Agency & Lankelly Chase. 2016.
35. Beresford P. User involvement, research and health inequalities: developing new directions. *Health Soc Care Community*. 2007;15(4):306-12.
36. Clover D. Successes and challenges of feminist arts-based participatory methodologies with homeless/street-involved women in Victoria. *Action Research*. 2011;9(1):12-26.
37. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol*. 2013;13(1):1-8.
38. Ritchie J, Spencer L. Qualitative data analysis for applied policy research. *The qualitative researcher's companion*. 2002;573(2002):305-29.
39. Dixon-Woods M, Cavers D, Agarwal S, et al. Conducting a critical interpretive synthesis of the literature on access to healthcare by vulnerable groups. *BMC Med Res Methodol*. 2006;6(1):1-13.
40. Fine M, Torre ME. Re-membering exclusions: Participatory action research in public institutions. *Qual Res Psychol*. 2004;1(1):15-37.
41. Asgary R, Sckell B, Alcabes A, et al. Perceptions, attitudes, and experience regarding mHealth among homeless persons in New York City shelters. *J heal commun*. 2015;20(12):1473-80.
42. Donaghy E, Atherton H, Hammersley V, et al. Acceptability, benefits, and challenges of video consulting: a qualitative study in primary care. *Br J Gen Pract*. 2019;69(686):e586-e94.
43. Ignatowicz A, Atherton H, Bernstein CJ, et al. Internet videoconferencing for patient-clinician consultations in long-term conditions: A review of reviews and applications in line with guidelines and recommendations. *Digit health*. 2019;5:2055207619845831.
44. Murphy M, Scott LJ, Salisbury C, et al. Implementation of remote consulting in UK primary care following the COVID-19 pandemic: a mixed-methods longitudinal study. *Br J Gen Pract*. 2021;71(704):e166-e77.
45. Checkland K. Who needs to see a doctor face to face? 35th Annual HSR Lecture. 2021.
46. England N. Personalised care: An induction guide for care-coordinator workers in primary care networks <https://www.england.nhs.uk/wp-content/uploads/2021/05/care-coordinator-welcome-pack.pdf>. 2021.

47. Huang H, Jefferson E, Gotink M, et al. Collaborative improvement in Scottish GP clusters after the Quality and Outcomes Framework: a qualitative study. *Br J Gen Pract*. 2021.

Accepted Manuscript – BJGP – BJGP.2021.0596

Table 1: Description of case study sites

Case study sites	Service Pre COVID-19	Main changes during COVID-19 pandemic
CSS1- primary care	<ul style="list-style-type: none"> • Flexible drop in appointments • Outreach and hostel drop-ins • Hospital in reach service • Case management of frequent attenders 	<ul style="list-style-type: none"> • No drop in clinics except for wound care • Telephone led consultations • Outreach was initially stopped and reintroduced in November 2020
CSS2- primary care	<ul style="list-style-type: none"> • GP service offering flexible appointments and drop in service • Outreach at daycentres 	<ul style="list-style-type: none"> • No drop in clinics • Outreach was initially stopped but resumed in November 2020 • Dedicated care coordinator to schedule appointments • Video- consultations trialled
CSS3- NHS community nurse inclusion service	<ul style="list-style-type: none"> • Nurse led drop in clinics • Outreach in the community, Hostels/guest houses/town centre walk about • Weekly street kitchen 	<ul style="list-style-type: none"> • Outreach to hostels and town centre walk about initially stopped in the national 1st lockdown from end of March-mid June 2020. Street Kitchen stopped for the same period. • Nurse led 'face- to face' clinics continued and outreach to guest houses continued on a needs basis.

Table 2: **Table 2:** Demographic characteristics of people experiencing homelessness sample (n=21)

<i>Sex</i>	
Male	18
Female	3
<i>Age</i>	
25-34	5
35-44	6
45-54	8
55-64	2
<i>Ethnicity</i>	
White British	18
Mixed Race	2
Eastern European	1
<i>Qualifications</i>	
None	11
GCSE	8
Degree	1
Missing	1
<i>Number of chronic conditions</i>	
1	3
2	2
More than 3	16
Anxiety and depression	15
Alcohol and drug related issues	11