

Patient involvement in diagnosing cancer in primary care:

a systematic review of current interventions

Abstract

Background

Patients can play a role in achieving an earlier diagnosis of cancer by monitoring and re-appraising symptoms after initially presenting to primary care. It is not clear what interventions exist, or what the components of an intervention to engage patients at this diagnostic stage are.

Aim

The review had two aims: 1) to identify interventions that involve patients, and 2) to establish key components for engaging patients in the diagnosis of cancer in primary care at the post-presentation stage.

Design and setting

Empirical studies and non-empirical articles were identified searching Ovid MEDLINE, PsycINFO, and Embase databases, relevant journals, and available key author publication lists.

Method

Abstracts and titles were screened against inclusion and exclusion criteria. Qualitative synthesis of empirical research and current opinion from across all articles was used to select, organise, and interpret findings.

Results

No interventions were found. Sixteen articles provided suggestions for potential interventions and components important at the post-presentation stage. Factors contributing to patients not always being engaged in assisting with diagnosis, strategies to foster patient involvement, and moderators and benefits to patients and health services (proximal and distal outcomes) were captured in a logic model.

Conclusion

There is an absence of interventions involving patients during the post-presentation stage of the diagnostic process. Limited literature was drawn upon to identify potential barriers and facilitators for engaging patients at this diagnostic stage, and to establish possible mechanisms of change and measurable outcomes. Findings can direct future research and the development of interventions.

Keywords

cancer; diagnosis; patient involvement; primary health care; systematic review.

INTRODUCTION

It is estimated that 46% of all cancers in England are diagnosed at a late stage, resulting in lower survival and additional treatment costs.¹ Delays in cancer diagnosis may occur at a number of stages during the diagnostic process.²⁻⁴ In a primary care setting, the patient interval (before presentation) rather than the primary care interval (after presentation) has been found to make a larger contribution to delay in 18 out of 28 cancers assessed.⁵ Work on factors involved in emergency presentations of cancer has also examined patients' previous use of primary care.^{6,7} One finding was that some patients diagnosed with cancer during emergency presentations had delayed re-presenting back to primary care after an initial consultation for symptoms that might be related to their cancer symptoms (referred to as post-presentation stage from this point).⁶ Delayed re-presentation may contribute to a diagnosis at a point where treatment is less effective. Clear communication between the GP and patient about the possible meaning of symptom severity and persistence, and the patient's role in following up and reviewing symptoms, are critical for early recognition, referral, and treatment for cancer.^{8,9}

Though diagnostic error is little researched in the field of patient safety,^{10,11} interest in the role that patients may

have in reducing delayed diagnosis and misdiagnosis, and improving health professionals' diagnostic performance, has begun to gather pace. This is reflected in the recent Institute of Medicine¹² report on diagnostic error, which proposes that patients are 'a diagnostic team member'. The importance of patients and healthcare professionals working in partnership to avoid delay in the diagnosis of cancer is also an emerging topic and a key component in the increasing focus on safety netting.^{8,13} Although there is a growing appreciation that engaging patients in their own health care can achieve better outcomes through involvement in disease management^{14,15} and improving patient safety,¹⁶ little is known about how patients currently assist with cancer diagnosis in primary care.

Here, the authors report a systematic review that addresses this gap in understanding. The aims of the review were to consolidate existing knowledge on interventions that involve patients in the diagnosis of cancer in primary care at the post-presentation stage (Aim 1), and to identify the components necessary to engage patients to be actively involved in achieving an earlier diagnosis of cancer at the post-presentation stage (Aim 2). By developing a logic model of these components, the authors' ambition was to guide further research and intervention development in this field.

J Heyhoe, PhD, senior research fellow;
C Reynolds, BSc, research practitioner, Bradford Institute for Health Research, Bradford. **A Dunning**, BSc, PhD student, School of Psychology, University of Leeds, Leeds. **O Johnson**, BSc, formerly School of Psychology, University of Leeds, Leeds. **A Howat**, MSc, formerly Institute of Psychiatry, Psychology, and Neuroscience, King's College London, London. **R Lawton**, PhD, professor, Psychology of Healthcare, School of Psychology, University of Leeds, Leeds, and Bradford Institute for Health Research, Bradford.

Address for correspondence

Jane Heyhoe, Bradford Institute for Health Research, Bradford Royal Infirmary, Duckworth Lane, Bradford, BD9 6RJ, UK.

E-mail: Jane.Heyhoe@bthft.nhs.uk

Submitted: 22 June 2017; **Editor's response:** 31 July 2017; **final acceptance:** 2 November 2017.

©British Journal of General Practice

This is the full-length article (published online 13 Feb 2018) of an abridged version published in print. Cite this version as: **Br J Gen Pract 2018**; DOI: <https://doi.org/10.3399/bjgp18X695045>

How this fits in

Patients have a role to play in achieving an earlier diagnosis of cancer, but little is known about how patients can be engaged in monitoring and re-appraising symptoms after an initial presentation to primary care, and before a referral or a diagnosis is obtained. This is reflected in the absence of interventions that involve patients at this stage. Drawing upon current thinking and empirical data, this systematic review identifies potential barriers and facilitators to patient involvement at the post-presentation stage, possible mechanisms of change, and measurable outcomes. The findings have implications for research and practice in the area of patient involvement in achieving an earlier cancer diagnosis in primary care.


METHOD

This systematic review was carried out using the preferred reporting items in systematic reviews and meta-analyses (PRISMA) guidelines.¹⁷

Search strategy: Aim 1

Databases were selected to cover both medical and psychological literatures.

Figure 1. Flow diagram of search strategy for Aim 1.



One researcher, with medical librarian assistance, searched three electronic databases: Ovid MEDLINE (1946 to February 2016), PsycINFO (1806 to February 2016), and Embase (1974 to February 2016). Articles were not restricted by publication type or study design but were limited to the English language. Four blocks of search terms (Appendices 1 and 2) to capture diagnosis (using previously used search terms),^{18–20} patient involvement, primary health care, and cancer, were combined for the database search. The authors did not search using the specific term 'safety netting', a relatively new concept in the diagnosis literature. A recent search using this term found no safety-netting interventions.⁸ The authors used a broader set of terms in the hope of identifying interventions with characteristics of 'safety netting' but which did not make reference to this specific form of intervention. In addition, two researchers hand searched relevant journal articles from 2010 to February 2016 in relevant fields (Appendix 3). Two reviewers independently reviewed the titles and abstracts of retrieved citations, screening half of the abstracts each. Two researchers reviewed 10% of the citations. The full paper was obtained if the inclusion criteria were met, or no abstract was available. Articles were retained or excluded using inclusion and exclusion criteria for Aim 1 (Appendix 4). Together, two reviewers screened all full-text articles obtained (n = 76) to identify those that reported the evaluation of an intervention using any method at the post-presentation stage (Aim 1). At all stages, disagreements on whether to include or exclude an article were resolved through discussion and consensus.

Search strategy: Aim 2

To address Aim 2, two reviewers screened articles that met the original inclusion/exclusion criteria (n = 76). Articles were retained or excluded using inclusion and exclusion criteria for Aim 2 (Appendix 5). An additional secondary strategy involved two researchers manually searching the reference list of all included articles. Publication lists of key researchers in the field were also searched. The decision to include or exclude an article was made through discussion, and a third researcher arbitrated when consensus could not be reached. At this stage, five articles fulfilling the inclusion criteria were removed due to data duplication.^{21–25} One article⁶ used some of the data reported in another article.²⁶ However, as the data was analysed to answer different research questions the




Figure 2. Flow diagram of search strategy for Aim 2.

decision was made to include the article. Two articles could not be retrieved for full article screening.

Data extraction and synthesis: Aim 2

General characteristics data were extracted for each article/study. To identify potential strategies or key components for an intervention at the post-presentation stage, the authors conducted a qualitative synthesis of research findings and current opinion. This facilitated the selection, organisation, and interpretation of nuanced data and thinking from articles with different objectives and which employed a variety of methods. This involved four distinct stages. First, all text relevant to the inclusion criteria in each article was highlighted by one of two researchers and then double-coded by the other researcher. Second, the same two researchers discussed and reviewed the highlighted text and developed a data extraction table (available from the authors upon request) that most meaningfully captured the key categories for all highlighted text and together organised text from each article into each category. Third, the two researchers and one other interrogated the data extraction table and, drawing on a programme theory approach,^{27,28} began to

develop a logic model that illustrated all considerations, possible mechanisms, and outcomes suggested in the articles. Fourth, the logic model was refined to illustrate key considerations, possible mechanisms, and outcomes across the articles.

RESULTS

No published studies met the inclusion criteria for Aim 1 of the review; 16 articles met the inclusion criteria for Aim 2. PRISMA flow charts for Aim 1 and Aim 2 are detailed in Figure 1 and 2.

General characteristics of included articles

The general characteristics of the articles included in the systematic review are detailed in Table 1. Four articles were non-empirical,^{29–32} whereas the majority of articles ($n = 12$)^{6,26,33–42} reported empirical findings. Qualitative methods, including interviews ($n = 6$),^{34–36,39,41,42} focus groups ($n = 2$),^{40,42} and qualitative synthesis of significant event audits ($n = 2$)^{6,26} were used in nine studies. Quantitative methods were used in six studies,^{26,33,37,38,40,42} and included responses to vignettes ($n = 1$),³³ responses to a questionnaire ($n = 3$),^{37,38,42} statistical review of incident reports ($n = 1$),⁴⁰ and clinical audit ($n = 2$).^{26,42} Three articles employed mixed methods using both quantitative and qualitative data.^{26,40,42} The majority of articles reported studies that were conducted in the UK ($n = 9$),^{6,26,33–36,38–40} and one study each was conducted in Denmark,³⁷ the US,⁴¹ and New Zealand.⁴²

Though three of the non-empirical articles^{29–31} and the thematic review⁴⁰ did not focus on a specific type of cancer, the other articles varied in the type of cancer studied or discussed. Lung cancer was most frequently studied, featuring in eight articles,^{6,26,33–35,37,39,42} and was the sole focus in two of these articles.^{34,42} Gastrointestinal (colorectal, colon, rectal, upper GI) was studied in seven articles,^{6,26,33,35,37,39,41} in one of which it was the sole focus.⁴¹ Ovarian cancer was the sole focus of one article,³⁶ and cervical cancer was the sole focus of two articles.^{32,38} Six articles considered a number of different cancers.^{6,26,33,35,37,39}

The study population in articles varied. Patients only were involved in eight studies,^{33–36,38,39,41,42} six of which consisted of patients with a diagnosis of cancer^{35,36,38,39,41,42} and one which involved general patients.³³ Two articles included both patients with a diagnosis of cancer and patients suspected of cancer but who had been diagnosed with other conditions.^{34,42} GPs only were involved in three studies,^{6,26,37}

Table 1. General characteristics of articles included in systematic review

Author/year	Type of cancer	Country	Article type	Aim — text taken from paper	Study design	Setting	Population characteristics
Banks <i>et al.</i> , 2014 ³³	Lung, pancreatic, colorectal	England	Research	To establish the likelihood that individuals would choose to be tested for cancer at various levels of risk	Vignette-based study	26 general practices across England	3469 individuals (≥40 years)
Birt <i>et al.</i> , 2014 ³⁴	Lung	England	Research	To understand the symptom evaluation, or 'appraisal', and help-seeking decisions of patients with symptoms suggestive of lung cancer	Prospective cohort face-to-face interview study	Specialist respiratory clinics in secondary and tertiary care, England	35 patients (male = 20, female = 15; mean age 58.5 years; range 41–88 years; 17 with lung cancer, 18 with other conditions)
Black <i>et al.</i> , 2015 ³⁵	Colorectal, upper GI, lung, haematological, head and neck, gynaecological, unknown primary	England	Research	To capture in detail the experiences of patients whose cancer was diagnosed following an ED visit to understand how emergency presentations arise, and identify where there is scope to improve outcomes	Interview	11 EDs across nine acute NHS trusts (participants came from seven NHS trusts)	27 patients (male = 15, female = 12; median age 59 years; range 18–92 years)
Evans <i>et al.</i> , 2006 ³⁶	Ovarian	UK	Research	1) To use accounts of provider delay from a sample of British women with ovarian cancer to enhance Andersen's model of total patient delay; 2) To suggest what GPs might do to minimise delays	Semi-structured interviews with a maximum variation sample	Recruitment through GPs, clinicians, support organisations, and personal contacts	43 women diagnosed with ovarian cancer (mean age 54 years; range 33–80 years)
Jensen <i>et al.</i> , 2014 ³⁷	Bladder, breast, colon, lung, malignant melanoma, others, ovarian, prostate, rectal, uterus	Denmark	Research	1) To describe the prevalence and types of QDs that arose during the diagnostic pathway for Danish patients with cancer as assessed by the GPs; 2) To analyse the associations between QDs and the type of cancer; between QDs and the GP interpretation of presenting symptoms; 3) Impact of QDs on diagnostic interval length	Nationwide retrospective cohort questionnaire-based study	1446 GPs identified through Danish national patient registry	4036 newly diagnosed cancer patients (male = 2176, female = 1860)
Lim <i>et al.</i> , 2014 ³⁸	Cervical	England	Research	To collect data on nature and duration of symptoms and risk factors for delay in presentation and diagnosis in young females with cervical cancer. To inform approaches to promote early presentation and prompt referral among young females with symptoms of cervical cancer	Semi-structured interview-based measure	Eligible participants identified by 116 hospitals and publicising of study through charity websites	128 women (18–29 years) with recent diagnosis of cervical cancer
Lyrtzopoulos <i>et al.</i> , 2015 ³⁹	Not specified	N/A	Perspective	1) To highlight factors involved in missed opportunities for cancer diagnosis among symptomatic patients; 2) Discuss potential mechanisms and approaches to accelerating progress towards minimising diagnostic delays post-presentation	N/A	All settings involved in diagnostic process including general practice and primary care	N/A

... continued

Table 1 continued. General characteristics of articles included in systematic review

Mitchell <i>et al</i> , 2012 ²⁶	Lung, ovarian, upper GI	England	Toolkit	Developed as a resource for use when considering ways to improve the diagnosis of cancer in primary care	1) National audit of cancer diagnosis in primary care; 2) Analysis of SEA; 3) Action plans for early cancer diagnosis derived using cancer-related practice profiles	Primary care	1) 1170 practices across 20 cancer networks; 2) SEA reports for 132 lung, 68 ovarian, and 78 upper GI cancer diagnoses; 3) general practices across England
Mitchell <i>et al</i> , 2015 ⁶	Brain, breast, carcinoid, CNS, colorectal, gynaecological, haematological, head and neck, lung, melanoma, sarcoma, upper GI, urological, unknown primary	England	Research	1) Understanding the cause of emergency presentations; 2) Determining the degree to which earlier intervention by general practice was possible	Qualitative synthesis of SEAs	203 primary care practices across North of England Cancer Network	222 SEAs for most recent patients diagnosed with cancer as a result of an emergency presentation
Molassiotis <i>et al</i> , 2010 ³⁹	Breast, GI, gynaecological, lung, brain, lymphoma, head and neck	UK	Research	1) To map the pathway from initial persistent change in health to diagnosis of cancer in a sample of patients from seven diagnostic groups in the UK; 2) Explore the patient and system factors mediating this process	In-depth qualitative interviews	Outpatient clinics at one large cancer hospital	74 patients (mean age 58.5 years; range 18–93 years)
National Reporting and Learning Service, 2010 ⁴⁰	All cancers	UK	Thematic review	1) To explore issues of patient safety around delayed diagnosis of cancer; 2) Provide the NHS with potential solutions	1) Thematic review; 2) Review of incidents reported to the NPSAs NRLS; 3) Consultation with stakeholders (focus groups, discussions, presentations); 4) Consideration of other patient safety data, such as complaints, litigation, and audits	Primary and secondary care	1) N/A; 2) 508 patient safety incidents from random sample of 1500; 3) Two focus groups (n = 50); 4) NHS Litigation Authority; MDU Report (2003); MPS (analysis of 1000 cases of delayed or missed diagnosis); Scottish Primary Care Cancer Group reports of 4181 cancer cases
Round <i>et al</i> , 2013 ³⁰	Not specified	UK	Essay	To highlight clinician- and system-related factors that may be contributing to a delayed diagnosis of cancer in primary care	N/A	Primary care	N/A

... continued

Table 1 continued. General characteristics of articles included in systematic review

Rubin <i>et al</i> , 2015 ³¹	Not specified	High-income countries/ countries with universal healthcare systems	Commission	To distil the evidence for the effectiveness of interventions for cancer control based in primary care at each stage of the cancer journey, and to consider how cancer care might be delivered differently in the future. Discuss how and whether health policy for cancer control will help or hinder such change. Examines the implications for the future education and training of doctors, and identifies emerging examples of good practice worldwide	N/A	Primary and specialist care	N/A
Siminoff <i>et al</i> , 2011 ⁴¹	Colorectal	US	Research	1) To describe the communication and symptom appraisal strategies of CRC patients; 2) Determine the patient and physician characteristics, consultation communication factors, follow-up recommendations, and the referral delay factors associated with DD	Semi-structured interview and review of medical records for verification of patient-provided information	Five academic and community oncology practices in two states	242 patients of any age (mean = 58.07 years) or sex (male = 126, female = 116)
Stevens <i>et al</i> , 2012 ⁴²	Lung	New Zealand	Report	To develop recommendations to improve the clinical pathway from presentation to diagnosis for people with suspected lung cancer	Mixed methods utilising both quantitative and qualitative data: 1) Literature review; 2) National Stocktake; 3) Clinical audit; 4) Practice survey; 5) GP survey; 6) GP focus group; 7) Patient interviews; 8) Patient and family focus groups	Four primary care organisations and three DHBs within the Northern and Midland Cancer Networks	1) N/A; 2) Primary or secondary care services that could be relevant to people with lung cancer; 3) Clinical pathway from initial presentation to diagnosis (272 cases); 4) All general practices across two regions (<i>n</i> = 170); 5) 102 GPs across two regions; 6) 33 GPs across two regions; 7) 19 people suspected of having lung cancer who presented to a hospital ED; 8) 20 people (11 patients with lung cancer diagnosis [male = 5; female = 6] and nine family members)
Walter <i>et al</i> , 2014 ²²	Cervical	UK and international	Editorial	Not specified	N/A	Primary care	N/A

CNS = central nervous system. CRC = colorectal cancer. DD = diagnostic delay. DHB = district health board. ED = emergency department. GI = gastrointestinal. MDU = Medical Defence Union. MPS = Medical Protection Society. N/A = not applicable. NPSA = National Patient Safety Agency. NPLS = National Reporting and Learning System. QD = quality deviation. SEA = significant event audit.

all of which included patients with a diagnosis of cancer. Both patients, GPs, and/or primary care stakeholders were included in two studies.^{40,42} And in one article, though patients with cancer and primary care stakeholders were included, it was difficult to establish the exact study population.⁴⁰

Suggestions for potential interventions/strategies

Though none of the 16 articles directly set out to answer questions about how patients could be involved in the diagnosis of cancer at the post-presentation stage, all articles did consider the patient's role in diagnostic delay, and identified how they may be involved in the diagnostic process. Articles also considered the contribution of other factors attributable to the healthcare provider (HCP) or system issues. Other than one article that explored referral preferences,³³ all articles incorporated findings across a number of diagnostic stages, including pre-initial presentation and/or post-referral factors. As the focus of this review is the post-presentation stage, findings and suggestions that were solely related to other diagnostic stages were disregarded.

Key considerations, possible mechanisms, and outcomes across the articles

The logic model (Figure 3) presents key components and potential mechanisms for involving patients in achieving an earlier diagnosis of cancer and other desirable outcomes.

'Patients' and healthcare professionals' difficulty in identifying potential symptoms' was identified across most articles, and included distinguishing between symptoms associated with comorbid conditions and new symptoms that may be due to cancer,^{6,26,29-31,34,36,38-42} and the significance of vague or non-specific symptoms, as well as established 'red flag' symptoms.^{6,26,29-32,34-42} Issues around 'poor patient-healthcare professional communication during consultation' were also prominent. These included the healthcare professionals not eliciting a thorough history or asking relevant questions,^{26,29,36,37,40,41} or the patient's ability to communicate their health problem.^{26,29,36,40,41} The emotions of patients were also critical: patients may not re-present due to feeling they were 'overburdening doctors' or wasting their time,^{29,31,38,39,41} or could be influenced by worries and fear concerning tests and potential diagnosis.^{26,29,31,33,36,38,41,42} The emotional response towards healthcare

professionals was also a factor when patients believed that their symptoms and concerns were not being taken seriously or misattributed.^{31,34-36,39,40}

There was a variety of suggestions for strategies to foster patient involvement in the post-presentation stage. Most common was the view that healthcare professionals would benefit from further improvement to their clinical practice. This was primarily around information gathering for diagnosis, such as history taking, physical examination, and record keeping.^{6,26,29-32,36,37,39} Many articles advocated interventions to involve the patient. The importance of 'safety netting' at the end of a consultation was a strong message. It was considered that patients and healthcare professionals agreeing together on a clear symptom follow-up and re-appraisal plan was important.^{6,26,29-32,34-38,40-42} Some articles discussed having more of an 'open door' approach,^{29,37} and a number also highlighted the importance of setting a timescale for review.^{6,26,29,30,32,34-36}

Many articles also suggested ways for facilitating more open dialogue between the patient and healthcare professional. This included communication about the potential meaning of symptoms and/or the patient's personal risk^{31-33,39-41} and healthcare professionals sharing with patients the reasoning behind diagnostic decisions.^{26,36,37,41} Some articles also stressed the importance of having procedures to ensure continuity of diagnostic information. This involved the patient seeing the same healthcare professional,^{6,26,30} and improved record keeping to facilitate the linking of previous presentations and/or access to management plans and potential diagnoses, should a patient present to a different healthcare professional.^{6,26,30,31}

Potential moderators of the relationship between suggested strategies and the benefits to patients and health services were identified at three distinct levels. At the patient level, characteristics such as the extent to which patients adhered to the recommendations of the healthcare professional (for example, choosing or not to re-attend to review symptoms)^{6,26,29,31,37,39,41,42} or patients' communication skills, such as their ability to express their health problems and concerns,^{26,29-31,36,41,42} were seen to have an impact on engagement and outcomes. Characteristics of the healthcare professional were also viewed as influencing patient engagement and outcomes, and included the healthcare professional's experience of cancer to aid recognition.^{29,31,36,39} There were also a




Figure 3. Logic model — patient involvement in diagnosing cancer in primary care. HCP = healthcare professional.

number of moderators at the practice level, which included the impact of consultation length on ability to adequately discuss and address health concerns.^{29,30,36,42}

Both proximal and distal benefits to patients and health services emerged across the articles. Proximal outcomes resulting from proposed strategies include improved

monitoring of symptoms^{6,26,29–32,34–41} and improved quality of the patient–healthcare professional consultation.^{6,26,29–31,33,35–37,39–41} It is proposed that proximal outcomes such as these may serve as mediators between proposed strategies and distal outcomes, which include reduced diagnostic delay.^{26,29,32,34–38,40–42}

DISCUSSION

Summary

This systematic review did not identify any interventions that involve patients in achieving an earlier diagnosis of cancer at the post-presentation stage and, indeed, none of the articles set out to do this. In general, articles tended to identify and describe barriers and facilitators to achieving an early diagnosis of cancer in primary care, and the role that patients play in diagnostic delay, and provided some general ideas about ways in which patients can assist in this process.

At present, it is unclear whether healthcare professionals and patients sharing responsibility in reviewing symptoms is effective,⁸ and there is currently no consensus on the practices involved in safety netting,^{8,13,43} a strategy that is likely to require sustained patient engagement if it is to influence diagnostic outcomes. Based on the authors' reading of empirical data and current thinking drawn from 16 articles, they identified common components and mechanisms considered to be key to involve patients in diagnosis at the post-presentation stage. This has enabled them to develop and illustrate a first attempt at a logic model for patient involvement in diagnosis in primary care.

Strengths and limitations

There are limitations to this review. Though the authors' focus was on re-presentation following an initial presentation to primary care, they did not identify any article that focused solely on this aspect in the diagnostic process. To retrieve the information required, the authors had to include articles that incorporated and gathered data from a number of stages in the diagnostic process, and varied in the way diagnostic stages were distinguished. A number of articles also make very general suggestions about patient involvement without expanding on these ideas, or providing any detail concerning what any strategy might include or require. Due to this, gathering data to answer the research questions was a difficult process, and it is possible that information not specific to the diagnostic stage of interest was included. To mitigate this, the authors applied a rigorous and consistent approach to the selection and collation of data using cross-checking and consensus opinion. They also drew upon a variety of articles using different methods to synthesise data to represent an overview of current thinking and findings.

In addition, articles or sections of articles that focused on patients'

symptom perception and help-seeking behaviour were excluded because of these components traditionally being regarded as contributing towards the delay in patients' initial primary care consultation. One of the articles included in this review³⁹ suggests that symptom perception and help-seeking behaviour before the first presentation may re-emerge and influence decisions and behaviour when a patient re-evaluates their symptoms following an initial presentation. As it is feasible that components identified in this review may operate in a continuum across diagnostic stages (pre-consultation and consultation, as well as post-consultation), it may transpire that findings from work with a focus on the pre-presentation stage, as well as the primary care consultation, are relevant for the development of interventions once the patient has presented. More work in this area is required to establish whether this is the case.

Comparison with existing literature

Though a previous systematic review of patient safety strategies targeted at diagnostic error²⁰ found two studies that focused on patient education, the studies were for breast screening and mothers' symptom decisions for their child, both of which would not have fitted the criteria for this review. This indicates that there is currently a gap in the evidence base in this field. It also reflects a pattern of a lack of development and evaluation of evidence-based interventions in the field of diagnosis found previously in literature searches conducted for system-related,¹⁸ cognitive,¹⁹ and safety-netting interventions.⁸

Implications for research and practice

The findings from this review can be used to guide the direction of future research and intervention development. Though a limited number of qualitative studies assessing the feasibility of safety-netting strategies have been identified and are an important step in understanding how safety netting could be operationalised in practice, there is still no evidence of intervention development in the context of cancer.⁴⁴ Though this review found some evidence to suggest that 'active' (establishing a set time for a further appointment) rather than 'passive' (the patient making a further appointment if they feel it is required) safety netting was perceived to be important to ensure the patient returns, further work is required to establish whether one approach rather than the other fosters patient involvement at post-presentation.^{8,43}

In addition, though it is generally agreed that an important dimension in achieving better diagnostic outcomes is engaging patients in the diagnostic process, it is unclear whether patients will welcome such an initiative or what components are important to encourage and sustain patient involvement.^{45,46} In light of this, this review is timely and progresses the field by identifying barriers and facilitators, possible mechanisms of action, and measurable outcomes when considering patient involvement in achieving an earlier cancer

diagnosis in a primary care setting across published articles.

The logic model presented in this review may help to bridge a gap between qualitative findings and quantitative potential through facilitating the testing of hypotheses, and the development and evaluation of evidence-based interventions that are currently lacking in the field of diagnosis.^{18,19} It also provides an initial testable model that can now be refined through further research.

Funding

This paper presents research funded by Yorkshire and Humber Commissioning Support Research Capability Funding (RCF) (ref. RCF-2015-011) and the National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care Yorkshire and Humber (NIHR CLAHRC YH) (ref. 2016-17) (<http://clahrc-yh.nihr.ac.uk/>). The views expressed are those of the authors, and not necessarily those of the NHS, the NIHR, or the Department of Health.

Ethical approval

Not applicable.

Provenance

Freely submitted; externally peer reviewed.

Competing interests

The authors have declared no competing interests.

Acknowledgements

The authors would like to thank reviewers for their comments on an earlier draft of this paper.

Discuss this article

Contribute and read comments about this article: bjgp.org/letters

REFERENCES

1. Incisive Health for Cancer Research UK. *Saving lives, averting costs. An analysis of the financial implications of achieving earlier diagnosis of colorectal, lung, and ovarian cancer*. A report prepared for Cancer Research UK. CRUK, 2014.
2. Anderson BL, Cacioppo JT, Roberts DC. Delay in seeking a cancer diagnosis: delay stages and psychophysiological comparison processes. *Br J Soc Psychol* 1995; **34**(Pt 1): 33–52.
3. Hansen RP, Vedsted P, Sokolowski I, *et al*. General practitioner characteristics and delay in cancer diagnosis. A population-based cohort study. *BMC Fam Pract* 2011; **12**: 100. <https://doi.org/10.1186/1471-2296-12-100>.
4. Hiom SC. Diagnosing cancer earlier: reviewing the evidence for improving cancer survival. The National Awareness and Early Diagnosis Initiative in England: assessing the evidence 5 years on. *Br J Cancer* 2015; **112**(Suppl 1): S1–S5.
5. Lyratzopoulos G, Saunders CL, Abel GA, *et al*. The relative length of the patient and the primary care interval in patients with 28 common and rarer cancers. *Br J Cancer* 2015; DOI: 10.1038/bjc.2015.40.
6. Mitchell ED, Rubin G, Merriman L, Macleod U. The role of primary care in cancer diagnosis via emergency presentation: qualitative synthesis of significant event reports. *Br J Cancer* 2015; **112**: S50–S56.
7. Zhou Y, Abel GA, Hamilton W, *et al*. Diagnosis of cancer as an emergency: a critical review of current evidence. *Nat Rev Clin Oncol* 2017; DOI: 10.1038/nrclinonc.2016.155.
8. Nicholson BD, Mant D, Bankhead C. Can safety-netting improve cancer detection in patients with vague symptoms? *BMJ* 2016; **355**: i5515. DOI: 10.1136/bmj.i5515.
9. Cancer Research UK. *Early diagnosis of cancer. How do we make sure patients don't slip through the net?* CRUK, 2016. http://www.cancerresearchuk.org/sites/default/files/safety_netting_scotland_201607.pdf [accessed 11 Apr 2017].
10. Singh H. Diagnostic errors: moving beyond 'no respect' and getting ready for prime time. *BMJ Qual Saf* 2013; **22**(10): 789–792.
11. Wachter RM. Diagnostic errors: central to patient safety, yet still in the periphery of safety's radar screen. *Diagnosis* 2014; **1**(1): 19–21.
12. Institute of Medicine. *Improving diagnosis in health care*. IOM, 2015. <http://nas.edu/improvingdiagnosis> [accessed 17 Jan 2018].
13. Edwards PJ, Seddon JO, Barnes RK. Safety-netting: a time for guidelines? *BMJ* 2016; <http://www.bmj.com/content/355/bmj.i5515/rr> [accessed 17 Jan 2018].
14. Coulter A. Patient engagement — what works? *J Ambul Care Manage* 2012; **35**(2): 80–89.
15. NHS England. *Five year forward view*. NHSE, 2014.
16. Lawton R, O'Hara J, Sheard L, *et al*. Can staff and patient perspectives on hospital safety predict harm-free care? An analysis of staff and patient survey data and routinely collected outcomes. *BMJ Qual Saf* 2015; **24**(6): 369–376.
17. Moher D, Liberati A, Tetzlaff J, Altman DG, the PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**(7): e1000097.
18. Singh H, Graber ML, Kissam SM, *et al*. System-related interventions to reduce diagnostic errors: a narrative review. *BMJ Qual Saf* 2011; DOI: 10.1136/bmjqs-2011-000150.
19. Graber ML, Kissam S, Payne VL, *et al*. Cognitive interventions to reduce diagnostic error: a narrative review. *BMJ Qual Saf* 2012; DOI: 10.1136/bmjqs-2011-000149.
20. McDonald K, Matesic B, Contopoulos-Ioannidis DG, *et al*. Patient safety strategies targeted at diagnostic errors. A systematic review. *Ann Intern Med* 2013; **158**(5 pt 2): 381–390.
21. Mitchell ED, Rubin G, Macleod U. Understanding diagnosis of lung cancer in primary care: qualitative synthesis of significant event audit reports. *Br J Gen Pract* 2013; DOI: <https://doi.org/10.3399/bjgp13X660760>.
22. Mitchell E, Macleod U, Rubin G. *Cancer in primary care. Significant event analysis of cancer diagnosis I: Lung and teenager/young adult cancers. Report to the National Awareness and Early Diagnosis Initiative*. Dundee: Department of Health, NAEDI, 2009.
23. Northern Cancer Network. *Literature review for the project 'Identification of barriers to the early diagnosis of people with lung cancer within primary care and description of best practice solutions'*. Northern Cancer Network, 2009. <http://www.northerncancernetwork.org.nz/LinkClick.aspx?fileticket=ZVHv0FU1VX4%3D&tabid=149&language=en-NZ> [accessed 17 Jan 2018].
24. Stevens W, Murray M, Hulme R, McNeill R. Expert Advisory Group. *GP survey report. Identification of barriers to the early diagnosis of people with lung cancer and description of best practice solutions*. Northern Cancer Network, 2011. <http://www.northerncancernetwork.org.nz/LinkClick.aspx?fileticket=rq3P07FXoMI%3d&tabid=117&language=en-NZ> [accessed 17 Jan 2018].
25. Walton L, McNeill R, Stevens W, *et al*. Patient perceptions of barriers to the early diagnosis of lung cancer and advice for health service improvement. *Fam Pract* 2013; **30**(4): 436–444.
26. Mitchell E, Rubin G, Macleod U. *Improving diagnosis of cancer: A toolkit for general practice*. London: Royal College of General Practitioners, 2012.
27. Lipsey MW. Theory as method: small theories of treatments. *New Dir Programme Eval* 1993; **57**: 5–38.
28. Goeschel CA, Weiss WM, Pronovost PJ. Using a logic model to design and evaluate quality and patient safety improvement programmes. *Int J Qual Health Care* 2012; **24**(4): 330–337.
29. Lyratzopoulos G, Vedsted P, Singh H. Understanding missed opportunities for more timely diagnosis of cancer in symptomatic patients after presentation. *Br J Cancer* 2015; **112**(Suppl 1): S84–S91.
30. Round T, Steed L, Shankleman J, *et al*. Primary care delays in diagnosing cancer: what is causing them and what can we do about them? *J R Soc Med* 2013; DOI: 10.1177/0141076813504744.
31. Rubin G, Berendsen A, Crawford SM, *et al*. The expanding role of primary care in cancer control. *Lancet Oncol* 2015; **16**(12): 1231–1272.
32. Walter FM, Mwaka AD, Neal RD. Achieving earlier diagnosis of symptomatic cervical cancer. *Br J Gen Pract* 2014; <https://doi.org/10.3399/bjgp14X681649>.
33. Banks J, Hollinghurst S, Bigwood L, *et al*. Preferences for cancer investigation: a vignette-based study of primary-care attendees. *Lancet Oncol* 2014; **15**(2): 232–240.
34. Birt L, Hall N, Emery J, *et al*. Responding to symptoms suggestive of lung cancer: a qualitative interview study. *BMJ Open Respir Res* 2014; **1**(1): e000067.
35. Black G, Sheringham J, Spencer-Hughes V, *et al*. Patients' experiences of cancer diagnosis as a result of an emergency presentation: a qualitative study. *PLoS One* 2015; DOI: 10.1371/journal.pone.0135027.
36. Evans J, Zieband S, McPherson A. Minimizing delays in ovarian cancer diagnosis: an expansion of Andersen's model of 'total patient delay'. *Fam Pract* 2006; DOI: 10.1093/fampra/cml063.
37. Jensen H, Nissen A, Vedsted P. Quality deviations in cancer diagnosis: prevalence and time to diagnosis in general practice. *Br J Gen Pract* 2014; <https://doi.org/10.3399/bjgp14X677149>.
38. Lim AW, Ramirez AJ, Hamilton W, *et al*. Delays in diagnosis of young females with symptomatic cervical cancer in England: an interview-based study. *Br J Gen Pract* 2014; <https://doi.org/10.3399/bjgp14X681757>.
39. Molassiotis A, Wilson B, Brunton L, Chandler C. Mapping patients' experiences from initial change in health to cancer diagnosis: a qualitative exploration of patient and system factors mediating this process. *Eur J Cancer Care* 2010; **19**(1): 98–109.
40. National Patient Safety Agency, National Reporting and Learning Service. *Delayed diagnosis of cancer: thematic review*. NPSA, 2010. <http://www.nrls.npsa.nhs.uk/EasySiteWeb/getresource.axd?AssetID=69895&> [accessed 17 Jan 2018].
41. Siminoff LA, Rogers HL, Thomson MD, *et al*. Doctor, what's wrong with me? Factors that delay the diagnosis of colorectal cancer. *Patient Educ Couns* 2011; **84**(3): 352–358.
42. Stevens W. *Recommendations to expedite the diagnosis of lung cancer. Final report of the HRC_DHBNZ funded project: 'Identification of barriers to the early diagnosis of people with lung cancer and description of best practice solutions'*. Northern Cancer Network, 2012.
43. Almond S, Mant D, Thompson M. Diagnostic safety-netting. *Br J Gen Pract* 2009; <https://doi.org/10.3399/bjgp09X472971>.
44. Nicholson BD, Bhuiya A, Black G, *et al*. Author response to Edwards *et al*. 'Safety-netting: a time for guidelines?' *BMJ* 2016; <http://www.bmj.com/content/355/bmj.i5515/rr-0>.
45. Graber ML. The IOM report on improving diagnosis: new concepts. *Diagnosis* 2015; **2**(4): 201–203.
46. McDonald KM, Bryce CL, Graber ML. The patient is in: patient involvement strategies for diagnostic error mitigation. *BMJ Qual Saf* 2013; DOI: 10.1136/bmjqs-2012-001623.

Appendix 1. MEDLINE and Embase search terms

- 1 Diagnosis/
- 2 diagn*.mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 3 Affect/
- 4 Clinical competence/
- 5 Communication/
- 6 "Continuity of Patient Care"/
- 7 Decision Making/
- 8 Decision Making, Organizational
- 9 Decision Support Systems, Clinical/
- 10 Decision Support Techniques/
- 11 Human Engineering/
- 12 Judgment/
- 13 Medical Informatics/
- 14 Medical Records Systems, Computerized/
- 15 Mental Recall/
- 16 Organizational Culture/
- 17 Patient Access to Records/
- 18 Feedback/
- 19 "Forms and Records Control"/st [Standards]
- 20 Guidelines as Topic/
- 21 Health Knowledge, Attitudes, Practice/
- 22 Health Literacy/
- 23 Health Records, Personal/
- 24 Physician's Practice Patterns/
- 25 Problem Solving/
- 26 Professional-Patient Relations/
- 27 Reminder Systems/
- 28 Systems Analysis/
- 29 Time Factors/
- 30 Truth Disclosure/
- 31 Knowledge Bases/
- 32 (cognitive error or bias or metacognition).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 33 Patient Participation/
- 34 (patient adj2 involv*).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 35 involv*.mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 36 patient empowerment.mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 37 (((patient adj2 led) or patient) adj2 instigated).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 38 Primary Health Care/
- 39 Family Practice/
- 40 exp Community Health Services/
- 41 Partnership Practice/
- 42 Private Practice/
- 43 (medical adj2 office).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 44 Neoplasms/
- 45 (([cancer or neoplasm*] adj5 (diagnos* or detect*])).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 46 (neoplasm* or cancer* or carcinoma* or malignan* or lesion* or tumo?*r*).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 47 (((family adj2 practice*) or (general adj2 practice*) or (primary adj2 care) or community) adj2 (service* or care)).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 48 ((((((delay* adj2 diagnos*) or diagnos*) adj2 delay*) or diagnos*) adj2 error*) or misdiagnos*).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 49 (missed adj2 diagnos*).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 50 (patient adj2 engagement).mp. [mp=ti, ab, hw, tn, ot, dm, mf, dv, kw, nm, kf, px, rx, ui]
- 51 1 or 2 or 48 or 49
- 52 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 or 35 or 36 or 37 or 50
- 53 38 or 39 or 40 or 41 or 42 or 43
- 54 44 or 45 or 46
- 55 51 and 52 and 53 and 54
- 56 limit 55 to english language
- 57 remove duplicates from 56

Appendix 2. PsycINFO search terms

- 1 Diagnosis/
- 2 (((((((diagn* or delay*) adj2 diagnos*) or diagnos*) adj2 delay*) or diagnos*) adj2 error*) or misdiagnos* or missed) adj2 diagnos*).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
- 3 Emotions/
- 4 clinical competence.mp.
- 5 Communication/
- 6 "continuity of patient care".mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
- 7 Decision Making/
- 8 decision making, organizational.mp.
- 9 decision support systems, clinical.mp.
- 10 Decision Support Systems/ or decision support techniques.mp.
- 11 human engineering.mp.
- 12 Judgment/
- 13 medical informatics.mp.
- 14 medical records, computerised.mp.
- 15 mental recall.mp.
- 16 Organizational Climate/
- 17 patient access to records.mp.
- 18 Feedback/
- 19 (forms and records control).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
- 20 guidelines as topic.mp.
- 21 Health Attitudes/ or Health Knowledge/ or health knowledge, attitudes, practice.mp.
- 22 Health Literacy/
- 23 health records, personal.mp.
- 24 physician's practice patterns.mp.
- 25 Problem Solving/
- 26 professional-patient relations.mp.
- 27 reminder systems.mp.
- 28 Systems Analysis/
- 29 time factors.mp.
- 30 truth disclosure.mp.
- 31 knowledge bases.mp.
- 32 (cognitive error or bias or metacognition).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
- 33 (((((((patient adj2 involv*) or involv* or patient empowerment or patient) adj2 led) or patient) adj2 instigated) or patient) adj2 engagement).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
- 34 patient participation.mp. or Client Participation/
- 35 Primary Health Care/
- 36 Family Medicine/
- 37 Health Care Services/
- 38 partnership practice.mp.
- 39 Private Practice/
- 40 (((((((medical adj2 office) or family) adj2 practice*) or general) adj2 practice*) or primary) adj2 care) or community) adj2 (service* or care)).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
- 41 Neoplasms/
- 42 (((cancer or neoplasm*) adj5 (diagnos* or detect*)) or neoplasms* or cancer* or carcinoma* or malignan* or lesion* or tumor?r*).mp. [mp=title, abstract, heading word, table of contents, key concepts, original title, tests & measures]
- 43 1 or 2
- 44 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34
- 45 35 or 36 or 37 or 38 or 39 or 40
- 46 41 or 42
- 47 43 and 44 and 45 and 46
- 48 limit 47 to english language

Appendix 3. Journal search — field and journal title

Field	Journal title
Diagnosis and decision making	<i>Diagnosis</i> <i>Medical Decision Making</i> <i>Judgement and Decision Making</i> <i>Journal of Risk and Uncertainty</i>
Primary health care	<i>Family Practice</i> <i>British Journal of General Practice</i> <i>Annals of Family Medicine</i> <i>European Journal of General Practice</i> <i>Primary Health Care Research and Development</i> <i>Journal of Primary Care and Community Health</i>
Cancer	<i>British Journal of Cancer</i> <i>International Journal of Cancer</i> <i>European Journal of Cancer</i> <i>European Journal of Cancer Care</i> <i>Cancer</i> <i>BMC Cancer</i> <i>Journal of Cancer</i>
Patient Safety	<i>BMJ Quality & Safety</i> <i>Patient Education and Counselling</i>

Appendix 4. Aim 1 inclusion and exclusion criteria

Articles were retained if they met all of the following inclusion criteria:

- 1) The article was about diagnosis
- 2) The article discussed patient involvement in diagnosis
- 3) The article was in a primary care setting
- 4) The article was about cancer

Articles were excluded if:

- 1) The article reported or its focus was cancer screening programmes
- 2) The article was concerned only with paediatric patients

Appendix 5. Aim 2 inclusion and exclusion criteria

Articles were retained if they met all of the following inclusion criteria:

- 1) The article provided suggestions for potential interventions/strategies to involve patients in diagnosis at the post-presentation stage
- 2) The article identified components considered to be key for potential interventions/strategies to involve patients in diagnosis at the post-presentation stage

Articles were excluded if:

- 1) The article only evaluated symptom perception before presentation to primary care
- 2) The article only evaluated help-seeking behaviour before presentation to primary care
- 3) The focus of the article was on epidemiology (for example, patterns, causes, and effects for identifying risk factors and targets for prevention)
- 4) The purpose of the article was to identify which symptoms are most common, or algorithms for diagnosis
- 5) The article involved the monitoring or surveillance of high-risk and predisposed patients identified as being at increased risk of cancer
- 6) The article was about metastasis in patients who already had cancer
- 7) The article was a case study
- 8) The article was about the relatives of cancer patients
- 9) The article included no elaboration or detailed examples and suggestions for patient involvement