

Accepted Manuscript

# *British Journal of General Practice*

## Measuring the complexity of general practice consultations: development and validation of a complexity measure

Salisbury, Chris; Lay-Flurrie, Sarah; Bankhead, Clare; Fuller, Alice;  
Murphy, Mairead; Caddick, Barbara; Ordóñez-Mena, Jose; Holt, Tim;  
Nicholson, Brian; Perera, Rafael; Hobbs, FD Richard

DOI: <https://doi.org/10.3399/BJGP.2020.0486>

To access the most recent version of this article, please click the DOI URL in the line above.

Received 21 May 2020

Revised 05 November 2020

Accepted 12 November 2020

© 2020 The Author(s). This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>). Published by British Journal of General Practice. For editorial process and policies, see: <https://bjgp.org/authors/bjgp-editorial-process-and-policies>

When citing this article please include the DOI provided above.

### **Author Accepted Manuscript**

This is an 'author accepted manuscript': a manuscript that has been accepted for publication in British Journal of General Practice, but which has not yet undergone subediting, typesetting, or correction. Errors discovered and corrected during this process may materially alter the content of this manuscript, and the latest published version (the Version of Record) should be used in preference to any preceding versions

# Measuring the complexity of general practice consultations: development and validation of a complexity measure

Chris Salisbury MD FRCGP<sup>1</sup>, Sarah Lay-Flurrie DPhil<sup>2</sup>, Clare R Bankhead MSc DPhil<sup>2</sup>, Alice Fuller BSc<sup>2</sup>, Mairead Murphy PhD<sup>1</sup>, Barbara Caddick PhD<sup>1</sup>, José M Ordóñez-Mena<sup>2,3</sup>, Tim Holt PhD FRCGP<sup>2</sup>, Brian D Nicholson DPhil MRCGP<sup>2</sup>, Rafael Perera DPhil<sup>2</sup>, FD Richard Hobbs FMedSci<sup>2</sup>

<sup>1</sup> Centre for Academic Primary Care, NIHR School for Primary Care Research, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

<sup>2</sup> Nuffield Department of Primary Care Health Sciences, Medical Sciences Division, University of Oxford, Oxford, UK

<sup>3</sup> NIHR Oxford Biomedical Research Centre, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

## **Correspondence to:**

Professor Chris Salisbury, Centre for Academic Primary Care, NIHR School for Primary Care Research, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, BS8 2PS UK

Email [c.salisbury@bristol.ac.uk](mailto:c.salisbury@bristol.ac.uk)

Tel 0117 3314530

Twitter: @prof\_tweet; @OxPrimaryCare; @capcbristol

Word count 2769

## Abstract

**Background:** The complexity of general practice consultations may be increasing and vary in different settings. Testing these hypotheses requires a measure of complexity.

**Aim:** To develop a valid measure of general practice consultation complexity applicable to routine medical records.

**Design:** Delphi study to select potential indicators of complexity followed by cross-sectional study to develop and validate a complexity measure.

**Setting:** English general practices.

**Method:** An online Delphi study over two rounds involved 32 general practitioners to identify potential indicators of consultation complexity. The cross-sectional study used an age-sex stratified random sample of 173,130 patients and 725,616 general practice face-to-face consultations from 2013/14 in the Clinical Practice Research Datalink. We explored independent relationships between each indicator and consultation duration using mixed effects regression models, and revalidated findings using data from 2017/18. We assessed the proportion of complex consultations in different age-sex groups.

**Results:** After two rounds, the Delphi panel endorsed 34 of 45 possible complexity indicators. In the cross-sectional study, after excluding factors because of low prevalence or confounding, 17 indicators were retained. Defining complexity as the presence of any of these factors, 308,370 consultations (42.5%) were complex. Mean duration of complex consultations was 10.49 minutes, compared to 9.64 minutes for non-complex consultations. The proportion of complex consultations was similar in men and women but increased with age.

**Conclusion:** Our consultation complexity measure has face and construct validity. It may be useful for research, management and policy, informing decisions about the range of resources needed in different practices.

**Keywords**

General practice; Delphi technique; Cross-sectional studies; Office visits; Risk adjustment

**How this fits in**

Increasing general practice workload due to rising consultation rates may be compounded by increasing complexity of consultations. Exploring these effects requires a valid and reliable measure of consultation complexity but there are no well-accepted measures. We have developed a suitable measure, starting with factors that general practitioners believe increase complexity and demonstrating those which are associated with longer consultations. Our complexity measure may be useful for research, management and policy, for example in allocating resources.

Accepted Manuscript – BJGP – BJGP 2020:0486

## **Introduction**

In the United Kingdom, general practitioners report increasing pressure from rising workload.<sup>1,2</sup> The number of consultations increased by 14% between 2007 and 2014, and the mean duration of face-to-face consultations increased by 7%.<sup>3</sup> Doctors' perceptions of increasing workload may reflect an increase in the complexity as well as number of consultations. This may be associated with an ageing population, increasing prevalence of multimorbidity and polypharmacy, transfer of activities from secondary to primary care, increasingly complex clinical guidelines, and growing policy expectations of what can be achieved within each consultation. The increasing delegation of routine tasks to other staff is also likely to increase the proportion of general practice consultations which are complex and intellectually demanding.<sup>1,4</sup>

To test this hypothesis, it is necessary to define and measure complexity within general practice consultations. A measure suitable for research and analysis needs to be applicable to routine electronic medical records, making it possible to explore changes in complexity over time and how consultation complexity varies in different practices, areas, and populations. A complexity measure would also be useful for resource allocation formulae, planning staffing needs, and as a case-mix variable within models to predict patient outcomes or the use of hospital and other services.

The aim of this study was to develop a valid and reliable measure of the complexity of general practice consultations that can be applied to routine medical records.

## **Methods**

We defined complex consultations as those which are more difficult to conduct, challenging, multi-faceted, intensive, or time-consuming than average. This study was conducted in two stages. First, we conducted a Delphi study to agree characteristics of consultations which were perceived by general practitioners to be complex. Second, we devised a valid and reliable measure using cross-sectional data from a large sample of routine general practice consultations, and re-validated the measure in a separate dataset of consultations from a different year.

### *Delphi study*

The research team created a long-list of variables that might increase the complexity of consultations based on previous literature,<sup>5-15</sup> their own experience, and informal discussion with general practice colleagues. We only included characteristics that were likely to be coded in routine medical records. We

did not include demographic factors such as age, sex or deprivation since the intention was to explore how the final complexity measure varied according to these characteristics.

Development and piloting of the Delphi study identified two conceptual issues. First, we found it necessary to distinguish between consultation complexity factors and patient complexity factors. We defined complex consultation factors as problems addressed within the consultation which made it complex. However, some patients have enduring characteristics which are likely to make most of their consultations complex irrespective of the problems presented – these were defined as complex patient factors. Second, we found that some practitioners felt that almost all their consultations were complex. Therefore, when designing the Delphi questionnaire, we asked clinicians whether each characteristic made a consultation ‘more complex than average’.

We asked colleagues from eight English universities to each recruit five clinically active general practitioners to participate in the Delphi study. These doctors were asked to complete an online questionnaire in two rounds. In the first round, they were presented with 14 consultation factors and 19 patient factors and asked to indicate whether or not each factor made consultations more complex than average, on a five point scale from 1 (no more complex than average) to 5 (very much more complex than average). Respondents to the first-round questionnaire could add comments about individual factors or suggest additional factors that we had not included.

Factors which received strong endorsement by the panel in the first round were accepted as markers of complexity. We considered scores of 3 to 5 (moderate to extreme complexity) to indicate endorsement of a characteristic and a score of 1 (no more complex than the average patient) to indicate rejection. Factors which more than 70% of participants endorsed, and fewer than 20% rejected, were accepted as markers of complexity. Factors which fewer than 40% of participants endorsed and more than 20% rejected were not accepted. All other factors were designated uncertain and were taken forward to a second round of voting. In some cases, the wording of items was revised before the second round to provide greater clarity in the light of respondents’ comments.

In the second round, participants were sent an individualised report which showed how their responses for each characteristic and overall compared with the median and inter-quartile range from other members of the panel. The report included a summary of comments from participants in round 1 about factors which had been designated uncertain. In round 2, participants were invited to vote again on the

uncertain factors and on new factors which had been proposed by participants. Factors were accepted or rejected using the same criteria as for round 1. Factors that remained uncertain were rejected.

#### *Creating and validating a complexity measure*

We created Read code sets for each of the patient and consultation complexity factors endorsed following the Delphi study. A researcher with extensive experience of coding general practice consultations (AF) created an initial code set for each characteristic. These code sets were checked independently by two academic general practitioners (BN and CS), with discrepancies resolved by discussion or involving a third academic general practitioner (RH). The final code sets are available at <https://doi.org/10.5287/bodleian:8gq7zbb8w>.

We assessed the prevalence of each characteristic based on an age-sex stratified sample from the Clinical Practice Research Datalink (CPRD) Gold database of non-temporary patients in England who were registered for any amount of time between 1/4/2013 and 31/3/2014 and had at least one face-to-face surgery consultation with a GP. We excluded from further consideration any characteristic which applied to less than 0.05% of consultations or patients, to simplify the measure. We used frequency data to specify factors which had been described qualitatively in the Delphi process. For example, 'frequent attender' was re-specified as patients with  $\geq 14$  GP consultations in the previous year, based on the 95<sup>th</sup> centile for number of consultations.

To assess construct validity, we explored the independent relationship between each complexity factor and consultation duration using mixed effects regression models of mean general practice consultation duration on patient and consultation complexity factors, with random effects for patient and practice. Consultation and patient factors were considered in separate models. Factors with a prevalence lower than 0.05% or which appeared to reduce the length of consultations were removed from the initial models. Remaining factors were removed in a backwards stepwise fashion using  $p < 0.05$  as the threshold. For a given consultation, the consultation factor applied if the topic was coded within the consultation and the patient factor applied if the consultation was with a patient with this factor.

We defined a complex consultation as one in which one or more complexity factor was present. We compared the mean duration of complex and non-complex consultations, and described the proportion of complex consultations by age-group.

As a further re-validation, we repeated the above analyses of construct validity in a separate dataset of patients from the CPRD comprising 67,829 patients who consulted at least once in 2017/18.

## Results

### *Delphi study*

Of 41 general practitioners sent details of the study, 32 agreed to participate and completed the first round of the survey. Participants included 10 (31%) men and 22 (69%) women with a mean of 11.2 years' (median 6; range <1 to 29 years) experience in general practice. The potential complexity factors in the first-round survey included 14 consultation factors and 19 patient factors. After the first round of the Delphi process, seven consultation factors were endorsed and none were rejected, while nine patient factors were endorsed and five were rejected. Seven consultation and five patient factors were uncertain and taken forward to round two, along with five new consultation factors and seven new patient factors suggested by panel participants.

In round two, 30 of the 32 round one participants (94%) completed the survey. A further 10 consultation factors and eight patient factors were endorsed, with the others being rejected or remaining uncertain (and therefore rejected). Therefore, after two rounds of the Delphi survey, 17 consultation factors and 17 patient factors were endorsed (Tables 1 and 2).

### *Creating and validating a complexity measure*

The demographic characteristics of the 173,130 patients included in the 2013/14 CPRD sample are shown in Supplementary Table 1. These patients had a total of 725,616 face-to-face consultations with a GP in 2013/14. Supplementary Tables 2 and 3 show the prevalence of consultation complexity factors and patient complexity factors respectively, along with the final wording used to define each factor.

Omission of factors coded in less than 0.05% of consultations or patients excluded two consultation factors (consultations about medically unexplained symptoms, resulting in urgent secondary care assessment) and two patient factors (paraplegia, medically unexplained symptoms within last year). Four further factors were excluded as consultation factors but included in the modelling as patient factors (palliative care, homelessness, domestic violence, safeguarding).

The results of the initial mixed effects regression models of consultation and patient factors against consultation duration for 2013/14 are shown in Table 3, with equivalent data for 2017/18 in

Supplementary Table 4. The final models, omitting variables with no significant relationship with consultation duration, include 17 factors (Table 4).

The re-validation analysis used data on consultations between 1/4/2017 and 31/3/2018 and included 234,447 consultations with 58,528 independent patients. In the final model, five factors were no longer significantly associated with consultation duration (Table 4). However, we decided to retain these factors in our complexity measure because they had all been endorsed by GPs in the Delphi study, and the coefficients for duration were all positive with confidence intervals that overlapped in the development and validation data sets. The higher p values in 2017/18 are likely to be due to the smaller total sample size. Defining complexity as the presence of any of these factors at the consultation, 308,370 consultations (42.5%) were defined as complex in 2013/14. The mean duration of complex consultations was 10.49 minutes, compared to 9.64 for non-complex consultations, with a difference of 0.85 (0.81 to 0.89) minutes. Equivalent analyses for the revalidation dataset in 2017/18 provided very similar results, with 41.6% (97,547/234,447) of consultations defined as complex. The mean duration of complex consultations in 2017/18 was 10.32 minutes, compared to 9.70 for non-complex consultations (difference 0.62 (0.55 to 0.69) minutes).

The proportion of complex consultations was strongly associated with increasing age, and was slightly higher in men than in women except in patients aged over 85 years (Figure 1).

## **Discussion**

### *Summary*

In this paper we have defined, created and applied a measure of the complexity of general practice consultations which can be applied to routine electronic medical records. This measure was constructed using characteristics of patients and problems selected by a consensus process involving experienced general practitioners, demonstrating face validity. The measure has been validated by showing that each of the characteristics in the final selection, and the overall complexity measure, are associated with consultation duration in two independent samples of consultations.

### *Comparison with existing literature*

Our research builds on previous research. Two studies<sup>4,7</sup> and an online survey<sup>2</sup> have asked primary care clinicians to record the complexity of their consultations subjectively, for example using a five-point scale from very simple to very complex, while another study quantified the number and range of

problems discussed within consultations.<sup>16</sup> Three studies have asked general practitioners about features that make patients complex, and we build on this by considering aspects of consultations as well as patients.<sup>12,14,15,17</sup> A few previous authors have devised case-mix measures applicable to primary care, but these have either not taken account of clinicians' perceptions of the complexity of different factors<sup>18-21</sup> or not been designed for analysis of routine medical records.<sup>13 22</sup>

There is some overlap between measures of complexity and case-mix measures such as Adjusted Clinical Groups,<sup>23</sup> Rx-Risk<sup>24</sup> and the Charlson score<sup>25</sup> which have been designed to predict health outcomes, resource utilisation or mortality. These case-mix measures are based on combinations of diagnostic information, medication data and/or demographic factors but do not account for social, behavioural or other psychological factors<sup>11</sup> which often create the greatest demands on general practitioners within consultations<sup>12,15,17,26</sup> and are captured by our complexity measure.

#### *Strengths and limitations*

Our study has several limitations. The concept of complexity in consultations is nebulous, and although widely recognised by clinicians it is hard to define.<sup>13,26</sup> Our definition of complexity encompasses intellectual, emotional and workload demands, but other definitions of complexity would lead to different measurement tools. The choice of complexity factors was based on the experience of the research team and the literature, with additional factors suggested by the general practitioners in the Delphi panel, but other factors could have been considered. Some factors may add complexity to consultations but are not coded within electronic medical records. We dropped some factors (such as medically unexplained symptoms) which almost certainly generate complexity within consultations but are rarely coded, so inclusion would add little to the measure when used for analysis at a population level. Two variables (patients with dementia or housebound) had a significant negative association with consultation duration. In post-hoc analysis we found that these characteristics were associated with more consulting time over a whole year, resulting from a higher number of consultations which are shorter than average.

The development of the complexity measure was conducted in England, and factors which cause consultation complexity may be different in other countries (for example insurance status, in the United States).<sup>13,14</sup> The complexity measure we developed was based on a sample of consultations which is six years old. This was deliberate, to create a baseline against which to assess changes in complexity over

time in a subsequent paper. However, we revalidated the findings in a more recent dataset (2017/18) and this analysis largely confirmed our findings.

We recognise that mean duration of consultations is not a gold standard for complexity, since the length of a consultation is only partly related to complexity and not all complex consultations are lengthy. However, it was the best and simplest (while imperfect) proxy available within routine medical records. The purpose of the cross-sectional analysis was not to derive a model to predict consultation duration, but to provide evidence for the construct validity of our complexity measure by showing a positive association with a variable (duration) that we hypothesised would be related to it. The analysis fulfilled our aims by confirming (a) that each of the included complexity factors was independently associated with longer consultations, (b) that a measure defined as the presence of one or more of these factors was discriminating, with complex consultations being on average 9% longer than non-complex consultations and (c) that these findings were robust when repeated in a different data sample. Although our complexity measure is useful as a binary 'complex/non-complex' variable, we do not propose combining the factors to create a cumulative score (see statistical note in appendix).

Our measure is reliable in that it is based on objective analysis of medical records and defined code sets for complexity factors, unlike measures which require subjective judgements.<sup>4,7,13,22</sup> Basing the measure on the views of practicing GPs and assessing the relationship with consultation duration provides evidence of face and construct validity respectively.

Further validation exercises could explore the relationship between our complexity measure and other variables, such as practitioners' self-assessment of the complexity of a sample of consultations. Future research should also explore the relationship between complexity and risk prediction models for health care utilisation. We anticipate some, but not complete, overlap.<sup>14</sup> It is likely that different tools will be best at predicting different outcomes and measures should be used in combination to understand population healthcare needs.<sup>27</sup>

#### *Implications for research and practice*

This paper describes a valid and reliable measure of the complexity of general practitioners' consultations. In future research we plan to explore the complexity of consultations in different settings and populations, and how complexity has changed over time. This may be relevant to the development of resource allocation formulae. The current UK formula for allocating payments to primary care takes account of the number of expected consultations based on characteristics of the practice population,

but not the complexity of those consultations.<sup>28</sup> Practices which have a high proportion of complex consultations may need a different mix of staff from practices with few complex consultations. There is growing interest in creating population health management systems by linking health and social care datasets to understand current and future health and care needs.<sup>29</sup> Use of a complexity measure may support this aim, providing greater nuance and understanding by taking account of the different workforce, workload and resource implications of consultations with different levels of complexity.

### **Funding**

This research was supported by the NIHR School for Primary Care Research grant number 243 and the Nuffield Department of Primary Care Health Sciences, University of Oxford.

### **Ethical approval**

The protocol was approved by the Independent Scientific Advisory Committee (ISAC) of the MHRA (ISAC protocol number 15\_120R; available from the authors on request). Ethics approval for observational research using the CPRD with approval from ISAC was granted by a National Research Ethics Service committee (Trent MultiResearch Ethics Committee, REC reference number 05/MRE04/87).

### **Competing interests**

None

### **Contributions**

CS and FDRH conceived the research and drafted the protocol with CB, which all then contributed to. CS drafted the grant application with CS and FDRH as co-principal investigators. MM designed and analysed the Delphi study which was co-ordinated by BC. AF and SLF developed initial code lists and conducted data management. SLF and JMOM conducted the statistical analysis, with advice from RP. TH and BN contributed to clinical coding and study design. CB managed the overall project. CS wrote the first draft of the paper and all authors provided critique of the output for important intellectual content. CS is guarantor and corresponding author.

## **Acknowledgements**

We would like to thank Chelu Mfalila for her help with creating coding sets. CS acknowledges part funding from National Institute for Health Research (NIHR) Applied Research Collaboration (ARC) West at University Hospitals Bristol NHS Foundation Trust and from Bristol, North Somerset and South Gloucestershire Clinical Commissioning Group. Authors acknowledge part-funding from the NIHR ARC Oxford and Thames Valley (FDRH, CB, SLF, JMOM, RP), NIHR Oxford Biomedical Research Centre (FDRH, CB, JMOM, RP), NIHR Oxford Medtech and the In-Vitro Diagnostics Co-operative (FDRH, RP), NIHR School for Primary Care Research (FDRH) and the Oxford Martin School (RP). BDN acknowledges funding from the NIHR. This study is based in part on data from the Clinical Practice Research Datalink obtained under licence from the UK Medicines and Healthcare products Regulatory Agency. The data is provided by patients and collected by the NHS as part of their care and support. The interpretation and conclusions contained in this study are those of the author/s alone. CS and FDRH are National Institute for Health Research (NIHR) Senior Investigators (FDRH is emeritus). The views expressed in this article are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

## **Competing interests**

None


Accepted Manuscript – BJGP – BJCPR 2020.2485

## References

1. Pressures in general practice: BMA; [Available from: <https://www.bma.org.uk/advice-and-support/nhs-delivery-and-workforce/pressures/pressures-in-general-practice> accessed 10/08/2020.
2. Anderson A. Pulse 2019 review: GP workload laid bare: Pulse; 2019 [Available from: <http://www.pulsetoday.co.uk/pulse-2019-review/pulse-2019-review-gp-workload-laid-bare/20039883.article> accessed 10/08/2020.
3. Hobbs FDR, Bankhead C, Mukhtar T, et al. Clinical workload in UK primary care: a retrospective analysis of 100 million consultations in England, 2007-14. *Lancet* 2016;387(10035):2323-30. doi: 10.1016/S0140-6736(16)00620-6
4. Gemmell I, Campbell S, Hann M, et al. Assessing workload in general practice in England before and after the introduction of the pay-for-performance contract. *J Adv Nurs* 2009;65(3):509-15. doi: 10.1111/j.1365-2648.2008.04902.x
5. Salisbury C, Procter S, Stewart K, et al. The content of general practice consultations: cross-sectional study based on video recordings. *Br J Gen Pract* 2013;63(616):e751-9. doi: 10.3399/bjgp13X674431
6. Baird B, Charles A, Honeyman M, et al. Understanding pressures in general practice London 2016 [Available from: [https://www.kingsfund.org.uk/sites/default/files/field/field\\_publication\\_file/Understanding-GP-pressures-Kings-Fund-May-2016.pdf](https://www.kingsfund.org.uk/sites/default/files/field/field_publication_file/Understanding-GP-pressures-Kings-Fund-May-2016.pdf) accessed 21/05/2020.
7. Mercer SW, Fitzpatrick B, Gourlay G, et al. More time for complex consultations in a high-deprivation practice is associated with increased patient enablement. *Br J Gen Pract* 2007;57(545):960-66. doi: 10.3399/096016407782604910
8. Vedsted P, Christensen MB. Frequent attenders in general practice care: a literature review with special reference to methodological considerations. *Public Health* 2005;119(2):118-37.
9. Deveugele M, Derese A, van den Brink-Muinen A, et al. Consultation length in general practice: cross sectional study in six European countries. *BMJ* 2002;325(7362):472.
10. The 2022 GP: Compendium of evidence London: Royal College of General Practitioners; 2013 [Available from: <https://www.rcgp.org.uk/~media/Files/Policy/A-Z-policy/The-2022-GP-Compendium-of-Evidence.ashx> accessed 21/05/2020.
11. Rosen AK, Reid R, Broemeling AM, et al. Applying a risk-adjustment framework to primary care: can we improve on existing measures? *Ann Fam Med* 2003;1(1):44-51.
12. Grant RW, Ashburner JM, Hong CS, et al. Defining Patient Complexity From the Primary Care Physician's Perspective: A Cohort Study. *Ann Intern Med* 2011;155(12):797-804. doi: 10.7326/0003-4819-155-12-201112200-00001
13. Peek CJ, Baird MA, Coleman E. Primary care for patient complexity, not only disease. *Fam Syst Health* 2009;27(4):287-302. doi: 10.1037/a0018048
14. Hong CS, Atlas SJ, Ashburner JM, et al. Evaluating a Model to Predict Primary Care Physician-Defined Complexity in a Large Academic Primary Care Practice-Based Research Network. *J Gen Intern Med* 2015;30(12):1741-47. doi: 10.1007/s11606-015-3357-8
15. Loeb DF, Binswanger IA, Candrian C, et al. Primary Care Physician Insights Into a Typology of the Complex Patient in Primary Care. *Ann Fam Med* 2015;13(5):451-55. doi: 10.1370/afm.1840
16. Procter S, Stewart K, Reeves D, et al. Complex consultations in primary care: a tool for assessing the range of health problems and issues addressed in general practice consultations. *BMC family practice* 2014;15:105. doi: 10.1186/1471-2296-15-105

17. Webster F, Rice K, Bhattacharyya O, et al. The mismeasurement of complexity: provider narratives of patients with complex needs in primary care settings. *Int J Equity Health* 2019;18(1):107. doi: 10.1186/s12939-019-1010-6
18. Tonelli M, Wiebe N, Manns BJ, et al. Comparison of the Complexity of Patients Seen by Different Medical Subspecialists in a Universal Health Care System. *JAMA Netw Open* 2018;1(7):e184852-e52. doi: 10.1001/jamanetworkopen.2018.4852
19. Halter M, Joly L, de Lusignan S, et al. Capturing complexity in clinician case-mix: classification system development using GP and physician associate data. *BJGP Open* 2018;2(1):bjgpopen18X101277-bjgpopen18X77. doi: 10.3399/bjgpopen18X101277
20. Katerndahl D, Wood R, Jaén CR. Family Medicine Outpatient Encounters are More Complex Than Those of Cardiology and Psychiatry. *J Am Board Fam Med* 2011;24(1):6. doi: 10.3122/jabfm.2011.01.100057
21. Katerndahl DA, Wood R, Jaen CR. A method for estimating relative complexity of ambulatory care. *Ann Fam Med* 2010;8(4):341-7. doi: 10.1370/afm.1157
22. Pratt R, Hibberd C, Cameron IM, et al. The Patient Centered Assessment Method (PCAM): integrating the social dimensions of health into primary care. *J Comorb* 2015;5:110-19. doi: 10.15256/joc.2015.5.35
23. Weiner JP, Starfield BH, Steinwachs DM, et al. Development and application of a population-oriented measure of ambulatory care case-mix. *Med Care* 1991;29(5):452-72.
24. Sloan KL, Sales AE, Liu CF, et al. Construction and characteristics of the RxRisk-V: a VA-adapted pharmacy-based case-mix instrument. *Med Care* 2003;41(6):761-74. doi: 10.1097/01.Mlr.0000064641.84967.B7
25. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40(5):373-83.
26. Safford MM, Allison JJ, Kiefe CI. Patient complexity: more than comorbidity. the vector model of complexity. *J Gen Intern Med* 2007;22 Suppl 3(Suppl 3):382-90. doi: 10.1007/s11606-007-0307-0
27. Safford MM. The Complexity of Complex Patients. *J Gen Intern Med* 2015;30(12):1724-25. doi: 10.1007/s11606-015-3472-6
28. DH Financial Planning and Allocations Division. Resource Allocation: Weighted Capitation Formula (Seventh Edition). London, 2011.
29. Population Health and the Population Health Management Programme: NHS England; [Available from: <https://www.england.nhs.uk/integratedcare/building-blocks/phm/> accessed 23/04/2020.
30. Payne RA, Mendonca SC, Elliott MN, et al. Development and validation of the Cambridge Multimorbidity Score. *CMAJ* 2020;192(5):E107-E14. doi: 10.1503/cmaj.190757

Figure 1 Percentage of consultations which are complex, by age and sex.



Accepted Manuscript – BJG

2020.0486

Table 1 Endorsement of consultation complexity factors in two rounds of Delphi study

Consultation Variables: Final Wording	Round 1 result	Median Score	Round 2 result	Median Score	Final inclusion
<b>Factors accepted or rejected in round 1</b>					
Patient presents with problem of being homeless	✓	3.0			Included
Patient presents with problem which raises child protection or adult safeguarding concerns	✓	4.0			Included
Patient presents with problem which raises concerns about domestic violence	✓	4.0			Included
Consultation about learning disability/autism	✓	3.0			Included
Discussion about end of life issues in current consultation	✓	3.0			Included
Consultation about Mental Health Problems	✓	4.0			Included
Multiple diagnoses or problems being managed in the current consultation	✓	3.0			Included
<b>Factors carried forward to round 2</b>					
Consultation about dementia	?	3.0	✓	3.0	Included
Discussion about problematic drug or alcohol use in current consultation	?	3.0	✓	3.5	Included
Several preventive health care and routine monitoring tasks being conducted in same consultation	?	3.0	✓	3.0	Included
Procedures or minor surgery carried out in the current consultation	?	2.0	?	2.0	Rejected
Needing to prescribe many drugs in the current consultation	?	3.0	✓	3.0	Included
First GP consultation following a diagnosis of cancer	?	3.0	✓	4.0	Included
First GP consultation following a diagnosis of diabetes	?	3.0	✓	3.0	Included
<b>Factors suggested by participants and included in round 2</b>					
Medically unexplained symptoms raised in consultation			✓	4.0	Included
Consultation results in outpatient referral			✗	2.0	Rejected
Consultation results in an emergency hospital admission			✓	4.0	Included
Consultation results in urgent secondary care assessment (e.g. crisis team/A&E)			✓	4.0	Included
Consultation about chronic pain management			✓	3.0	Included

Key: ✓ endorsement ? uncertain ✗ rejection

Table 2 Endorsement of patient complexity factors in two rounds of Delphi study

Patient Variables: Final Wording	Round 1 result	Median Score	Round 2 result	Median Score	Final inclusion
<b>Factors accepted or rejected in round 1</b>					
Homelessness (noted in the previous year)	✓	3.0			Included
Child protection/adult safeguarding (until resolved code)	✓	4.0			Included
Domestic Violence (recorded in the previous year)	✓	3.0			Included
Interpreter needed/no English (noted in last three years)	✓	3.0			Included
Learning disability/autism (noted ever)	✓	3.0			Included
Dementia (noted ever)	✓	3.0			Included
Receiving Palliative Care (noted ever)	✓	3.0			Included
Substance misuse / alcoholism (noted in the previous year)	✓	3.5			Included
Severe mental illness (in the previous three years)	✓	4.0			Included
Wheelchair use (in the previous three years)	✗	2.0			Rejected
Recent Outpatient referral	✗	2.0			Rejected
Patient Currently on Warfarin	✗	2.0			Rejected
Cancer (noted ever)	✗	2.0			Rejected
Diabetes (noted ever)	✗	2.0			Rejected
<b>Factors carried forward to round 2</b>					
Patient has 3 or more major long term chronic conditions	?	2.0	✓	3.0	Included
Deafness (noted ever)	?	2.0	?	2.0	Rejected
Paraplegic (noted ever)	?	2.5	✓	3.0	Included
Blind or partially sighted (noted ever)	?	2.0	?	2.0	Rejected
Patient on immunosuppressant medication (currently)	?	2.0	?	3.0	Rejected
<b>Factors suggested by participants and included in round 2</b>					
Housebound or nursing home patient			✓	4.0	Included
Personality disorder or disruptive behaviour (noted ever)			✓	4.0	Included
Diagnostic code for 'Medically Unexplained Symptoms' entered in last year			✓	3.0	Included
Patient is morbidly obese (BMI > 40)			?	3.0	Rejected

Frequent attender (high number of GP consultations in the last year).	✓	3.0	Included
Chronic pain recorded as a code in the last year	✓	3.0	Included
Polypharmacy (high number of drugs prescribed in the last 2 months).	✓	3.5	Included

Key: ✓ endorsement ? uncertain ✗ rejection

Accepted Manuscript – BJGP – BJGP.2020.0486

Table 3 Initial mixed effects regression of consultation and patient factors against consultation duration. Random effects for patient and practice.<sup>a</sup> Data from 2013/14.

	Univariable <sup>c</sup>			Multivariable		
	Change in mins	95% CI	p-value	Change in mins	95% CI	p-value
<b>Consultation factors<sup>a</sup></b>						
Mean duration	NA	NA	NA	9.78	9.56 to 10.00	<0.001
About drug/ alcohol use	4.49	3.99 to 4.98	<0.001	4.19	3.68 to 4.69	<0.001
About chronic pain	1.48	1.37 to 1.6	<0.001	0.98	0.86 to 1.1	<0.001
About dementia	1.45	0.92 to 1.98	<0.001	1.42	0.90 to 1.94	<0.001
Results in emergency hospital admission	7.81	7.12 to 8.5	<0.001	7.76	7.09 to 8.43	<0.001
About learning disability/ autism	4.54	3.86 to 5.22	<0.001	3.84	3.17 to 4.52	<0.001
About mental health problems	4.06	3.92 to 4.21	<0.001	3.85	3.70 to 3.99	<0.001
Two or more diagnoses from unique Read chapters	2.99	2.87 to 3.12	<0.001	2.54	2.42 to 2.67	<0.001
Three or more unique substance prescribed	1.93	1.87 to 1.99	<0.001	1.73	1.67 to 1.8	<0.001
Two or more preventive/ routine tasks carried out	3.94	3.75 to 4.13	<0.001	3.73	3.55 to 3.92	<0.001
First consultation after cancer diagnosis	0.59	0.04 to 1.14	0.037	0.43	-0.11 to 0.97	0.118
First consultation after diabetes diagnosis	3.59	3.00 to 4.17	<0.001	3.05	2.48 to 3.63	<0.001
<b>Patient factors<sup>a</sup></b>						
Mean duration of consultation	NA	NA	NA	10.02	9.79 to 10.25	<0.001
Drug/ alcohol abuse in previous year	2.03	1.58 to 2.48	<0.001	1.89	1.44 to 2.33	<0.001
Chronic pain in previous year	0.87	0.72 to 1.02	<0.001	0.73	0.58 to 0.89	<0.001
Dementia (ever)	-0.78	-1.19 to -0.38	<0.001	NA <sup>b</sup>	NA <sup>b</sup>	NA <sup>b</sup>
Domestic violence in last year	1.46	0.40 to 2.52	0.007	1.43	0.37 to 2.49	0.008
Frequent attender (14+ consultations in last year)	0.35	0.19 to 0.52	<0.001	0.01	-0.17 to 0.19	0.902
Homelessness in previous year	1.64	0.67 to 2.61	<0.001	1.36	0.39 to 2.33	0.006
Housebound or nursing home in previous 3 years	-3.72	-4.37 to -3.06	<0.001	NA <sup>b</sup>	NA <sup>b</sup>	NA <sup>b</sup>
No English noted in last 3 years	1.02	0.29 to 1.76	0.006	0.98	0.25 to 1.72	0.009
Learning disability/ autism (ever)	0.10	-0.17 to 0.36	0.481	0.06	-0.21 to 0.33	0.654
Severe mental illness in previous 3 years	0.18	-0.36 to 0.72	0.506	-0.10	-0.64 to 0.44	0.727
3 or more long term conditions <sup>d</sup>	0.45	0.36 to 0.54	<0.001	0.32	0.21 to 0.43	<0.001
Receiving palliative care (ever)	-0.58	-1.22 to 0.05	0.07	NA <sup>b</sup>	NA <sup>b</sup>	NA <sup>b</sup>
Personality/ disruptive disorder (ever)	0.75	0.37 to 1.13	<0.001	0.51	0.12 to 0.89	0.01
Polypharmacy (9 or more unique substances prescribed in previous 3 months)	0.40	0.25 to 0.56	<0.001	-0.07	-0.25 to 0.11	0.447
Child protection/ safeguarding in previous 3 years	-0.33	-0.69 to 0.04	0.079	NA <sup>b</sup>	NA <sup>b</sup>	NA <sup>b</sup>

<sup>a</sup> based on separate regressions for consultation and patient factors

<sup>b</sup> Factors which had a negative relationship with consultation duration were excluded

<sup>c</sup> mixed effect model with random intercepts for practice and patient, and a fixed effect for each patient or consultation factor at a time.

<sup>d</sup> Based on conditions included in the Cambridge Multimorbidity Score.<sup>30</sup>

Table 4 Final mixed effects regression models of patient and consultation factors against consultation duration in the development and validation data sets.<sup>a</sup> Random effects for patient and practice.

	2013/14 (development N = 725,616 consultations)			2017/18 (validation N = 234,447 consultations)		
	Change in mins	95% CI	p	Change in mins	95% CI	p
<b>Consultation factors<sup>a</sup></b>						
Mean duration of consultation	9.78	9.56 to 10	<0.001	9.81	9.38 to 10.24	<0.001
About drug/ alcohol use	4.19	3.68 to 4.69	<0.001	3.73	2.54 to 4.92	<0.001
About chronic pain	0.98	0.86 to 1.1	<0.001	1.00	0.79 to 1.21	<0.001
About dementia <sup>b</sup>	1.42	0.9 to 1.94	<0.001	NA	NA	NA
Results in emergency hospital admission	7.76	7.09 to 8.43	<0.001	4.69	3.47 to 5.92	<0.001
About learning disability/ autism	3.84	3.17 to 4.52	<0.001	3.05	1.98 to 4.12	<0.001
About mental health problems	3.85	3.7 to 3.99	<0.001	3.83	3.58 to 4.08	<0.001
Two or more diagnoses from unique Read chapters recorded	2.54	2.42 to 2.67	<0.001	2.86	2.61 to 3.12	<0.001
Three or more unique substance prescribed	1.73	1.67 to 1.8	<0.001	1.82	1.69 to 1.94	<0.001
Two or more preventive/ routine tasks carried out	3.73	3.55 to 3.92	<0.001	4.81	4.44 to 5.19	<0.001
First consultation after diabetes diagnosis	3.05	2.48 to 3.62	<0.001	2.39	1.6 to 3.19	<0.001
<b>Patient factors<sup>a</sup></b>						
Mean duration of consultation	10.02	9.8 to 10.25	<0.001	10.03	9.6 to 10.46	<0.001
Drug/ alcohol abuse in previous year <sup>b</sup>	1.89	1.44 to 2.33	<0.001	NA	NA	NA
Chronic pain in previous year	0.72	0.57 to 0.87	<0.001	0.75	0.51 to 1	<0.001
Domestic violence in last year	1.43	0.37 to 2.49	0.008	2.37	0.99 to 3.74	<0.001
Homelessness in previous year <sup>b</sup>	1.36	0.39 to 2.33	0.006	NA	NA	NA
No English noted in last 3 years	0.98	0.25 to 1.72	0.009	1.01	0.12 to 1.91	0.026
3 or more long term conditions <sup>b</sup>	0.30	0.21 to 0.4	<0.001	NA	NA	NA
Personality/ disruptive disorder (ever) <sup>b</sup>	0.51	0.12 to 0.89	0.01	NA	NA	NA

<sup>a</sup> based on separate regressions for consultation and patient factors

<sup>b</sup> These variables were retained in our complexity measure despite low prevalence or lack of significant effect in the 2017/18 model. See text of article.